

Product Name: ALBAUGH ADJUST 38 COTTON GROWTH REGULATOR
APVMA Approval No.: 56331/141418



Label Name:	ALBAUGH ADJUST 38 COTTON GROWTH REGULATOR
Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	ACTIVE CONSTITUENT: 38 g/L MEPIQUAT PRESENT AS THE CHLORIDE
Mode of Action:	
Statement of Claims:	For the management of fruiting and excessive vegetative growth in cotton
Net Contents:	NET CONTENTS 1000L 110L 200L 20L
Restrains:	RESTRAINTS: DO NOT apply to crops that have restricted growth due to drought stress, waterlogging, nutrient deficiency, poor soil structure or any other cause as this can result in yield decreases. DO NOT apply if adequate insect / mite control and supply of water / nutrients later in the crop is in doubt. DO NOT apply 'Albaugh Adjust' if rainfall is expected within 8 hours of application as product effectiveness may be reduced. DO NOT use on new varieties not available commercially in Australia without first checking with the manufacturer. DO NOT apply to cotton less than 50cm high if using the high rate single application spray.

Directions for Use:	This section contains file attachment.
---------------------	--

Other Limitations:	
--------------------	--

Withholding Periods:	<p>WITHHOLDING PERIOD:</p> <p>DO NOT APPLY LATER THAN 4 WEEKS BEFORE HARVEST</p>
----------------------	--

Trade Advice:	
---------------	--

General Instructions:	<p>GENERAL INSTRUCTIONS: 'Albaugh Adjust' is a foliar applied plant growth regulator, which alters the cotton plant growth pattern and behaviour. One or more of the following effects may be produced in cotton: darker leaf colour, reduction of internodal length, more open canopy, better boll retention, earlier maturity. Some of these effects may favourably influence the yield potential of the cotton plant.</p> <p>APPLICATION: 'Albaugh Adjust' may be applied using a split application technique, i.e. 3 separate low rate sprays applied at strategic times during the growing period. It can also be applied as a single application at the 1st flower stage. Information on rates and timing is given in the "Directions for Use Table". Use a total spray volume to ensure adequate cover of foliage. Add an agricultural wetting agent to improve foliar coverage. If 'Albaugh Adjust' is mixed with another product, the addition may not be necessary as one may be present.</p> <p>MIXING: 'Albaugh Adjust' mixes readily with water. Add the required amount to the partly filled tank, agitate and fill to the desired level.</p> <p>LODGING: Cotton treated with 'Albaugh Adjust' that has set bolls packed tightly together against the stem may be more susceptible to lodging should adverse conditions occur.</p> <p>COMPATIBILITY: Do not tank mix with foliar fertilisers. Although 'Albaugh Adjust' is compatible with insecticides normally applied to cotton, always check compatibility in a small container prior to use.</p>
-----------------------	---

Resistance Warning:	
---------------------	--

Precautions:	
--------------	--

Protections:	<p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT: DO NOT contaminate streams, rivers or waterways with the chemical or used containers</p>
--------------	--

Storage and Disposal:	<p>STORAGE AND DISPOSAL: Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.</p> <p>Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or a designated collection point.</p> <p>If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.</p> <p>Refillable containers: empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.</p>
-----------------------	---

Safety Directions:	<p>SAFETY DIRECTIONS: Avoid Contact with eyes and skin. Wash hands after use.</p>
--------------------	--

First Aid Instructions:	<p>FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131126</p>
-------------------------	--

First Aid Warnings:	
---------------------	--

DIRECTIONS FOR USE

Crop	Purpose	State	Application Rate	WHP	Critical Comments
Cotton	To regulate Fruiting and vegetive growth	QLD and NSW only	Split Application Rate:	4 weeks	Split Applications: Apply no more than 1.5L in total, spread over 2=3 application times as suggested below.
			250 to 600ml /ha		Apply the 1 st spray at early squaring-squares the size of match heads.
			250 to 600ml /ha		Apply the 2 nd spray at approximately first flow.
			250 to 600ml /ha		Apply the 3 rd spray at approximately 14 days later then 1 st flower.
	Control of excessive growth		Single Application Rate:		Single Application:
			750 to 1L /ha		Apply at 1 st flower/ use the rate where crop growth is excessive.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.