

Product Name: Albaugh Polca 125 SC Fungicide
APVMA Approval No: 94627/143009



Label Name:	Albaugh Polca 125 SC Fungicide
Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	ACTIVE CONSTITUENT 125 g/L EPOXICONAZOLE
Mode of Action:	GROUP 3 FUNGICIDE
Statement of Claims:	For the control of leaf rust, stripe rust and septoria nodorum blotch of wheat and leaf rust of barley, as specified in the DIRECTIONS FOR USE table
Net Contents:	5 L - 20 L
Restrains:	This section contains file attachment.
Directions for Use:	This section contains file attachment.
Other Limitations:	
Withholding Periods:	WITHHOLDING PERIOD(S) DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION. DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 6 WEEKS AFTER APPLICATION.

Trade Advice:	<p>CAUTION</p> <p>Export of treated cereals</p> <p>Growers should note that Maximum Residue Limits (MRLs) or import tolerances do not exist in all markets for cereals treated with Albaugh Polca 125 SC Fungicide. Additionally, some export markets have established MRLs different to those in Australia. If you are growing cereals for export, please check with ALBAUGH ASIA PACIFIC LIMITED or your grain exporter for the latest information on MRLs and import tolerances BEFORE using Albaugh Polca 125 SC Fungicide.</p> <p>LIVESTOCK DESTINED FOR EXPORT MARKETS</p> <p>The grazing withholding period only applies to stock slaughtered for the domestic market. Some export markets apply different standards. To meet these standards, ensure that in addition to complying with the grazing withholding period, the Export Slaughter Interval is observed before stock are sold or slaughtered.</p> <p>EXPORT SLAUGHTER INTERVAL (ESI): AFTER OBSERVING THE PERIOD FOR GRAZING OR CUTTING FOR STOCKFOOD, LIVESTOCK THAT HAVE BEEN GRAZED ON OR FED TREATED CROPS SHOULD BE PLACED ON CLEAN FEED FOR 3 DAYS PRIOR TO SLAUGHTER.</p>
---------------	---

General Instructions:	<p>Albaugh Polca 125 SC Fungicide is a broad-spectrum contact and systemic fungicide. It is taken up via the stem and foliage and is translocated upwards and outwards to provide protection to new growth. Thorough coverage of the crop is necessary for best results. Check the crop regularly for any symptoms of disease infection. Apply as soon as disease is present on the flag leaf or flag-2. Application beyond flag leaf emergence (ZGS39) may not provide any yield benefit, particularly under dry seasonal conditions.</p> <p>MIXING</p> <p>Albaugh Polca 125 SC Fungicide is a suspension concentrate formulation. Add the product to the half-filled spray tank while agitating. Continue to agitate while topping up the tank and during spraying.</p> <p>APPLICATION</p> <p>Ground Application</p> <p>Apply in a water volume of between 50 and 100L/ha, using flat fan nozzles operating at around 50 cm above the top of the crop. Use the higher water volume in crops with heavier canopies.</p> <p>Aerial Application</p> <p>Apply with suitable aircraft set up and operated to apply fungicides to cereal crops in a minimum water volume of 20L/ha.</p> <p>COMPATIBILITY</p> <p>For information on compatibility please contact your local reseller or ALBAUGH ASIA PACIFIC LIMITED representative.</p>
-----------------------	--

Resistance Warning:	<p>FUNGICIDE RESISTANCE WARNING</p> <p>GROUP 3 FUNGICIDE</p> <p>Albaugh Polca 125 SC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Albaugh Asia Pacific Limited accepts no liability for any losses that may result from the failure of this product to control resistant fungi.</p>
---------------------	--

Precautions:	<p>RE-ENTRY PERIOD</p> <p>DO NOT allow entry into treated areas until the spray has dried.</p>
--------------	---

Protections:	<p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT</p> <p>Highly toxic to aquatic organisms. Do NOT apply under weather conditions or from spraying equipment which could be expected to cause spray drift onto adjacent crops, croplands, livestock, pastures, wetlands, waterbodies or watercourses. Do NOT contaminate streams, rivers or watercourses with the chemical or used containers.</p>
--------------	--

Storage and Disposal:	<p>Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.</p> <p>Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.</p> <p>If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p>
-----------------------	--

Safety Directions:	Harmful if inhaled or swallowed. Will irritate the eyes, nose, throat and skin. Avoid contact with eyes and skin. Do not inhale vapour. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and half facepiece respirator. When using the prepared spray wear cotton overalls buttoned to the neck and wrist or equivalent clothing and a washable hat. If product on skin, immediately wash area with soap and water. Wash hands after use. After each days use, wash gloves and respirator and if rubber wash with detergent and warm water. Wash contaminated clothing.
--------------------	---

First Aid Instructions:	FIRST AID If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.
-------------------------	--

First Aid Warnings:	
---------------------	--

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/Spraydrift

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by aircraft unless the following requirements are met:

For maximum release heights above the target canopy of 3m or 25% of wingspan or 25% of rotor diameter whichever is the greatest, minimum distances between the application site and downwind sensitive areas (see 'Buffer zones' section of the following table titled 'Buffer zones for aircraft') are observed.

Buffer zones for aircraft

Application rate	Mandatory downwind buffer zones
	Aquatic areas
Up to a maximum label rate	30 metres

DIRECTIONS FOR USE

Crop	Disease	Rate per ha	Critical Comments
Wheat	Leaf rust (<i>Puccinia recondita</i>)	500 mL plus 200ml non-ionic surfactant per 100L spray- mix	Apply when conditions favour disease development and preferably prior to development of high levels of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59). DO NOT apply later than ZGS 59. Repeat spraying may be required for stripe rust. Regularly monitor the crop from 3-4 weeks after spraying for signs of re-infection and treat accordingly.
	Stripe rust (<i>Puccinia striiformis</i>) Septoria nodorum blotch (<i>Septoria nodorum</i>)		
Barley	Leaf rust (<i>Puccinia hordei</i>)	250 to 500 mL plus 200 ml non-ionic surfactant per 100L spray- mix	Apply when conditions favour disease development and preferably prior to development of high levels of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59). DO NOT apply later than ZGS 59. Apply the higher rate when disease is present on the top leaf or conditions are favourable for disease development at these stages of crop development.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION