

ALBAUGH VERSADO 800 WG FUNGICIDE

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply product in combination with spraying oils except to deciduous fruit in dormant period.
DO NOT use on sulfur sensitive crops when temperature exceeds 24°C
DO NOT apply this product during the heat of the day.
Tree and vine crops

Note for Rate column: In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Mixing/Application section.

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/praydrift

DO NOT allow bystanders to come into contact with the spray cloud
DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.
DO NOT apply unless the wind speed is between 3 and 20 Kilometres per hour at the application site during the time of application.
DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings 1 to 2 hours before sunrise and persist until 1 to 2 hours after sunrise.

Crop	Disease/Pest	State	Rate/100L	Critical Comments
Citrus	Citrus Rust Mite, Brown Citrus Rust Mite, Rust Mite, Mosic, Citrus Melanose	NSW, Vic, Qld, SA, WA only	400-500g	For all uses in this table: Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying method.
	White Loose Scale	NSW, WA only	200-300g	Apply during July-August. Leave at least 2 weeks between sprays of oil and product. Will assist control when applied late August.
				May be added to copper spray at peak fall or applied separately during Spring or Autumn.
Grapes (table grapes)	Vine Mite, Grapeleaf Blister Mite	NSW, Vic, SA, WA, Tas only	300g	Apply before spraying. Ensure thorough coverage.
	Vine Mite	Old only		
Grapes (wine grapes only)	Powdery Mildew (Dudium spp.) Mites	NSW, Vic, SA, WA, Tas only	100-200g	Apply immediately after budburst, then every 2-3 weeks or as required.
	Vine Mite, Grapeleaf Blister Mite	NSW, Vic, SA, WA, Tas only	300g	Apply at late budburst. During summer, build up of mites is checked by sulfur sprays for powdery mildew. An application immediately after picking may aid control in vineyards where mites are a severe problem.
Kiwifruit	Vine Mite	Old only		
	Powdery Mildew (Dudium spp.) Mites	NSW, Vic, SA, WA, Tas only	200-600g	Apply when shoots are 20 to 25cm long and repeat at intervals of 10 to 14 days during humid weather. Use the higher rates where flower pressure is high and/or a higher level of control is required. Crop phytotoxicity may occur when applications are made during hot conditions. DO NOT apply during the heat of the day. Use the higher rates early in the season before the temperatures rise. DO NOT apply more than 6kg per hectare.
Mangoes	Two spotted Mite	NSW, WA only	150-200g	Apply when the pest is present. Repeat application only when necessary.
	Powdery Mildew Mites	All States	400g	Apply as a foliar spray. Spray when pests are present. Repeat applications only when necessary. DO NOT apply during hot weather.
Peaches, Nectarines, Plums	Rust	NSW, Vic, SA, Tas only	200g	Apply as required during November to January.
		Old only	350g	
Pome Fruit	Brown Rot (<i>Monilinia fructicola</i>), Blossom Blight (<i>Monilinia laxa</i>)	NSW, Vic, SA, WA, Tas only	200g	Apply at petal fall and then as a cover spray at regular intervals beginning 3-4 weeks after petal fall.
	Powdery Mildew	All States	400g	Apply at petal fall only.
Rambutans	Powdery Mildew, Black Spot (SDB)	Vic, SA, Tas only	200g	Start spraying early. Apply at least 2 sprays before blossom. Apply every 2 weeks after blossom or as required. Regular sprays will also aid in the control of Black Spot (SDB).
	Powdery Mildew, Mites	All States	400g	Apply as a foliar spray. Spray when pests are present. Repeat applications only when necessary. DO NOT apply during hot weather.
Other Crops	Powdery Mildew, Two spotted Mite	Vic, SA, WA, Tas only	200g	Apply when pest or disease is first seen and repeat every 10-21 days.
	Ornamentals	Old only	350g	
Strawberries	Powdery Mildew	NSW only	200g	
	Powdery Mildew	NSW only	200-300g	Apply when the disease is first noticed then at 10-14 day intervals during humid weather.
Tomatoes for processing	Powdery Mildew, Bean Rust, Tomato Rust Mite, Bean Spider Mite, Two spotted Mite	Vic, SA, WA, Tas only	2 g/ha	Apply when pest or disease is first seen and repeat as necessary.
	Powdery Mildew, Bean Rust, Tomato Rust Mite, Bean Spider Mite	NSW only		Crop phytotoxicity may occur when applications occur during hot conditions. DO NOT apply during the heat of the day.
Vegetables (except Cucumbers or Backbeetles)	Powdery Mildew, Bean Rust, Tomato Rust Mite, Bean Spider Mite	Vic, SA, WA, Tas only	200g	
	Powdery Mildew, Bean Rust, Tomato Rust Mite, Bean Spider Mite	NSW only		
	Powdery Mildew, Bean Rust, Tomato Rust Mite, Bean Spider Mite	Old only	300g	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

MIXING

Measure the required amount in a small quantity of water. Half fill spray tank and commence agitation. Slowly add well stored product. Top up with required quantity of water. Maintain agitation throughout spraying procedure.

Application in the Crop and Wine

DILUTE SPRAYING

Use a sprayer designed to apply high volumes of water up to the point of runoff and matched to the crop being sprayed.

Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Directions for Use table for each 100L of water. Spray to the point of run-off.

The required dilute spray volume will change as the sprayer set up and operation may also need to be changed, as the crop grows.

CONCENTRATE SPRAYING

Use a sprayer designed and set up for concentrate spraying that is a sprayer which applies water volumes less than those required to reach the point of run-off and matched to the crop being sprayed.

Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.

Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy.

This is needed to calculate the concentrate mixing rate.

The mixing rate for concentrate spraying can then be calculated in the following way:

CONCENTRATE MIXING

(a) Dilute spray volume as determined above: For example 1500L/ha

(b) Your chosen concentrate spray volume: For example 500L/ha

(c) The concentration factor in this example is: 3 (i.e. 1500L ÷ 500L = 3).

(d) If the dilute label rate is 10 mL/100L, then the concentrate rate becomes 3 x 10, that is 30mL/100L of concentrate spray.

(e) The chosen spray volume, amount of product per 100L of water, and the sprayer set up and operation may need to be changed as the crop grows.

For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

COMPATIBILITY

This product is compatible with most commonly used fungicides and insecticides.

Any preparation of mixtures is entirely at the risk of the user and neither manufacturer or seller shall assume any responsibility in such cases. A preliminary test with small quantities is recommended prior to use. DO NOT mix with dimethylphthal derivatives such as DNOC or any oil containing products. DO NOT apply tank mixes to crops suffering from stress.

APVMA Approval Number: 94208/141819

Additional statements required by the WHS Regulations

Causes skin irritation. Very toxic to aquatic life.

If ON SKIN: Wash with plenty of water. Specific treatment (see supplemental first aid instruction on this label). If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash it before reuse. Collect spillage. Wash hands, forearms and face thoroughly after handling. Avoid release to the environment. Wear protective gloves. Dispose of container/contents to hazardous or special waste collection point, in accordance with local, regional, national and/or international regulation.

FUNGICIDE RESISTANCE WARNING

GROUP M2 FUNGICIDE

Albaugh VERSADO 800 WG fungicide is a member of the multi-site activity group of fungicides. For fungicide resistance management the product is a Group M2 fungicide. Some naturally occurring individual fungi resistant to the product and other Group M2 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product and other Group M2 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Albaugh Australia Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or watercourses with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

Single rinse or double rinse into spray tank. Do not dispose of undiluted chemicals on-site. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty container or product.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia: 13 11 21, New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS) which is available from the supplier and accessible from the Albaugh website albaugh.com.au

TERMS AND CONDITIONS OF SUPPLY, SALE AND USE

Many factors can affect or influence the activity of this product, including, but not limited to: weather and soil conditions, crop variety, treatment timing, water volume, application rates, spraying techniques, crop rotation, regional factors and the occurrence and development of strains resistant to the active ingredient.

Under certain circumstances, changes in activity or crop damage can occur. The manufacturer or supplier is unable to accept liability in these circumstances. All goods supplied by us are of a high grade and we believe them to be suitable for the purpose for which we regularly supply them, but we cannot exercise any control over their mixing, use or application which may affect the performance of the goods. All conditions and warranties statutory or otherwise as to the quality of fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff, agents, or the re-sellers of the product whether or not they supervise or assist in the use of such goods.

APVMA Approval Number: 94208/141819

ALBAUGH VERSADO 800 WG FUNGICIDE

ALBAUGH VERSADO 800 WG FUNGICIDE

ACTIVE CONSTITUENT: 800 g/kg SULFUR (S) present as elemental Sulfur

GROUP **M2** FUNGICIDE

For the control of powdery mildew, rust and mites in pome and stone fruit, citrus, grapevines, kiwifruit, strawberries and some vegetables as per the Directions for Use table

IMPORTANT: READ THE LABEL INSTRUCTIONS AND DIRECTIONS FOR USE BEFORE OPENING AND USING.

Net Contents
15 KG

FUNGICIDE



Manufactured globally to an American standard.

ALBAUGH AUSTRALIA PTY LTD
Level 1, 530 Little Collins Street,
Melbourne VIC 3000
Australia
Tel 03 9909 7183
www.albaugh.com.au

ALBAUGH
your alternative