

# BOGEN<sup>®</sup> 600 EC

**CONTROLS A VARIETY OF UNWANTED HARD TO KILL TREES,  
BUSH AND VARIOUS BROADLEAF WEEDS.**



Albaugh BOGEN<sup>®</sup> 600 EC Herbicide is a member of the pyridyloxy-carboxylate group of herbicides. BOGEN<sup>®</sup> 600 EC has the disruptors of plant cell growth mode of action.

## PRODUCT OVERVIEW

- **Application**  
Apply with hand directed equipment such as high volume spraying equipment, knapsack sprayer, knapsack misting machines with the following exceptions:  
Mistblower: Eucalyptus, Blackberry, Brigalow only  
Aircraft: Blackberry only  
Controlled Droplet Application (CDA): Blackberry only  
Gas Gun, Sprinkler Sprayer: Eucalyptus seedlings only  
Boomsprayer: Paddy Melon, Afghan, Bitter or Camel Melon only
- Highly effective on melon weed species.
- Weeds controlled include blackberry, camphor laural, English broom, gorse, common prickly pear, saffron thistle, capeweed and lantana (refer to the label to see full weed list).
- Fallow crop land weed control.
- Firebreak weed control.
- Commercial/industrial land weed control.

### ACTIVE INGREDIENT

600 g/L Triclopyr

### H.R.A.C. GROUP

GROUP 4 HERBICIDE

### FORMULATION

Emulsifiable Concentrate

### POISON SCHEDULE

Schedule 6

### COMPARE TO

Garlon<sup>®</sup>

### MODE OF ACTION

Synthetic auxin

### SIGNAL HEADING

POISON  
KEEP OUT OF REACH OF  
CHILDREN  
READ SAFETY DIRECTIONS  
BEFORE OPENING OR USING

### PACKAGING

20LTR, 110LTR

### APVMA APPROVAL NO.

94156/141713

### STATES AND TERRITORIES REGISTERED

ACT, NSW, NT, QLD, SA, TAS, VIC,  
WA



Garlon<sup>®</sup> is a registered trademark of Corteva Agriscience