

# DITTO<sup>®</sup>

**CONTROLS WILD OATS, PARADOXA GRASS (ANNUAL PHALARIS), CANARY GRASS AND ANNUAL RYEGRASS IN WHEAT. HIGHLY ACTIVE ON WILD OATS AND CAN BE MIXED WITH MANY BROADLEAF HERBICIDES.**



## PRODUCT OVERVIEW

- Apply to wheat at 2 leaf to late jointing (Zadocks 12-37).
- May be tank mixed with compatible herbicides.

## KEY WEEDS CONTROLLED OR SUPPRESSED

Wild oats

Paradoxa grass (*Phalaris paradoxa*)

Canary grass (*Phalaris minor*)

Annual ryegrass

Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.

### KEY CROPS

Wheat

### ACTIVE INGREDIENT

240 g/L Clodinafop-propargyl  
60 g/L Cloquintocet-mexyl

### H.R.A.C. GROUP

GROUP 1 HERBICIDE

### FORMULATION

Emulsifiable Concentrate

### POISON SCHEDULE

Schedule 6

### COMPARE TO

Topik<sup>®</sup>

### MODE OF ACTION

Inhibitors of acetyl CoA  
carboxylase

### SIGNAL HEADING

POISON  
KEEP OUT OF REACH OF  
CHILDREN  
READ SAFETY DIRECTIONS  
BEFORE OPENING OR USING

### PACKAGING

5LTR

### APVMA APPROVAL NO.

81130/141541

### STATES AND TERRITORIES REGISTERED

ACT, NSW, NT, QLD, SA, TAS, VIC,  
WA



Topik<sup>®</sup> is a registered trademark of Syngenta