

# FLUROPIX® 400 EC

A SELECTIVE HERBICIDE FOR POST-EMERGENT BROADLEAF WEED CONTROL



## PRODUCT OVERVIEW

- Use Albaugh FLUROPIX® 400 EC Herbicide for the control of a wide range of broadleaf weeds in fallow, lucerne, maize, millets, pastures, poppies, sorghum, sugar cane, sweet corn, winter cereals. Also for the control of woody weeds in agricultural non-crop areas, commercial and industrial areas, forests, pastures and rights-of-way.
- Controls annual broadleaf weeds.
- FLUROPIX® 400 EC is a systemic herbicide that moves within the plant, controlling exposed and underground plant tissue.
- Broad tank mix flexibility with most grass and broadleaf herbicides.

## KEY WEEDS CONTROLLED OR SUPPRESSED

Bathurst burr, noogoora burr, black bindweed (climbing buckwheat), mimosa pigra, bellyache bush, blackberry nightshade, bokhara clover, caltrop (yellow vine) (tribulus terrestris) (T. micrococcus), cobblers pegs, cockspur thorn, creeping lantana, crofton weed, mistflower, docks (rumex spp.), hexham scent, honey locust, small flowered mallow (marshmallow) (malva parviflora), lantana.

Woody weed situations: Weeds must be actively growing to attain optimal effect. Delay the treatment of regrowth following bulldozing, slashing, burning, ploughing or a previous chemical treatment until it has at least 1 metre of new, vigorous, growth.

### ACTIVE INGREDIENT

400g/L Fluroxypyr present as the methyl heptyl ester

### H.R.A.C. GROUP

**GROUP 4 HERBICIDE**

### FORMULATION

Emulsifiable Concentrate

### POISON SCHEDULE

Schedule 5

### COMPARE TO

Various

### MODE OF ACTION

Pyridine group, disrupters of plant cell growth

### SIGNAL HEADING

CAUTION  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

### PACKAGING

20LTR, 110LTR, 1000LTR

### APVMA APPROVAL NO.

81092/141581

### STATES AND TERRITORIES REGISTERED

ACT, NSW, NT, QLD, SA, TAS, VIC, WA

See label for complete application guidelines and restrictions.

