

# HEXANIL<sup>®</sup> 250 SL

**FOR THE CONTROL OF CERTAIN BROADLEAF WEEDS, PERENNIAL AND ANNUAL GRASSES, WOODY WEEDS IN PINUS RADIATA PLANTATIONS, PASTURE SITUATIONS AND COMMERCIAL AND INDUSTRIAL AREAS AND RIGHTS OF WAY.**



## PRODUCT OVERVIEW

- Albaugh HEXANIL<sup>®</sup> 250 SL Herbicide is both foliar and root absorbed and will give residual soil activity. For best foliar and residual control, apply during periods of active plant growth.
- Application when vegetation is dormant, semi-dormant, under moisture stress or during winter months may not be as effective.
- In temperate winter rainfall areas where temperatures are generally below 25°C during periods of active growth, overall effect may be slower to appear.
- For best control of perennials, apply during the months of spring in these areas.
- In WA and SA apply only during April-August when adequate rainfall is expected.
- Sufficient moisture is required after treatment to promote root absorption by the weed species.

## KEY USES

Pinus radiata.

Commercial and industrial areas, rights of way, around agricultural buildings and pasture situations.

### ACTIVE INGREDIENT

250 g/L Hexazinone

### H.R.A.C. GROUP

**GROUP 5 HERBICIDE**

### FORMULATION

Soluble Concentrate

### POISON SCHEDULE

Schedule 5

### COMPARE TO

various

### MODE OF ACTION

Triazinones; inhibitors of photosynthesis at photosystem II

### SIGNAL HEADING

CAUTION  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

### PACKAGING

10LTR, 20LTR

### APVMA APPROVAL NO.

81083/141683

### STATES AND TERRITORIES REGISTERED

ACT, NSW, NT, QLD, SA, TAS, VIC, WA



See label for complete application guidelines and restrictions.