

# TUKEN<sup>®</sup> EXTRA

**FOR THE CONTROL OF CERTAIN BROADLEAF WEEDS IN WINTER CEREALS AND CLOVER AS SPECIFIED**



## PRODUCT OVERVIEW

- Cereals: Wheat, barley, oats, triticale, cereal rye.
- Pasture: Newly sown and established clover-based pasture, clover for hay and seed production.
- Cereals: After application some transient crop yellowing may occur. This usually appears as yellow or white banding on leaves. Provided the crop is not under stress from pre-emergent herbicide, root disease, insect damage, frost, dry or excessively moist conditions, the development of the crop and subsequent growth will be unaffected.
- Oats: The tolerance of varieties Esk and Nile (the two main varieties grown in Tasmania) to Albaugh TUKEN<sup>®</sup> Extra Herbicide has not been tested. Test a small area of crop before using TUKEN<sup>®</sup> Extra over large areas. Consult your local Albaugh representative for advice on specific varieties.

## KEY WEEDS CONTROLLED

Canola (Rape), Brassica napus  
 Charlock, Sinapis arvensis  
 Common Sowthistle (Milk Thistle), Sonchus oleraceus  
 Cowvine, Ipomoea lonchophylla  
 Deadnettle, Lamium amplexicaule  
 Dock, Rumex spp.  
 Fat Hen, Chenopodium album  
 Fumitory, Fumitory spp.  
 Hexham Scent (King Island Melilot), Melilotus indicus  
 Hyssop Loosestrife, Lythrum hyssopifolia  
 Indian Hedge Mustard, Sisymbrium orientale  
 Long Storksbill, Erodium botrys  
 Mouse-eared Chickweed, Cerastium glomeratum  
 Paterson's Curse, Echium plantagineum  
 Prickly Lettuce, Lactuca serriola  
 Rough Poppy, Papaver hybridum  
 Scarlet Pimpernel, Anagallis arvensis  
 Skeleton Weed, Chondrilla juncea  
 Stemless Thistle, Onopordum acaulon  
 Tree Hogweed, Polygonum patulum  
 Variegated Thistle, Silybum marianum  
 Volunteer Lupins, Lupinus spp.  
 Wild Radish, Raphanus raphanistrum  
 Wireweed (Hogweed), Polygonum aviculare

Capeweed, Arctotheca calendula  
 Chickweed, Stellaria media  
 Corn Gromwell, Buglossoides arvensis  
 Crassula, Crassula spp.  
 Dense-flower Fumitory, Fumaria densiflora  
 Doublegee (Spiny Emex), Emex australis  
 Fireweed, Senecio spp.  
 Hedge Mustard, Sisymbrium officinale  
 Horehound, Marrubium vulgare  
 Iceplant, Mesembryanthemum spp.  
 London Rocket, Sisymbrium irio  
 Marshmallow, Malva parviflora  
 Night-scented Stock, Matthiola longipetala  
 Peppergrass, Lepidium spp.  
 Purple Goosefoot, Scleroblitum atriplicinum  
 Saffron Thistle, Carthamus lanatus  
 Shepherd's Purse, Capsella bursa-pastoris  
 Sorrel, Rumex spp.  
 Toad Rush, Juncus bufonius  
 Turnip Weed, Rapistrum rugosum  
 Vetch (Tares), Vicia sativa  
 Ward's Weed, Carrichtera annua  
 Wild Turnip, Brassica tournefortii



Jaguar<sup>®</sup> is a registered trademark of Bayer

### ACTIVE INGREDIENT

250 g/L MCPA present as the ethyl hexyl ester  
 25 g/L Diflufenican

### H.R.A.C. GROUP

**GROUP 4 12 HERBICIDE**

### FORMULATION

Emulsifiable Concentrate

### POISON SCHEDULE

Schedule 5

### COMPARE TO

Jaguar<sup>®</sup>

### MODE OF ACTION

Phenoxy and nicotinanalide groups, acts by inhibiting carotenoid biosynthesis and disrupting plant cell growth

### SIGNAL HEADING

CAUTION  
 KEEP OUT OF REACH OF CHILDREN  
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING

### PACKAGING

5LTR, 10LTR, 20LTR, 110LTR, 1000LTR

### APVMA APPROVAL NO.

81044/141684

### STATES AND TERRITORIES REGISTERED

ACT, NSW, NT, QLD, SA, TAS, VIC, WA