

Audible®

THIS PRODUCT IS A COPACK

Products Labels Included:

Audible®B and Draft®

Audible®B

For label, see pages 2-23.

Registration Number: 33706

Draft®

For label, see pages 24-52.

Registration Number: 31904

Manufacturer:

Albaugh, LLC.
1525 NE 36th St.
Ankeny, IA 50021, USA

2022-3214
2022-08-04

(Container)

Audible B Herbicide

GROUP	4	HERBICIDE
-------	---	-----------

Audible B Herbicide is a selective herbicide for postemergent control of annual broadleaved weeds including cleavers and certain ALS-resistant broadleaved weeds such as kochia in spring wheat, durum wheat, winter wheat, spring barley and oats. Audible B Herbicide will also suppress hemp-nettle, common chickweed and wild buckwheat.

COMMERCIAL (AGRICULTURAL)

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: fluroxypyr, present as 1-methylheptyl ester 333 g a.e./L
Emulsifiable concentrate

REGISTRATION NO. 33706 PEST CONTROL PRODUCTS ACT

WARNING
EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER

NET CONTENTS: 1 L - bulk

Albaugh LLC
1525 NE 36th Street
Ankeny, IA 50021 USA
1-800-247-8013

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

Causes eye irritation. DO NOT get in eyes. May irritate the skin. Avoid contact with skin. Potential skin sensitizer.

For agricultural/crop scenarios: Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12-hours.

Do not use in residential areas. Residential areas are defined as any use site where bystanders including children could be exposed during or after application. This includes homes, schools, parks, playgrounds, playing fields, public buildings, or any other area where the general public including children could be exposed.

Apply only when potential for drift to areas of human habitation or activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversion, application equipment and sprayer settings.

PROTECTIVE CLOTHING AND EQUIPMENT

Wear a hat, coveralls over a long-sleeved shirt and long pants, socks, rubber boots or work boots and chemical-resistant gloves during spraying or application. In addition, wear goggles or face shield during mixing, loading, repair and clean-up or when handling the concentrate. Applicators using a closed cab are not required to wear chemical-resistant gloves.

OPERATOR USE PRECAUTIONS

Wear freshly laundered clothing and clean protective equipment daily. At completion of spraying or end of the day take a shower immediately. Wash thoroughly with soap and water before eating, drinking, smoking or using the toilet. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse. If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Avoid contact with eyes, skin and clothing. Avoid breathing spray mist.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE. Do not use or store near heat or open flame.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

2022-3214
2022-08-04

TOXICOLOGICAL INFORMATION

THIS PRODUCT CONTAINS A PETROLEUM DISTILLATE. Vomiting may cause aspiration pneumonia. The decision of whether to induce vomiting or not should be made by the attending physician. If lavage is performed, suggest endotracheal and/or oesophageal control. Danger from lung aspiration of petroleum based solvents must be weighed against toxicity when considering emptying the stomach. No specific antidote. Employ supportive care. Treatment should be based on the judgment of the physician in response to reactions of the patient.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants and aquatic organisms. Observe buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

This product contains an active ingredient and aromatic petroleum distillates which are toxic to aquatic organisms.

STORAGE

Store in original containers in a secure, dry heated storage. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. To prevent contamination, store this product away from food or feed. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL

Recyclable Containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Returnable Containers

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

2022-3214
2022-08-04

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

Causes eye irritation. DO NOT get in eyes. May irritate the skin. Avoid contact with skin. Potential skin sensitizer.

For agricultural/crop scenarios: Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12-hours.

Do not use in residential areas. Residential areas are defined as any use site where bystanders including children could be exposed during or after application. This includes homes, schools, parks, playgrounds, playing fields, public buildings, or any other area where the general public including children could be exposed.

Apply only when potential for drift to areas of human habitation or activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversion, application equipment and sprayer settings.

PROTECTIVE CLOTHING AND EQUIPMENT

Wear a hat, coveralls over a long-sleeved shirt and long pants, socks, rubber boots or work boots and chemical-resistant gloves during spraying or application. In addition, wear goggles or face shield during mixing, loading, repair and clean-up or when handling the concentrate. Applicators using a closed cab are not required to wear chemical-resistant gloves.

OPERATOR USE PRECAUTIONS

Wear freshly laundered clothing and clean protective equipment daily. At completion of spraying or end of the day take a shower immediately. Wash thoroughly with soap and water before eating, drinking, smoking or using the toilet. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse. If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Avoid contact with eyes, skin and clothing. Avoid breathing spray mist.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE. Do not use or store near heat or open flame.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

2022-3214
2022-08-04

TOXICOLOGICAL INFORMATION

THIS PRODUCT CONTAINS A PETROLEUM DISTILLATE. Vomiting may cause aspiration pneumonia. The decision of whether to induce vomiting or not should be made by the attending physician. If lavage is performed, suggest endotracheal and/or oesophageal control. Danger from lung aspiration of petroleum based solvents must be weighed against toxicity when considering emptying the stomach. No specific antidote. Employ supportive care. Treatment should be based on the judgment of the physician in response to reactions of the patient.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants and aquatic organisms. Observe buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

This product contains an active ingredient and aromatic petroleum distillates which are toxic to aquatic organisms.

STORAGE

Store in original containers in a secure, dry heated storage. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. To prevent contamination, store this product away from food or feed. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL

Recyclable Containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Returnable Containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

GENERAL INFORMATION

Audible B Herbicide is a selective postemergent herbicide for the control of annual broadleaved weeds in spring wheat, durum wheat, winter wheat, spring barley and oats, not underseeded with legumes. Audible B Herbicide offers a novel mode of action for controlling hard-to-kill annual broadleaved weeds such as kochia (including ALS resistant biotypes) and cleavers. Audible B Herbicide will also provide suppression of hempnettle, wild buckwheat and chickweed.

MODE OF ACTION

Audible B Herbicide is a systemic auxin-type herbicide and moves within the plant for control of exposed and underground plant tissues. The product controls weeds by disrupting normal plant growth patterns. Symptoms of weeds include epinasty (twisting of the stems) and swollen nodes.

GENERAL USE PRECAUTIONS

Sensitive Plants

Do not apply Audible B Herbicide directly to, or otherwise permit it to come in direct contact with susceptible crops or desirable plants including alfalfa, edible beans, canola, flowers and ornamentals, lentils, lettuce, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tomatoes or tobacco.

Non-Target Sites

Do not apply where proximity of susceptible crops or other desirable plants is likely to result in exposure to spray or spray drift. See Environmental Hazards section of the label.

Crop Rotation

Fields previously treated with Audible B Herbicide can be seeded the following year to alfalfa, barley, canola, corn, dry beans, fababeans, flax, forage grasses, lentils, mustard, oats, peas, potatoes, rye, soybeans, sugar beets, sunflowers or wheat or fields can be summerfallowed.

Tank Mixtures

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Albaugh LLC at 1-800-247-8013 for information before mixing any pesticide or fertilizer that is not specifically recommended on this label.

Spray Equipment Precaution

Do not apply through any type of irrigation system.

To Reduce Spray Drift

- Use nozzles delivering higher volumes and coarser droplets.
- Use low pressures (200 to 275 kPa).
- Spray when the wind velocity is 15 km/hr or less.
- Spot treatments should only be applied with a calibrated boom to prevent over-application.

Equipment Clean-Up

Equipment used to apply Audible B Herbicide should not be used to apply other pesticides to sensitive crops without thorough cleaning. Contact your Audible B Herbicide dealer for detailed equipment cleaning procedures.

DIRECTIONS FOR USE

READ THE ENTIRE LABEL BEFORE USE. FAILURE TO FOLLOW LABEL INSTRUCTIONS MAY RESULT IN ERRATIC WEED CONTROL OR CROP DAMAGE. DO NOT APPLY TO CROPS UNDERSEEDDED WITH LEGUMES.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

2022-3214
2022-08-04

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

Aerial application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing or rotor span.

Crops Registered

spring wheat
durum wheat
winter wheat
spring barley
oats

AUDIBLE B HERBICIDE ALONE

Apply Audible B Herbicide with ground or aerial application equipment (see Application directions section) for postemergent control of cleavers and kochia, including ALS resistant kochia and other broadleaved weeds infesting spring wheat, durum wheat, winter wheat, spring barley and oats. See following table for weeds controlled/suppressed and applications rates.

Weeds Controlled or Suppressed By Audible B Herbicide Alone

Weeds Controlled	Audible B Herbicide Rate (L/ha)
chickweed* (up to 8 cm)	0.41
cleavers (1-6 whorls)	0.21
cleavers (1-8 whorls)	0.31
kochia* (2-8 leaf)	0.31
round-leaved mallow (1-6 leaf)	0.41
volunteer flax (1-12 cm)	0.31
Weeds Suppressed	Audible B Herbicide Rate (L/ha)
hempnettle (2-6 leaf)	0.41
stork's-bill (1-8 leaf)	0.31
wild buckwheat (1-8 leaf)	0.21

* Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.

NOTE: Audible B Herbicide activity is influenced by weather conditions. Optimum activity requires active crop and weed growth. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost before application (3 days) or shortly after (3 days) may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions, e.g. drought, heat or cold stress, or if weeds have initiated flowering, or if heavy infestations exist.

Application Timing

Apply to actively growing spring wheat, durum wheat, spring barley and oats from the 2-leaf crop growth stage up to and including initiation of stem elongation. Apply to winter wheat in the spring from the 3 tiller stage to just before the flag leaf stage. Apply when weeds are actively growing. Best results are obtained from applications made to seedling weeds. Only weeds emerged at the time of treatment will be controlled. Extreme growing conditions such as drought or near freezing temperature prior to, and

2022-3214
2022-08-04

following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Foliage that is wet at the time of application may decrease control. Audible B Herbicide applications are rainfast within 1 hour after application. Under conditions of low crop and high weed density, control may be reduced.

Preharvest/Grazing Intervals (Audible B Herbicide applied alone)

- Livestock may be grazed on treated crops 3 days following application.
- Do not harvest the treated crop within 60 days after application.

APPLICATION DIRECTIONS (GROUND AND AERIAL APPLICATIONS)

(1) Ground Application

Using ground equipment, apply Audible B Herbicide alone or with tank mix partners as a broadcast treatment. Apply Audible B Herbicide alone at the recommended rates in a spray volume of 50-100 L water/ha. When tank mixing refer to the label of the tank mix partner label for additional instructions, water volumes, precautions and weeds controlled.

(2) Aerial Application

Do not use human flaggers.

Use Audible B Herbicide alone or tank mixed with a product registered for aerial application as a broadcast treatment. Apply the recommended rates of Audible B Herbicide and tank mix partner as indicated elsewhere on this label. Apply in a spray volume of 30-50 L water/ha. Refer to the tank mix partner label for additional instructions, precautions and weeds controlled.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate swath marking devices.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

2022-3214
2022-08-04

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-800-667-3852 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

- The interaction of many equipment-and-weather-related factors determine the potential for spray drift. Users are responsible for considering all these factors when making decisions.

The following drift management requirement must be followed to avoid off-target drift movement from aerial applications: Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

TANK-MIX COMBINATIONS - AUDIBLE B HERBICIDE + LIQUID ACHIEVE¹ SC HERBICIDE OR ACHIEVE LIQUID HERBICIDE (Do not apply to oats)

For control of wild oats and green foxtail in spring wheat, winter wheat and spring barley apply 0.5 L/ha of Liquid Achieve SC or Achieve Liquid with Audible B Herbicide. Refer to the above table for rates and broadleaved weeds controlled or suppressed with Audible B Herbicide. Read both product labels thoroughly for more information on precautions and additional directions for use. To control additional broadleaved weeds tank mix with either 2,4-D ester or MCPA ester up to a maximum of 560 g ae/ha. (equivalent to 1.0 L/ha of 2,4-D LV600 Herbicide (0.85 L/ha 2,4-D Ester 700 Herbicide) or 1.12 L/ha of MCPA LV500 Herbicide) (0.93 L/ha MCPA Ester 600 Herbicide) or 1.5 to 2.0 L/ha Curtail M Herbicide. Refer to the 2,4-D ester or MCPA ester tank mix partner label for rates less than 560 g ae/ha and weeds controlled.

Mixing Instructions

1. Begin to fill sprayer tank with clean water, and engage agitator. **Agitation must be continued throughout the entire mixing and spraying procedure.**
2. When the sprayer is three quarters full of water, add Liquid Achieve SC or Achieve Liquid. If more than one container of Liquid Achieve SC or Achieve Liquid is to be used, add all Liquid Achieve SC or Achieve Liquid containers prior to adding Audible B Herbicide.
3. Add Audible B Herbicide next. If more than one container of Audible B Herbicide is to be used, add all Audible B Herbicide containers prior to adding any other tank-mix partners.
4. If 2,4-D ester, MCPA ester or Curtail M are also being added, add these next. If more than one container of 2,4-D ester, MCPA ester or Curtail M is to be used, add all 2,4-D ester, MCPA ester or Curtail M containers prior to adding Turbocharge D or Turbocharge.
5. Lastly, add the required amount of Turbocharge D or Turbocharge and then continue to fill tank to desired level with water.

TANK-MIX COMBINATIONS - AUDIBLE B HERBICIDE + CURTAILTM M HERBICIDE

Audible B Herbicide mixed with Curtail M Herbicide will control a wide spectrum of broadleaf weeds. Refer to Curtail M Herbicide label for additional precautions and directions for use, particularly crop rotation restrictions.

Mixing Instructions

Add the correct amount of Audible B Herbicide to spray tank 1/2 filled with water and agitate. Add the correct amount of Curtail M Herbicide and continue to agitate.

Timing of Application

Apply postemergence when weeds are in the seedling stage (2-4 leaf) and when spring wheat, durum wheat, spring barley and oats are in the 3-leaf to just before the flag leaf emergence stage. Apply to winter wheat in the spring from the 3 tiller stage to just before the flag leaf stage. Do not apply later than the flag-leaf stage.

Low Rate: To control or suppress the following weeds apply a uniform spray containing Audible B Herbicide at 0.31 L/ha tank mixed with Curtail M Herbicide at 1.5 L/ha.

Weeds Controlled

Canada thistle (low infestations)	mustard, wild
cleavers (1-8 whorls)	shepherd's-purse
flax, volunteer (1-12 cm)	stinkweed
flixweed (spring seedling 2-4 leaf)	sunflower, volunteer
kochia***	wild buckwheat (1-8 leaf)
lamb's-quarters	

Weeds Suppressed

stork's-bill (1-8 leaf)

High Rate: To control or suppress the following weeds apply a uniform spray containing Audible B Herbicide at 0.41 L/ha tank mixed with Curtail M Herbicide at 2.0 L/ha.

Weeds Controlled

Canada thistle*	lamb's-quarters	sow-thistle (perennial) *
canola (volunteer)	redroot pigweed	stinkweed
cleavers (1-8 whorls)	round-leaved mallow (1-6 leaf)	stork's-bill (1-8 leaf)
common chickweed*	Russian pigweed	sunflower (volunteer)
common groundsel	scentsless chamomile	tartary buckwheat
dandelion**	shepherd's-purse	volunteer flax (1-12 cm)
flixweed**	smartweed	wild buckwheat (1-8 leaf)
hemp-nettle (2-6 leaf stage)	sow-thistle (annual)	wild mustard
kochia***		

* Season long control, with some regrowth in the fall (top growth control).

** Spring rosettes only.

*** Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.

The active ingredients in this tank mixture will only control chickweed that is emerged at TIME OF application. Chickweed plants which emerge after application will not be controlled. To improve the reduction in chickweed population at the end of season, delay the timing of application as late as possible to when the majority of chickweed plants have emerged. The reduction in chickweed population will also be improved in crops which are more competitive and which allow limited light penetration.

Pre-Harvest/Grazing Intervals (Audible B Herbicide + Curtail M Herbicide)

- Do not cut or graze treated fields within 7 days after application.
- Do not harvest the treated mature crop within 60 days after application.

When tank mixing with Curtail M Herbicide, wear goggles or face shield, coveralls and chemically resistant gloves, apron and boots when mixing, loading and during clean-up and repair.

Curtail M Herbicide contains an ingredient that is rapidly absorbed through the skin and respiratory tract.

2022-3214
2022-08-04

TANK MIXES WITH AUDIBLE B HERBICIDE + CURTAIL M HERBICIDE FOR ANNUAL GRASS CONTROL (Do not apply to oats)

Tank mixtures of Audible B Herbicide + Curtail M Herbicide with other herbicides will provide control of additional weeds. See following table for tank mixes with Audible B Herbicide + Curtail M Herbicide. Follow all precautions, minimal interval to harvest and directions for use on the Audible B Herbicide and tank-mix partner labels.

Tank-Mix Combinations with Audible B Herbicide at 0.31 L/ha + Curtail M Herbicide at 1.5 L/ha or Audible B Herbicide at 0.41 L/ha + Curtail M Herbicide at 2.0 L/ha

Herbicide Tank-Mix Partner	Crops Registered	Rate of Tank-Mix Partner	Adjuvant Rate	Additional Weeds
Liquid Achieve SC or Achieve Liquid	spring wheat, durum wheat, winter wheat, spring barley	0.5 L/ha	Turbocharge D or Turbocharge 0.5 L/100 L of spray volume (0.5% v/v)	green foxtail wild oats
Assert 300 SC	spring wheat, durum wheat, spring barley	1.3 L/ha	Refer to Assert 300 SC label	wild oats (1-3 leaf)
		1.6 L/ha		wild oats (4 leaf)
Puma 120 Super	spring wheat, durum wheat, spring barley	385 mL/ha	None required	green foxtail only
		770 mL/ha		green foxtail, wild oats & barnyard grass
Everest 70 WDG or Everest Solupak 70 DF	spring wheat, durum wheat	43 g/ha	Refer to Everest label	green foxtail wild oats

TANK-MIX COMBINATION - AUDIBLE B HERBICIDE + 2,4-D ESTER HERBICIDE (Do not apply to oats)

For control of a wide spectrum of broadleaved weeds (see following table) apply Audible B Herbicide tank mixed with 2,4-D ester herbicide. Apply postemergence when weeds are in the seedling stage (2-4 leaf) and when spring wheat, durum wheat, winter wheat and spring barley are in the 4-leaf to flag-leaf stage. Apply to winter wheat in the spring from the 3 tiller stage to just before the flag leaf stage. Application before the 4-leaf stage of wheat and barley may cause severe twisting of leaves and stem, and head deformities which may reduce yield up to 10%. Do not apply later than the flag-leaf stage.

Rates of Applications

I. Low Rate: To control the following weeds apply a uniform spray containing 0.23 L/ha Audible B Herbicide and 420 g a.e./ha 2,4-D ester herbicide (equivalent to 0.64 L/ha of 2,4-D Ester LV700 Herbicide).

Weeds Controlled

bluebur	goat's-beard	shepherd's-purse
burdock	hoary cress*	stinkweed
canola (volunteer)	kochia***	sunflower (annual)
cleavers (1-6 whorls)	lamb's-quarters	vetch
clovers (sweet)	mustards (except green, dog & tansy)	wild radish
cocklebur	plantain	wild mustard
flixweed	prickly lettuce	wild buckwheat (1-6 leaf)
field horsetail*	ragweed	

II. High Rate: To control the above weeds plus the additional weeds listed below, apply a uniform spray containing 0.31 L/ha of Audible B Herbicide and 560 g a.e./ha of 2,4-D ester herbicide (equivalent to 0.85 L/ha of 2,4-D Ester LV700 Herbicide).

2022-3214
2022-08-04

Additional Weeds Controlled:

blue lettuce*	hairy galinsoga	Russian thistle
cleavers (1-8 whorls)	hedge bindweed	smartweed
dandelion**	lady's-thumb	stork's-bill (1-8 leaf)
docks	leafy spurge*	tansy mustard
dog mustard	oak-leaved goosefoot	tartary buckwheat
field bindweed*	redroot pigweed	volunteer flax (1-12 cm)
field peppergrass	round-leaved mallow (1-6 Leaf)	wild buckwheat (1-8 leaf)
gumweed		

Weeds Suppressed:

common chickweed*** (up to 8 cm)	hemp-nettle (2-6 leaf stage)	sow-thistle (perennial) *
Canada thistle*	sow-thistle (annual)	

* Top growth control only

** Spring rosettes

*** Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme

NOTE: The activity of Audible B Herbicide combined with 2,4-D ester herbicide is influenced by weather conditions. Optimum activity requires active crop and weed growth. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost before application (3 days) or shortly after (3 days) may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions, e.g. drought, heat or cold stress, or if weeds have initiated flowering, or if heavy infestations exist.

Mixing Instructions

Add the correct amount of Audible B Herbicide to spray tank 1/2 filled with water and agitate. Add the correct amount of 2,4-D ester herbicide (e.g. 2,4-D Ester LV700 Herbicide) and continue to agitate.

Pre-Harvest/Grazing Intervals (Audible B Herbicide + 2,4-D ester herbicide)

- Do not permit lactating dairy animals to graze fields within 7 days after application.
- Do not harvest forage or cut hay within 30 days after application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.
- Do not harvest the treated mature crop within 60 days after application.

TANK-MIX COMBINATIONS - AUDIBLE B HERBICIDE + 2,4-D ESTER HERBICIDE + OTHER HERBICIDES (Do not apply to oats)

Tank mixtures of Audible B Herbicide + 2,4-D ester herbicide with other herbicides will provide control of additional weeds. For products which may be tank mixed with Audible B Herbicide + 2,4-D ester herbicide and the additional weeds controlled see following table. Follow all precautions, minimal interval to harvest and directions for use on the Audible B Herbicide and tank-mix partner labels.

2022-3214
2022-08-04

Tank-Mix Combinations with Audible B Herbicide (0.31 L/ha) + 2,4-D ester herbicide (560 g a.e./ha) or Audible B Herbicide (0.23 L/ha) + 2,4-D ester herbicide (420 g a.e./ha)

Herbicide Tank-Mix Partner	Crops Registered	Rate of Tank-Mix Partner	Adjuvant Rate	Additional Weeds
Liquid Achieve SC or Achieve Liquid ^{♦♦♦}	spring wheat, durum wheat, winter wheat, spring barley	0.5 L/ha	Turbocharge D or Turbocharge 0.5 L/100 L of spray volume (0.5% v/v)	green foxtail, wild oats
Assert 300 SC	spring wheat, durum wheat, spring barley	1.3 L/ha	Refer to Assert 300 SC label	wild oats (1-3 leaf stage)
		1.6 L/ha		wild oats (4 leaf stage)
Puma 120 Super	spring wheat, durum wheat	385 mL/ha	None required	green foxtail only
		770 mL/ha		green foxtail, plus wild oats & barnyard grass
Simplicity ^{TM♦♦♦}	spring wheat, durum wheat, winter wheat	0.5 L/ha	None required	barnyard grass (1-5 leaf), common chickweed (up to 10 cm), hemp nettle (1-8 leaf), Japanese brome (1-6 leaf), wild oats (up to 4 leaf, 2 tiller), wild buckwheat (1-4 leaf)
Everest 70 WDG or Everest Solupak 70 DF	spring wheat, durum wheat	43 g/ha	Refer to Everest label	green foxtail wild oats

♦ This tank-mix may cause temporary crop injury. However, yield will not normally be affected. This tank-mix may reduce control of wild oats compared to Liquid Achieve SC or Achieve Liquid Herbicide alone.

♦♦ Wild oat control may be reduced when tank mixed with Audible B Herbicide + 2,4-D ester herbicide

♦♦♦ Tank mixing with Simplicity only applies to the low rate of Audible B Herbicide + 2,4-D ester

TANK-MIX COMBINATIONS – AUDIBLE B HERBICIDE + REFINE EXTRA 75 DF HERBICIDE OR REFINE EXTRA 75 TOSS-N-GO HERBICIDE OR DRAFT OR REFINE SG HERBICIDE OR REFINE SG TOSS-NGO HERBICIDE

For control of a wide spectrum of broadleaved weeds (see following table) Audible B Herbicide can be tank mixed with Refine Extra 75 DF Herbicide or Refine Extra 75 Toss-N-Go Herbicide or Draft or Refine SG Herbicide or Refine SG Toss-N-Go Herbicide. Apply post-emergence when weeds are actively growing, with best results on seedling weeds that are less than 10 cm in height. Only weeds emerged at the time of treatment will be controlled. Thorough covering of the target weeds is essential. The tank-mix can be applied to actively growing spring wheat, including durum, spring barley and oats from the 2-leaf crop growth stage up to and including initiation of stem elongation. The tank-mix of Refine Extra 75 DF Herbicide or Refine Extra 75 Toss-N-Go Herbicide or Draft or Refine SG Herbicide or Refine SG Toss-N-Go Herbicide with Audible B Herbicide allows for the control of Group 2 resistant cleavers at 0.21 L/ha and kochia 0.31 L/ha. Do not apply to crops underseeded to legumes or grasses.

Rates of Application To control the following weeds apply a uniform spray containing 0.21 to 0.41 litres of Audible B Herbicide with 20 g/ha Refine Extra 75 DF Herbicide or Refine Extra 75 Toss-N-Go Herbicide or Draft or 30 g/ha of Refine SG Herbicide or Refine SG Toss-N-Go Herbicide and 2 litres per 1000 L spray solution of non-ionic surfactant (Agral 90, Ag-surf or Citowett Plus) with a spray volume of 100 L/ha.

Weeds Controlled or Suppressed

Weeds Controlled at 0.21 L/ha of Audible B Herbicide and 20 g/ha Refine Extra 75 DF Herbicide or Refine Extra 75 Toss-N-Go Herbicide or Draft, or 30 g/ha of Refine SG Herbicide or Refine SG Toss-N-Go Herbicide:

annual smartweed (green smartweed, lady's thumb)	groundsel (common)	shepherd's purse
ball mustard	hemp-nettle	stinkweed
chickweed (1-6 leaf) ¹	kochia ²	tartary buckwheat
cleavers (1-4 whorls)	lamb's-quarters	volunteer rapeseed ³
corn spurry	narrow-leaved hawk's beard	volunteer sunflower
cow cockle	redroot pigweed	wild buckwheat ⁴
flixweed	Russian thistle	wild mustard

¹ Spray before canopy prevents thorough coverage of weeds. Chickweed emerging after application will not be controlled.

² Excluding Group 2 resistant kochia

³ The tank-mix will not control imazethapyr tolerant canola varieties (e.g. Clearfield varieties or other varieties with the Pursuit Smart trait)

⁴ Apply to actively growing wild buckwheat at the cotyledon to 3-leaf stage. Control may be reduced under environmental stress. Large plants may regrow after treatment.

Weeds suppressed at 0.21 L/ha Audible B Herbicide and 20 g/ha Refine Extra 75 DF Herbicide or Refine Extra 75 Toss-N-Go Herbicide or Draft, or 30 g/ha of Refine SG Herbicide or Refine SG Toss-N-Go Herbicide:

Canada thistle ⁵	sow thistle ⁵
round leaved mallow (2-6 leaf) ⁶	stork's bill (2-6 leaf stage)
scentless chamomile	toadflax (up to 15 cm) ⁷

⁵ Apply when most of the thistles have emerged and are actively growing. Apply before bud stage and before crop canopy prevents thorough coverage of thistles, when thistles are no larger than 15 cm in height for best top growth control. A single application will effectively inhibit the ability of emerged thistles to compete with the crop. Later emerging thistles will not be controlled.

⁶ Apply to actively growing round leaved mallow. Will keep mallow stunted, but may not be able to reduce overall populations

⁷ A control program for this weed includes chemical application and frequent tillage.

2022-3214
2022-08-04

Additional Weeds Controlled at 0.31 L/ha of Audible B Herbicide and 20 g/ha Refine Extra 75 DF Herbicide or Refine Extra 75 Toss-N-Go Herbicide or Draft, or 30 g/ha of Refine SG Herbicide or Refine SG Toss-N-Go Herbicide:

kochia (2-8 leaf; including group 2 resistant biotypes)
volunteer flax (1-12 cm)

Additional Weeds Controlled and Suppressed at 0.41 L/ha Audible B Herbicide and 20 g/ha Refine Extra 75 DF Herbicide or Refine Extra 75 Toss-N-Go Herbicide or Draft, or 30 g/ha of Refine SG Herbicide or Refine SG Toss-N-Go Herbicide:

Weeds Controlled	Weeds Suppressed
round leaved mallow (1-6 leaf)	chickweed (up to 8 cm; including group 2 resistant biotypes)

Mixing Instructions

Fill the tank to 1/3 full of clean water, and with the agitator running, add the required amount of Refine Extra 75 DF Herbicide or Refine Extra 75 Toss-N-Go Herbicide or Draft or Refine SG Herbicide or Refine SG Toss-N-Go Herbicide and continue to agitate for a minimum of 5 minutes to ensure that Refine Extra 75 DF Herbicide or Refine Extra 75 Toss-N-Go Herbicide or Draft or Refine SG Herbicide or Refine SG Toss-N-Go Herbicide is completely dissolved. Add the required amount of Audible B Herbicide at a reduced agitation. Add the rest of the water and then add adjuvant. Use a spray volume of 100 L/ha.

Pre-Harvest/Grazing Intervals

- Livestock may not be grazed or fed crops within 7 days of application
- Do not harvest the treated crop within 60 days of application

**TANK-MIX COMBINATIONS – AUDIBLE B HERBICIDE + 2,4-D ESTER HERBICIDE + REFINE EXTRA 75 DF HERBICIDE OR REFINE EXTRA 75 TOSS-N-GO HERBICIDE OR DRAFT OR REFINE SG HERBICIDE OR REFINE SG TOSS-N-GO HERBICIDE OR RETAIN A HERBICIDE
(Do not apply to oats)**

Spring Wheat (excluding durum) and spring barley

For control of a wide spectrum of broadleaf weeds, including cleavers, Audible B Herbicide may be tank mixed with 2,4-D ester herbicide and Refine Extra 75 DF Herbicide or Refine Extra 75 Toss-N-Go Herbicide or Draft or Refine SG Herbicide or Refine SG Toss-N-Go Herbicide or Retain A Herbicide. Apply Audible B Herbicide at 0.16 L/ha tank mixed with Refine Extra 75 DF Herbicide or Refine Extra 75 Toss-N-Go Herbicide or Draft at 20 g/ha, or Refine SG Herbicide or Refine SG Toss-N-Go Herbicide or Retain A Herbicide at 30 g/ha, 2,4-D ester herbicide at 280 g a.e./ha (equal to 0.5 L/ha of 2,4-D Ester LV 600 or 0.42 L/ha of 2,4-D Ester LV 700) and a non-ionic surfactant (Agral 90, Ag-Surf or Citowett Plus) at 2 L per 1000 L of spray solution. A spray volume of 100 L/ha should be utilized. Apply postemergence when weeds are in the seedling up to 4-leaf/whorls stage, and when spring wheat (excluding durum) and spring barley are in the 4-leaf to flag-leaf stage. **Do not apply to crops underseeded to legumes.**

Weeds Controlled or Suppressed

Weeds Controlled:

annual smartweed (green smartweed, lady's thumb)	groundsel (common)	shepherd's purse
ball mustard	hemp-nettle	stinkweed
chickweed (1-6 leaf) ¹	kochia ³	tartary buckwheat
cleavers ²	lamb'-quarters	volunteer rapeseed ⁴
corn spurry	narrow-leaved hawk's beard	volunteer sunflower
cow cockle	redroot pigweed	wild buckwheat ⁵
flixweed	Russian thistle	wild mustard

¹ Spray before crop canopy prevents thorough coverage of weeds. Chickweed emerging after application will not be controlled.

² Excluding Group 2 resistant cleavers

2022-3214
2022-08-04

³ Excluding Group 2 resistant kochia

⁴ The tank-mix will not control imazethapyr tolerant canola varieties (e.g. Clearfield varieties or other varieties with the Pursuit Smart trait)

⁵ Apply to actively growing wild buckwheat from the cotyledon to 3-leaf stage. Control may be reduced under environmental stress. Large plants may regrow after treatment

Weeds Suppressed:

Canada thistle ⁶	sow thistle ⁶
round leaved mallow (2-6 leaf) ⁷	stork's bill (2-6 leaf)
scentless chamomile	toadflax (to 15 cm) ⁸

⁶ Apply when most of the thistles have emerged and are actively growing. Apply before bud stage, and before crop canopy prevents thorough coverage of thistles, when thistles are no larger than 15 cm in height for best top growth control. A single application will effectively inhibit the ability of emerged thistles to compete with the crop. Later emerging thistles will not be controlled.

⁷ Apply to actively growing round leaved mallow. Will keep mallow stunted, but may not be able to reduce overall populations.

⁸ A control program for this weed includes chemical application and frequent tillage.

Mixing Instructions

Fill the tank to 1/3 full of clean water. With agitator running, add required amount of Refine Extra 75 DF Herbicide or Refine Extra 75 Toss-N-Go Herbicide or Draft or Refine SG Herbicide or Refine SG Toss-N-Go Herbicide or Retain A Herbicide and continue to agitate for a minimum of 5 minutes to ensure the Refine Extra 75 DF Herbicide or Refine Extra 75 Toss-N-Go Herbicide or Draft or Refine SG Herbicide or Refine SG Toss-N-Go Herbicide or Retain A Herbicide is completely dissolved. After it is fully dissolved add the Audible B Herbicide with continuous agitation. Add the 2,4-D ester herbicide and the rest of the water. Then add adjuvant. Use a spray volume of 100 L/ha. Consult the 2,4-D label for additional application instructions and use precautions.

Preharvest/Grazing Intervals (Audible B Herbicide + Refine Extra or Draft + 2,4-D ester)

- Allow 7 days between application and grazing of lactating dairy animals.
- Do not harvest forage or cut hay within 30 days after application.
- Remove meat animals from treated fields at least 3 days prior to slaughter.
- Do not harvest the treated mature crop within 60 days after application.

TANK-MIX COMBINATION - AUDIBLE B HERBICIDE + MCPA ESTER HERBICIDE

For control of a wide spectrum of broadleaved weeds (see following table) apply Audible B Herbicide at 0.31 L/ha tank mixed with MCPA LV500 Herbicide at 1.12 L/ha or MCPA Ester 600 Herbicide at 0.93 L/ha. Apply postemergence when weeds are in the seedling stage (2-4 leaf) and when spring wheat, durum wheat, spring barley and oats are in the 3-leaf to flag-leaf stage. Apply to winter wheat in the spring from the 3 tiller stage to just before the flag leaf stage. Application before the 3-leaf stage of wheat and barley may cause severe twisting of leaves and stem, and head deformities which may reduce yield up to 10%. Do not apply later than the flag-leaf stage. Do not apply to cereals underseeded with a legume.

Mixing Instructions

Add the correct amount of Audible B Herbicide to spray tank 1/2 filled with water and agitate. Add the correct amount of MCPA ester herbicide and continue to agitate.

Weeds Controlled or Suppressed by Audible B Herbicide+ MCPA ester herbicide Tank-Mix

Weeds Controlled

burdock, common	kochia*	shepherd's-purse
canola, volunteer	lamb's-quarters	stinkweed
cleavers (1-8 whorls)	mustard (except dog & green tansy)	sunflower, annual
cocklebur	pigweed, redroot (1-4 leaf)	vetch
flax, volunteer (1-12 cm)	prickly lettuce	wild radish
flixweed	ragweed, common	wild buckwheat (1-8 leaf)
hempnettle (2-6 leaf)		

Weeds Suppressed

green smartweed
stork's-bill (1-8 leaf)

*Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.

Pre-Harvest/Grazing Intervals (Audible B Herbicide + MCPA ester herbicide)

- Allow 7 days between application and grazing/cutting for hay.
- Do not harvest the treated mature crop within 60 days after application.

TANK-MIX COMBINATIONS - AUDIBLE B HERBICIDE + MCPA ESTER HERBICIDE + OTHER HERBICIDES (Do not apply to oats)

Tank mixtures of Audible B Herbicide+ MCPA ester herbicide with other herbicides will provide control of additional weeds. For products which may be tank mixed with Audible B Herbicide + MCPA ester herbicide and the additional weeds controlled see following table. Follow all precautions, minimal interval to harvest and directions for use (including timing of application) on the Audible B Herbicide and tank-mix partner labels.

Tank-Mix Combinations with Audible B Herbicide (0.31 L/ha) + MCPA LV500 Herbicide (1.12 L/ha) or MCPA Ester 600 Herbicide (0.93 L/ha)

Herbicide Tank-Mix Partner	Crops Registered	Rate/ha	Adjuvant Rate	Additional Weeds
Liquid Achieve SC or Achieve Liquid*	spring wheat, durum wheat, winter wheat, spring barley	0.5 L/ha	Turbocharge D or Turbocharge 0.5 L/ha (0.5% v/v)	green foxtail wild oats
Assert 300 SC	spring wheat, durum wheat, spring barley	1.3 L	Refer to Assert 300 SC label	wild oats (1-3 leaf)
		1.6 L		wild oats (4 leaf)
Everest 70 WDG or Everest Solupak 70 DF	spring wheat, durum wheat	43 g/ha	Refer to Everest label	green foxtail wild oats
Puma 120 Super	spring wheat, durum wheat, spring barley	0.77 L	None required	wild oats

*MCPA ester can be tank mixed at rates up to a maximum of 560 g ae/ha with this tank-mix combination.

TANK-MIX COMBINATION - AUDIBLE B HERBICIDE +SIMPLICITY™ (Do not apply to oats)

For control of a wide spectrum of broadleaf and grass weeds (see following table) in spring wheat, durum wheat and winter wheat, apply Audible B Herbicide tank mixed with Simplicity. Apply post-emergence when weeds are actively growing: grass weeds up to the 4 leaf, 2 tiller stage and broadleaf weeds as indicated in the following table. Best results are obtained from applications made to seedling weeds. Apply to actively growing spring or durum wheat from the 3 leaf stage to the first node stage. Apply to winter wheat in the 2-7 leaf, 4 tiller stage with a spring application. Extreme growing conditions such as

2022-3214
2022-08-04

drought or near freezing temperature prior to, at, or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Under conditions of low crop and high weed density, control may be reduced.

Mixing Instructions

Fill the sprayer tank ½ full of water and start sprayer tank agitation. Add the required amount of Simplicity. Add the correct amount of Audible B Herbicide and continue to agitate.

Preharvest/Grazing Intervals (Audible B Herbicide and Simplicity)

- Livestock may be grazed on treated crops 7 days following application.
- Do not harvest the treated crop within 60 days after application.

Rates of Application

I. Low Rate: To control or suppress the following weeds apply a uniform spray containing Audible B Herbicide at 0.21 L/ha tank mixed with Simplicity at 375 mL/ha.

Weeds Controlled

cleavers (1-6 whorls)

wild oats (up to 4 leaf, 2 tiller) *

*Under low wild oat populations (<75 plants/m²)

II. High Rate: To control the above weeds plus the additional weeds listed below, apply a uniform spray containing Audible B Herbicide at 0.31 L/ha with Simplicity at 500 mL/ha.

Additional Weeds Controlled

barnyard grass (1-5 leaf)	kochia* (2-8 leaf)
chickweed, common (up to 10 cm)	red root pigweed (1-8 leaf)
cleavers (1-8 whorls)	smartweed (lady's thumb, 1-5 leaf)
foxtail, yellow (1-5 leaf)	volunteer canola** (1-6 leaf)
hemp nettle (1-8 leaf)	volunteer flax (1-12 cm)
Japanese brome (1-6 leaf)	

*Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.

** will not control volunteer Imidazolinone-tolerant canola

Additional Weeds Suppressed

brome, downy (2-6 leaf, 2 tillers)	storks-bill (1-8 leaf)
foxtail, green (1-5 leaf)	wild buckwheat (1-4 leaf)

TANK-MIX COMBINATIONS –AUDIBLE B HERBICIDE + SIMPLICITY + OTHER HERBICIDES (Do not apply to oats)

When tank mixing Audible B Herbicide + Simplicity at either the low or high rates, the addition of other herbicides will provide control of additional weeds. For products which may be tank mixed with Audible B Herbicide + Simplicity and the additional weeds controlled see following table. Follow all precautions, minimal interval to harvest and directions for use on the Audible B Herbicide and tank-mix partner labels.

2022-3214
2022-08-04

Tank-Mix Combinations with Audible B Herbicide (0.21 L/ha) + Simplicity Herbicide (375 mL/ha) or Audible B Herbicide (0.31 L/ha) + Simplicity herbicide (500 mL/ha)

Herbicide Tank-Mix Partner	Crops Registered	Rate of Tank-Mix Partner	Additional Weeds Controlled
2,4-D ester*	spring wheat, durum wheat, winter wheat	Up to 420 g ae/ha	Susceptible weeds such as: mustard (except dog and green tansy), flixweed, bluebur, burdock, cocklebur, stinkweed, goatsbeard, prickly lettuce, lamb's-quarters, plantain, ragweeds, Russian thistle, shepherd's purse, annual sunflower, sweet clover, volunteer canola, wild buckwheat, wild radish * Consult 2,4-D ester labels for weed stages and rates
MCPA ester*	spring wheat, durum wheat, winter wheat	Up to 560 g ae/ha	Burdock (before the 4-leaf stage), cocklebur, flixweed, plantain, lamb's-quarters, mustards (except dog and tansy), prickly lettuce, ragweeds, Russian pigweed, shepherd's-purse, stinkweed, vetch, wild buckwheat, wild radish, wild (annual) sunflower. * Consult MCPA ester labels for weed stages and rates

BUFFER ZONES

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands) and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop		Buffer Zones (metres) Required for the Protection of:				
			Freshwater Habitat of Depths:		Estuarine/marine Habitat of Depths:		Terrestrial habitat
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	Spring wheat, durum wheat, winter wheat, spring barley and oats		1	0	1	1	3
Aerial	Spring wheat, durum wheat, winter wheat, spring barley and oats	Fixed Wing	5	0	1	1	95
		Rotary Wing	3	0	1	1	80

2022-3214
2022-08-04

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator on the Pesticides portion of the Canada.ca website

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, Audible B Herbicide is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to Audible B Herbicide and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of Audible B Herbicide or other Group 4 herbicide within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
 - Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
 - Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
 - Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
 - Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
 - Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Albaugh LLC at 1-800-247-8013.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

TMTrademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners.

All other products listed are registered trademarks of their respective companies.

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

<container label>

[indicates optional text]

Group	2	Herbicide
-------	---	-----------

DRAFT

DRY FLOWABLE

[FOR USE ON WHEAT (SPRING, WINTER OR DURUM), BARLEY, OATS AND SULFONYLUREA (SU)
TOLERANT CANOLA]

COMMERCIAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

ACTIVE INGREDIENT: Thifensulfuron methyl 50% Tribenuron methyl 25%

WARNING – EYE AND SKIN IRRITANT

REGISTRATION. NO.: 31904

PEST CONTROL PRODUCTS ACT

NET CONTENTS: 0.02 kg – 13.6 kg

Albaugh LLC
1525 NE 36th Street
Ankeny, IA 50021 USA
1-800-247-8013

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- Avoid breathing spray mist. Use with adequate ventilation.
- Avoid contact with skin, eyes and clothing.
- Do not contaminate any body of water.
- May be harmful if swallowed, inhaled or absorbed through skin.
- Causes eye and skin irritation.
- When using, do not eat, drink or smoke. Wash skin thoroughly with soap and water after handling.
- Remove contaminated clothing immediately after use. Store and wash contaminated clothing separately from household laundry before re-use.
- Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves and socks and chemical-resistant footwear during mixing, loading, application, clean-up and repair. In addition, wear a face shield during mixing and loading. Chemical-resistant gloves are not required while operating groundboom sprayers.
- **TOXIC** to non-target terrestrial plants and aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

FIRST AID:

If in eyes:

Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If swallowed:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If inhaled:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or the product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call CHEMTREC 1-800-424-9300 (24 hours).

TOXICOLOGICAL INFORMATION: Treat symptomatically.

STORAGE:

Store this product away from food or feed. Not for use or storage in or around the home. Keep container closed.

DISPOSAL:

Recyclable Containers

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

If this is a refillable container:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

If this is a returnable container:

Do not reuse this container for any other purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

Group	2	Herbicide
-------	---	-----------

booklet

DRAFT

DRY FLOWABLE

[FOR USE ON WHEAT (SPRING, WINTER OR DURUM), BARLEY, OATS AND SULFONYLUREA (SU)
TOLERANT CANOLA]

COMMERCIAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

ACTIVE INGREDIENT: Thifensulfuron methyl 50% Tribenuron methyl 25%

WARNING – EYE AND SKIN IRRITANT

REGISTRATION. NO.: 31904

PEST CONTROL PRODUCTS ACT

NET CONTENTS: 0.02 kg – 13.6 kg

Albaugh LLC
1525 NE 36th Street
Ankeny, IA 50021 USA
1-800-247-8013

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- Avoid breathing spray mist. Use with adequate ventilation.
- Avoid contact with skin, eyes and clothing.
- Do not contaminate any body of water.
- May be harmful if swallowed, inhaled or absorbed through skin.
- Causes eye and skin irritation.
- When using, do not eat, drink or smoke. Wash skin thoroughly with soap and water after handling.
- Remove contaminated clothing immediately after use. Store and wash contaminated clothing separately from household laundry before re-use.
- Do not apply by air.
- Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves and socks and chemical-resistant footwear during mixing, loading, application, clean-up and repair. In addition, wear a face shield during mixing and loading. Chemical-resistant gloves are not required while operating groundboom sprayers.
- **Do not** enter or allow entry into treated areas for 12 hours after application.

FIRST AID:

If in eyes:

Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If swallowed:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If inhaled:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or the product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call CHEMTREC 1-800-424-9300 (24 hours).

TOXICOLOGICAL INFORMATION: Treat symptomatically.

IMPORTANT

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, recreational grounds or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water, including irrigation water that may be used on other crops.

DO NOT apply by air.

Carefully observe sprayer clean-up instructions, as spray tank residues may damage crops other than wheat, barley, oats or SU Canola™.

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

APPLY WITH A SURFACTANT REGISTERED FOR USE WITH DRAFT

GENERAL INFORMATION

DRAFT is recommended for selective postemergence control or suppression of broadleaf weeds in wheat (spring, winter or durum), barley, oats and SU tolerant canola. DRAFT is a 75% active ingredient herbicide formulated as a dry flowable dispersible granule. DRAFT is to be mixed with water and a recommended surfactant and applied as a uniform broadcast spray. It is noncorrosive, nonflammable, nonvolatile and does not freeze.

DRAFT MUST be applied with a recommended surfactant, early postemergence, to the main flush of actively growing broadleaf weeds. Recommended surfactants are AGRAL* 90, AG-SURF*, HIACTIVATE NON-IONIC LIQUID SPREADER/ACTIVATOR and CITOWETT® PLUS.

DRAFT is a Group 2 herbicide and may only suppress or not control Group 2 resistant weeds.

Warm, moist growing conditions promote active weed growth and enhance the activity of DRAFT by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and regrowth may occur. APPLY DRAFT WITH GROUND EQUIPMENT ONLY. For best results, ensure thorough spray coverage of target weeds. See "APPLICATION INSTRUCTIONS" section of this label for complete use details.

DRAFT stops growth of susceptible weeds rapidly. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 3 weeks after application, depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent on weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.

The following crops may be planted two months after an application of DRAFT: canola, flax, lentils and alfalfa.

DIRECTIONS FOR USE

- DO NOT re-enter treated fields until 12 hours after application.
- Apply only when the potential for drift to areas of human habitation or areas of human activity (such as houses, cottages, schools and recreational areas) is minimal. Take into consideration wind speed, wind direction, temperature inversion, application equipment and sprayer settings.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application:

- **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty.
- **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Spray Buffer zones:

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, pastures, rangelands, riparian areas, and shrublands), sensitive

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of Application	Spray Buffer zone (metres) required for the protection of:		
	Terrestrial habitat	Aquatic habitats of water depths:	
		<1m	>1m
Field sprayer	15	1	0

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

ENVIRONMENTAL PRECAUTIONS:

- **TOXIC** to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under **DIRECTIONS FOR USE**.
- To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil or clay.
- Avoid application of this product when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

SULFONYLUREA TOLERANT CANOLA

(FOR USE ONLY ON SULFONYLUREA TOLERANT CANOLA THAT CONTAINS THE CIBUS SU CANOLA™ TRAIT)

APPLICATION TIMING

Apply 20 grams of DRAFT per hectare with a surfactant which is recommended for use with DRAFT at 2.0 litres per 1,000 litres of spray volume (see "Mixing Instructions" section of this label) when SU Canola™ is at the 2 to 5 leaf stage of development but prior to the beginning of bolting for the control or suppression of weeds listed below. For best results apply to emerged, young, actively growing weeds that are less than 10 cm tall or across. Thorough coverage of target weeds is essential.

RESTRICTIONS:

Only for use on canola that contains the Cibus sulfonylurea herbicide tolerant trait (SU Canola™ Trait). Do not apply to non-sulfonylurea tolerant canola as severe crop injury or death of the plants may occur. Do not exceed more than 20 g of product per hectare applied to any one crop of SU tolerant Canola (Cibus SU Canola™) during one growing season.

WEEDS CONTROLLED:

Annual smartweed (green smartweed, lady's thumb)	Narrow-leaved hawk's-beard
Ball mustard	Redroot pigweed
Chickweed ²	Russian thistle
Corn spurry	Shepherd's purse
Cow cockle	Stinkweed
Flixweed	Tartary buckwheat
Groundsel, common	Volunteer rapeseed ⁶
Hemp-nettle	Volunteer sunflower
Kochia	Wild buckwheat ³
Lamb's-quarters	Wild mustard

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

WEEDS SUPPRESSED:

Canada thistle¹

Cleavers (1 to 3 whorls)

Round-leaved mallow⁴

Scentless chamomile

Sow thistle¹

Stork's-bill (2-6 leaf)

Toadflax⁵

Weed suppression is a visual reduction in weed competition (reduced population and/or vigour) as compared to an untreated check. Degree of suppression will vary with size of weed and environmental conditions prior to and following treatment.

Specific Weed Recommendations:

1. Canada thistle, Sow thistle: Apply DRAFT plus recommended surfactant postemergence when the majority of the thistles have emerged and are actively growing. For best top growth control, apply when thistles are no larger than 15 cm tall, before bud stage, and before crop canopy prevents thorough coverage of thistles. A single application will effectively inhibit the ability of emerged thistles to compete with the crop. Later emerging thistles will not be controlled.

2. Chickweed: Apply DRAFT plus recommended surfactant when the chickweed is small (1-6 leaf) and actively growing but before crop canopy prevents thorough coverage of weeds. Chickweed emerging after application will not be controlled.

3. Wild Buckwheat: Apply DRAFT plus surfactant to actively growing wild buckwheat in the cotyledon - 3 leaf stage. Under stress conditions such as dry weather control may be reduced. Large plants may regrow after treatment.

4. Round-leaved Mallow: Apply DRAFT plus a surfactant to actively growing round leaved mallow in the 2-6 leaf stage (10-12 cm in height). DRAFT will keep mallow stunted but may not reduce the overall populations.

5. Toadflax: Apply DRAFT plus recommended surfactant when toadflax is no taller than 15 cm in height. A control program for this weed includes both frequent tillage and chemical application.

6. Volunteer rapeseed: DRAFT alone will not control imazamox and imazethapyr tolerant canola, (e.g. Clearfield* canola)

NOTE: Avoid making applications to weeds when rainfall is threatening. Rainfall within 4 hours will result in reduced weed control of DRAFT.

Tank Mixtures

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Albaugh at 1-800- 247-8013 for information before mixing any pesticide or fertilizer that is not specifically recommended on this label.

DRAFT + LONTREL* 360 or CT MIX 360

For Use on Sulfonylurea Tolerant Canola (for use only on sulfonylurea tolerant canola that contains the Cibus SU Canola™ trait).

APPLY:

DRAFT	at 20 grams per hectare
plus	

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

LONTREL* 360 or CT MIX 360	at 0.21 litres per hectare
plus	
a recommended surfactant	at 2 litres per 1000 litres of spray volume

See "Mixing Instructions" section of the label for proper mixing/application instructions. Ensure that DRAFT is completely in suspension in the spray tank before adding LONTREL* 360 or CT MIX 360.

Apply the tankmix from the full 2-leaf stage to the 5-leaf stage of SU tolerant canola. Consult the LONTREL* 360 or CT MIX 360 label for additional application instructions and use precautions.

WEEDS CONTROLLED:

Canada Thistle (seasonal control) †

Wild Buckwheat††

PLUS: All other broadleaf weeds controlled or suppressed by DRAFT alone at 20 grams/ha.

†Canada Thistle (seasonal control): Apply DRAFT plus LONTREL* 360 or CT MIX 360 plus a recommended surfactant postemergent, after all the thistles have emerged, and are actively growing. For best control, apply when thistles are no larger than 15 cm tall, before bud stage and before crop canopy prevents thorough coverage of thistles. Later emerging thistles will not be controlled.

†† Wild Buckwheat: Apply DRAFT plus LONTREL* 360 or CT MIX 360 plus a recommended surfactant to actively growing wild buckwheat in the cotyledon to 5-leaf stage. Under stress conditions such as dry weather, control may be reduced. Large plants may regrow after treatment.

DRAFT HERBICIDE + YUMA GL LIQUID EC HERBICIDE

For use on Sulfonylurea Tolerant Canola (for use only on sulfonylurea canola that contains the Cibus SU Canola™ trait)

For the control of annual grasses as well as broadleaf weeds, apply a tank mixture of:

DRAFT HERBICIDE at 20 grams per hectare
plus
YUMA GL LIQUID EC HERBICIDE at 0.38 - 0.5 litres per hectare
plus
MERGE at 0.5 % v/v or 5 litres per 1000 litres spray solution

Apply DRAFT HERBICIDE plus YUMA GL LIQUID EC HERBICIDE tank mixture when the sulfonylurea tolerant canola (Cibus SU Canola™ trait) is in the 3- to 5-leaf stage of growth but prior to beginning of bolting. Application of the DRAFT HERBICIDE plus YUMA GL LIQUID EC HERBICIDE tank mixture prior to the 3-leaf stage of SU tolerant canola could result in crop injury.

This tankmix will control the above broadleaf weeds as well as the grass weeds listed below:

RATE	WEEDS CONTROLLED	WEED LEAF STAGE
0.38 or 0.5 L/ha	Barnyard grass	2 leaf to early tillering*
	Green foxtail	2 leaf to early tillering
	Volunteer barley	2 leaf to early tillering
	Volunteer corn	2-6 leaf
	Volunteer oats	2 leaf to early tillering**

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

	Volunteer wheat	2 leaf to early tillering
		1-5 leaf without tillers (0.38 L/ha)
	Wild oats	1-5 leaf to early tillering (up to 2 tillers)** (0.5 L/ha)
	WEEDS SUPPRESSED	WEED LEAF STAGE
	Quackgrass	2-6 leaf*

* Use the higher rate of YUMA GL LIQUID EC HERBICIDE (0.5 L/ha) for control of barnyard grass, as well as suppression of quackgrass.

** Best results on volunteer/wild oats if application is made before tillering begins.

NOTE:

Consult the YUMA GL LIQUID EC HERBICIDE label for additional instructions regarding grassy weed size and/or height at time of application as well as other use precautions, and restrictions.

DRAFT HERBICIDE + ARROW 240 EC HERBICIDE

For use on Sulfonylurea Tolerant Canola (for use only on sulfonylurea tolerant canola that contains the Cibus SU Canola™ trait)

For the control of annual grasses as well as broadleaf weeds, apply a tank mixture of:

DRAFT HERBICIDE at 20 grams per hectare

plus

ARROW 240 EC HERBICIDE

at

0.19 litres per hectare

plus

ADAMA X-ACT

at

0.5 % v/v or 5 litres per 1000 litres spray

solution

Apply DRAFT HERBICIDE plus ARROW 240 EC tank mixture when the sulfonylurea tolerant canola (Cibus SU Canola™ trait) is in the 3- to 5-leaf stage of growth but prior to beginning of bolting. Application of the DRAFT HERBICIDE plus ARROW 240 EC tank mixture prior to the 3-leaf stage of SU tolerant canola could result in crop injury.

This tankmix will control the above broadleaf weeds as well as the grass weeds listed below:

Grass Species	Leaf Stage	Application Rate of ARROW 240 EC	Application Rate of X-ACT Adjuvant
Wild Oats Volunteer cereals (wheat, barley, oats) Foxtail (green, yellow) Persian darnel Crabgrass (smooth and large) Proso millet Witchgrass Fall panicum Barnyard grass Volunteer corn Volunteer canary grass Quackgrass suppression	2 to 6	0.19 L/ha	0.5% v/v

NOTE:

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

Consult the ARROW 240 EC label for additional instructions regarding grassy weed size and/or height at time of application as well as other use precautions, and restrictions.

DRAFT Alone

Wheat (Spring, winter and durum), Barley and Oats

Apply 20 grams of DRAFT per hectare with a surfactant which is recommended for use with DRAFT at 2.0 litres per 1,000 litres of spray volume (see "Mixing Instructions" section of this label). Apply postemergence when the crop is in the two-leaf to flag leaf (shot blade) stage of growth, (prior to head emergence) but before the crop canopy will prevent thorough coverage of target weeds. For best results apply to emerged, young, actively growing weeds that are less than 10 cm tall or across. Thorough coverage of target weeds is essential.

WEEDS CONTROLLED:

Annual smartweed (green smartweed, lady's thumb)	Narrow-leaved hawk's-beard
Ball mustard	Redroot pigweed
Chickweed ²	Russian thistle
Corn spurry	Shepherd's purse
Cow cockle	Stinkweed
Flixweed	Tartary buckwheat
Groundsel, common	Volunteer rapeseed ⁶
Hemp-nettle	Volunteer sunflower
Kochia	Wild buckwheat ³
Lamb's-quarters	Wild mustard

WEEDS SUPPRESSED:

Canada thistle¹
Cleavers (1 to 3 whorls)
Round-leaved mallow⁴
Scentless chamomile
Sow thistle¹
Stork's-bill (2-6 leaf) Toadflax⁵

Weed suppression is a visual reduction in weed competition (reduced population and/or vigour) as compared to an untreated check. Degree of suppression will vary with size of weed and environmental conditions prior to and following treatment.

Specific Weed Recommendations:

1. Canada thistle, Sow thistle: Apply DRAFT plus recommended surfactant postemergence when the majority of the thistles have emerged and are actively growing. For best top growth control, apply when thistles are no larger than 15 cm tall, before bud stage, and before crop canopy prevents thorough coverage of thistles. A single application will effectively inhibit the ability of emerged thistles to compete with the crop. Later emerging thistles will not be controlled.
2. Chickweed: Apply DRAFT plus recommended surfactant when the chickweed is small (1-6 leaf) and actively growing but before crop canopy prevents thorough coverage of weeds. Chickweed emerging after application will not be controlled.
3. Wild Buckwheat: Apply DRAFT plus surfactant to actively growing wild buckwheat in the cotyledon - 3 leaf stage. Under stress conditions such as dry weather control may be reduced. Large plants may regrow after treatment.

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

4. Round-leaved Mallow: Apply DRAFT plus a surfactant to actively growing round leaved mallow in the 2-6 leaf stage (10-12 cm in height). DRAFT will keep mallow stunted but may not reduce the overall populations.
5. Toadflax: Apply DRAFT plus recommended surfactant when toadflax is no taller than 15 cm in height. A control program for this weed includes both frequent tillage and chemical application.
6. Volunteer rapeseed: DRAFT alone will not control imazamox and imazethapyr tolerant canola, (e.g. Clearfield* canola)

NOTE:

Avoid making applications to weeds when rainfall is threatening. Rainfall within 4 hours will result in reduced weed control of DRAFT.

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

TANK MIXES WITH DRAFT

DRAFT PLUS MCPA

Wheat (Spring, winter and durum), Barley and Oats

Apply DRAFT at 20 grams per hectare plus MCPA Amine or Ester 500 at 0.7 - 1.1 litres per hectare (or acid equivalent to deliver 350 - 560 grams active ingredient per hectare) plus a recommended surfactant at 2 litres per 1,000 litres of spray volume (see "Mixing Instructions" section of the label). Apply the tank mix from the full 3-leaf stage to just before the flag leaf stage of wheat (spring, durum, winter), barley and oats. This tank mix will control:

Annual smartweed [†]	Prickly lettuce
Ball mustard	Ragweed ^{††}
Burdocks (seedling)	Redroot pigweed
Chickweed	Russian pigweed ^{††}
Cocklebur ^{††}	Russian thistle
Corn spurry	Shepherd's purse
Cow cockle	Stinkweed
False ragweed ^{††}	Tartary buckwheat
Flixweed	Tumble mustard
Giant ragweed ^{††}	Volunteer rapeseed ^{†††}
Hare's-ear mustard	Wild buckwheat
Hempnettle	Plaintain (common)
Indian mustard	Wild mustard
Kochia ^{††}	Wild radish
Lamb's-quarters	Wormseed mustard

[†] Green smartweed and lady's thumb

^{††} Use a minimum of 400 grams MCPA active ingredient per hectare.

^{†††} Including imazamox and imazethapyr tolerant canola (e.g., Clearfield* canola).

WEEDS SUPPRESSED:

Canada thistle

Round-leaved mallow

Scentless chamomile

Sow thistle

Stork's-bill (2-6 leaf) (MCPA Ester only)

Follow "Specific Weed Recommendations" for Canada and sow thistle, wild buckwheat, round-leaved mallow and chickweed.

WEEDS CONTROLLED BY DRAFT PLUS MCPA (0.56 L/HA)

Weeds controlled or suppressed by DRAFT alone plus volunteer imazamox and imazethapyr tolerant canola (e.g., Clearfield* canola) (2-4 leaf stage).

Consult the MCPA label for additional application instructions and use precautions .

DRAFT PLUS 2,4-D

Wheat or Barley –Do Not Use On Oats

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

Apply DRAFT at 20 grams per hectare tank mixed with 2,4-D and a recommended surfactant. Rates and formulation of 2,4-D include:

1. 2,4-D Amine 500 at 0.84 - 1.1 litres per hectare or
2. 2,4-D Ester LV700 at 0.6 - 0.8 litres per hectare or
3. 2,4-D Ester LV600 at 0.7 - 0.9 litres per hectare or
4. Acid equivalent.

A recommended surfactant at 2 litres per 1,000 litres of spray volume must be added (see "Mixing Instructions" section of the label). Apply the tank mix from the full 3-leaf stage to just before the flag leaf stage of wheat (spring, durum, winter) and barley. DO NOT USE ON OATS.

WEEDS CONTROLLED:

Annual smartweed †	Prickly lettuce
Ball mustard	Redroot pigweed
Burdocks (seedling)	Russian pigweed
Chickweed	Russian thistle
Cocklebur	Shepherd's purse
Common ragweed	Stinkweed
Corn spurry	Stork's-bill
Cow cockle	(2-6 leaf)
False ragweed	(2,4-D ester only)
Flixweed	Sweet clover
Giant ragweed	Tartary buckwheat
Hare's-ear mustard	Tumble mustard
Hemp-nettle	Volunteer rapeseed ††
Indian mustard	Wild buckwheat
Kochia	Wild mustard
Lamb's-quarters	Wild radish
Plantain (common)	Wormseed mustard

† green smartweed and lady's thumb

†† including imazamox and imazethapyr tolerant canola varieties (e.g., Clearfield* canola). Consult the 2,4-D label for appropriate use rates.

WEEDS SUPPRESSED:

Canada thistle
Round-leaved mallow
Scentless chamomile
Sow thistle
Toadflax (2,4-D Ester only)

Follow "Specific Weed Recommendations" for Canada and sow thistle, chickweed, wild buckwheat and round-leaved mallow.

Consult the 2,4-D label for additional application instructions and use precautions.

DRAFT PLUS ASSERT* 300 Herbicide

Spring Wheat and Spring Barley Only

For Sale for Use in the Prairie Provinces and Interior of British Columbia (Including the Peace River Region) Only

Apply DRAFT at 20 grams per hectare plus ASSERT* 300 Herbicide at 1.3 to 1.6 litres of product per

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

hectare for the control of broadleaf weeds controlled by DRAFT, and wild oats.

Use the low rate of ASSERT* 300 Herbicide when the majority of wild oats are in the 1 - 3 leaf stage and the higher rate if wild oats are in the 4-leaf stage. Application can be made from the 2-leaf up to and including the 6-leaf stage of the crop.

Consult the ASSERT* 300 Herbicide label for additional application instructions; use precautions and recropping information.

Use the spray preparation within 12 hours or product degradation may occur.

DRAFT + ASSERT* 300 Herbicide + MCPA ESTER 500

Spring Wheat and Spring Barley Only

For Sale for Use in the Prairie Provinces and Interior of British Columbia (Including the Peace River Region) Only

Apply DRAFT at 20 grams per hectare plus ASSERT* 300 Herbicide at 1.3 - 1.6 litres of product per hectare plus MCPA Ester 500 at 0.7 - 1.1 litres per hectare (or acid equivalent to deliver 350 - 560 grams active ingredient per hectare) for the control of broadleaf weeds controlled by DRAFT plus MCPA, and wild oats.

Apply at the 4-leaf to 6-leaf stage of the crop.

SOIL ZONES	ASSERT* 300 Herbicide RATE
Black and Grey Wooded	1.6 L/ha
Brown and Dark Brown	1.3 L/ha if the majority of the wild oats are in the 1-3 leaf stage. Apply Assert* 300

Consult the ASSERT* 300 Herbicide label for additional application instructions, use precautions and recropping information.

Use the spray preparation within 12 hours or product degradation may occur.

DRAFT + PUMA* SUPER

Spring Wheat (Including Durum) and Barley

For Sale for Use in the Prairie Provinces and Interior of British Columbia (Including the Peace River Region) Only

APPLY:

DRAFT	at 20 grams per hectare
Plus	
PUMA* SUPER	at 0.5 or 1.0 Litres per hectare

A non-ionic surfactant is NOT required when DRAFT is tankmixed with PUMA* SUPER.

Apply the DRAFT + PUMA* SUPER tankmix to spring and durum wheat at the 1 to 6-leaf plus 3 tillers stage or to barley at the 1 to 5-leaf plus 2 tillers stage. Apply the tankmix to annual grasses in the 1-6 leaf stage (up to emergence of third tiller). Consult the PUMA* SUPER label for additional information on appropriate weed staging, application instructions, environmental and use precautions.

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

WEEDS CONTROLLED:

Green Foxtail (PUMA* SUPER at 0.5 or 1.0 L/ha)
Wild Oats (PUMA* SUPER at 1.0 L/ha)
Barnyard Grass (PUMA* SUPER at 1.0 L/ha)

This tankmix will also control or suppress the broadleaf weeds listed for DRAFT alone at 20 grams/ha.

DRAFT + MCPA ESTER 500 + PUMA* SUPER

Spring Wheat (Including Durum) and Barley

For Sale for Use in the Prairie Provinces and Interior of British Columbia (Including the Peace River Region) Only

APPLY:

DRAFT	at 20 grams per hectare
Plus	
MCPA ESTER 500	at 0.84 Litres per hectare
Plus	
PUMA* SUPER	at 0.5 OR 1.0 Litres per hectare

A non-ionic surfactant is NOT required when DRAFT is tankmixed with MCPA ESTER 500 + PUMA* SUPER. Use ONLY MCPA ESTER 500 in this tankmix and only at the recommended rate or a decrease in wild oat control may be observed.

Apply the DRAFT + MCPA ESTER 500 + PUMA* SUPER tankmix to spring and durum wheat at the 1 to 6-leaf plus 3 tillers stage or to barley at the 1 to 5-leaf plus 2 tillers stage. Apply to annual grasses in the 1-6 leaf stage (up to emergence of third tiller). Consult the MCPA ESTER 500 and PUMA* SUPER labels for additional information on appropriate weed staging, application instructions, environmental and use precautions.

WEEDS CONTROLLED:

Green Foxtail (PUMA* SUPER at 0.5 or 1.0 L/ha)
Wild Oats (PUMA* SUPER at 1.0 L/ha)
Barnyard Grass (PUMA* SUPER at 1.0 L/ha)

This tankmix will also control or suppress the broadleaf weeds listed for DRAFT at 20 grams/ha + MCPA ESTER 500 at 0.84 Litres/ha.

DRAFT + PUMA* 120 SUPER

Spring Wheat (Including Durum) and Barley

For Sale for Use in the Prairie Provinces and Interior of British Columbia (Including the Peace River Region) Only

APPLY:

DRAFT	at 20 grams per hectare
Plus	
PUMA* 120 SUPER	at 770 mL/Ha or 385 mL/Ha

A non-ionic surfactant is NOT required when DRAFT is tank mixed with PUMA* 120 SUPER.

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

Apply the DRAFT + PUMA* 120 SUPER tankmix to spring and durum wheat at the 1 to 6-leaf plus 3 tillers stage or to barley at the 1 to 5-leaf plus 2 tillers stage. Apply the tankmix to annual grasses in the 1-6 leaf stage (up to emergence of third tiller). Consult the PUMA* 120 SUPER label for additional information on appropriate weed staging, application instructions, environmental and use precautions.

WEEDS CONTROLLED:

Green Foxtail (PUMA* 120 SUPER at 385 mL/ha)
Wild Oats, Green Foxtail, Yellow (PUMA* 120 SUPER at 770 mL/ha)
Foxtail, Barnyard Grass
Broadleaf weeds controlled or suppressed by DRAFT alone

DRAFT + PUMA* 120 SUPER + MCPA Ester 500

Spring Wheat (Including Durum) and Barley

For Sale for Use in the Prairie Provinces and Interior of British Columbia (Including the Peace River Region) Only

APPLY:

DRAFT	at 20 grams per hectare
	Plus
PUMA* 120 SUPER	at 770 mL/Ha or 385 mL/Ha
	Plus
MCPA Ester 500	0.84 or 0.56 L/Ha

A non-ionic surfactant is NOT required when DRAFT is tank mixed with PUMA* 120 SUPER and MCPA Ester 500.

Use ONLY MCPA ESTER 500 in this tank mix and only at the recommended rate or a decrease in wild oat control may be observed.

Apply the DRAFT + PUMA* 120 SUPER + MCPA Ester 500 tank mix to spring and durum wheat at the 1 to 6-leaf plus 3 tillers stage or to barley at the 1 to 5-leaf plus 2 tillers stage. Apply the tank mix to annual grasses in the 1-6 leaf stage (up to emergence of third tiller). Consult the MCPA Ester 500 and PUMA* 120 SUPER label for additional information on appropriate weed staging, application instructions, environmental and use precautions.

Green Foxtail (PUMA* 120 SUPER at 385 mL/ha)
Wild Oats, Green Foxtail, Yellow (PUMA* 120 SUPER at 770 mL/ha)
Foxtail, Barnyard Grass
Broadleaf weeds controlled or suppressed by DRAFT alone
Broadleaf weeds controlled by MCPA Ester 500 at 0.84 L/Ha

OR

Volunteer imazamox and imazethapyr tolerant canola (e.g., Clearfield* canola) (2-4 leaf stage) when MCPA Ester 500 is applied at 0.56 L/ha

DRAFT + LONTREL* 360 or CT MIX 360 + 2,4-D ESTER or MCPA ESTER

Spring Wheat (Excluding Durum) and Barley Only. For Use in Western Canada Only.

APPLY:

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

DRAFT	at 20 grams per hectare
plus	
LONTREL* 360 or CT MIX 360	at 0.21 litres per hectare
plus	
MCPA Ester (500 g/L)	at 0.84 litres per hectare
or	
2,4-D Ester (600 g/L)	at 0.7 litres per hectare

Add a recommended non-ionic surfactant such as AGRAL* 90, AG-SURF*, HIACTIVATE NON-IONIC LIQUID SPREADER/ACTIVATOR or CITOWETT® PLUS at a rate of 0.2% of the final spray volume (i.e., 2.0 litres per 1000 litres of spray solution).
See "Mixing Instructions" section of the label.

WEEDS CONTROLLED:

Canada thistle[†]
Lady's thumb
Perennial sow thistle
Stinkweed
Volunteer canola
Wild buckwheat
Wild mustard

[†] Provides seasonal control of Canada thistle

WEEDS SUPPRESSED:

Cleavers

PLUS: all other broadleaf weeds controlled or suppressed by DRAFT alone at 20 grams per hectare. Refer to the MCPA Ester or 2,4-D Ester labels for additional broadleaf weeds controlled at the rates recommended above. Consult the LONTREL* or CT MIX 360 Herbicide and 2,4-D Ester or MCPA Ester labels for additional application instructions, use precautions, and other label restrictions.

DO NOT USE ON OATS.

DRAFT + CURTAIL* M

Spring Wheat (excluding Durum) and Spring Barley Only.

For Sale for Use in the Prairie Provinces and Interior of British Columbia (Including the Peace River Region) Only

APPLY:

DRAFT	20 grams per hectare
plus	
CURTAIL* M	1.5 litres per hectare
plus	
a recommended surfactant	at 2 litres per 1,000 litres of spray volume

See "Mixing Instructions" section of the label for proper mixing/application instructions. Ensure that DRAFT is completely in suspension in the spray tank before adding CURTAIL*M.

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

Apply the tank mix from the full 3-leaf stage of spring wheat and spring barley. Consult the CURTAIL* M label for additional application instructions and use precautions.

WEEDS CONTROLLED:

Canada thistle[†]

Wild buckwheat^{††}

PLUS all other broadleaf weeds controlled or suppressed by DRAFT alone at 20 grams per hectare.

[†] Canada thistle: Apply DRAFT plus CURTAIL* M plus a recommended surfactant postemergence after all the thistles have emerged, and are actively growing. For best control, apply when thistles are no larger than 15 cm tall, before bud stage and before canopy prevents thorough coverage of thistles. Later emerging thistles will not be controlled.

^{††} Wild buckwheat: Apply DRAFT plus CURTAIL* M plus a recommended surfactant to actively growing wild buckwheat in the cotyledon to 5 leaf stage. Under stress conditions such as dry weather, control may be reduced. Large plants may regrow after treatment.

DRAFT + ACCORD* DF HERBICIDE

SPRING WHEAT (including Durum) AND SPRING BARLEY (grown for feed purposes only). Not Underseeded to Forages.

For Sale for Use in the Prairie Provinces and Interior of British Columbia (Including the Peace River Region) Only

APPLY:

DRAFT	at 20 grams per hectare
plus	
ACCORD* DF Herbicide	At 67 OR at 135-165 grams per hectare
plus	
MERGE* adjuvant	at 1.0% v/v (10 litres per 1000 litres spray volume)

A non-ionic surfactant is NOT required when DRAFT is tank mixed with ACCORD* DF Herbicide plus the MERGE* adjuvant.

Make application when crop is in the 2-5 leaf stage. Use the 165 grams per hectare rate of ACCORD* DF Herbicide when green foxtail is the target weed. Thorough coverage of foliage is important for consistent weed control. Do not apply this tank mixture more often than once every two years. Prior to use, consult the ACCORD* DF Herbicide label for additional rotational cropping restrictions, application instructions, use precautions and other restrictions. Do not graze the treated crops or cut for hay within 77 days of application.

WEEDS CONTROLLED WHEN ACCORD* DF HERBICIDE IS USED AT 135-165 GRAMS/HA	APPLICATION TIMING
Green Foxtail	1-5 leaf, 2 tillers
Barnyard Grass	1-5 leaf
Volunteer Flax	1-8 cm
Cleavers	1-4 whorls
Annual Sowthistle	1-4 leaf
PLUS: Weeds controlled or suppressed by DRAFT alone	

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

Refer to the ACCORD* DF Herbicide label for additional information on appropriate weed staging at application.

Do not graze the treated crops or cut for hay within 77 days (spring wheat, durum) or 80 days (spring barley) of application.

WEEDS CONTROLLED WHEN ACCORD* DF Herbicide IS USED AT 67 GRAMS/HA	APPLICATION TIMING
Cleavers	1-3 whorls
Annual Sowthistle	1-4 leaf
PLUS: Weeds controlled or suppressed by DRAFT alone	

DRAFT + ACCORD* DF HERBICIDE + PUMA* 120 SUPER

SPRING WHEAT (Including Durum) AND SPRING BARLEY (grown for feed purposes only). Not Underseeded to Forages.

For Sale for Use in the Prairie Provinces and Interior of British Columbia (Including the Peace River Region) Only

APPLY:

DRAFT	20 grams per hectare
Accord* DF Herbicide	67 grams per hectare
Puma* 120 Super	0.77 Litres per hectare

Merge* adjuvant is NOT required when DRAFT and Accord* DF Herbicide are tank mixed with Puma* 120 Super.

Apply the DRAFT + Accord* DF Herbicide + Puma* 120 Super tank mix to spring and durum wheat at the 2 to 5-leaf stage or to barley at the 2 to 4-leaf stage. Apply the tank mix to annual grasses in the 1-6 leaf stage (up to emergence of third tiller), or to cleavers in the 1-4 whorl. Consult the Accord* DF Herbicide and Puma* 120 Super labels for additional information on appropriate weed staging, application instructions, environmental and use precautions.

WEEDS CONTROLLED:

Weeds controlled or suppressed by DRAFT alone
Green Foxtail
Wild Oats
Barnyard Grass
Cleavers

Do not graze the treated crops or cut for hay within 77 days (spring wheat, durum) or 80 days (spring barley) of application.

DRAFT + ATTAIN* HERBICIDE TANK MIX

For Use on Spring Wheat (excluding Durum), and Spring Barley Only. Not Underseeded to Legumes. For Sale for Use in the Prairie Provinces and Interior of British Columbia (Including the Peace River Region) Only

APPLY:

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

DRAFT	at 20 grams per hectare
plus	
ATTAIN* Herbicide Tank Mix	Attain A (fluroxypyr) at 0.30 L/ha Attain B (2,4-D ester) at 0.50 L/ha
plus	
Adjuvants: AGRAL* 90, AG-SURF*, HIACTIVATE NON-IONIC LIQUID SPREADER/ACTIVATOR or CITOWETT*	at 2L/1000 litres of spray solution

Apply from the 4-leaf to flag-leaf stage of spring wheat or spring barley and seedling stage to 4-leaf/whorls of weed.

Mixing order is DRAFT, then Attain* A then Attain* B then adjuvant. Refer to MIXING DIRECTIONS section for general instructions. Use a spray volume of 100 L/ha.

Do not permit lactating dairy animals to graze fields within 7 days of application. Do not harvest forage or cut hay within 30 days after application. Withdraw meat animals from treated fields at least 3 days before slaughter. Do not harvest the treated mature crop within 60 days after application. Consult the Attain* Herbicide Tank Mix labels for additional application instructions and use precautions.

WEEDS CONTROLLED:

Weeds controlled or suppressed by DRAFT alone

Cleavers (excluding Group 2 resistant cleavers).

DRAFT + LONTREL* 360 or CT MIX 360

**For Use on Spring Wheat (excluding Durum), and Spring Barley Only
For Sale for Use in the Prairie Provinces and Interior of British Columbia (Including the Peace River Region) Only**

APPLY:

DRAFT	at 20 grams per hectare
plus	
LONTREL* 360 or CT MIX 360	at 0.21 litres per hectare
plus	
a recommended surfactant	at 2 litres per 1000 litres of spray volume

See "Mixing Instructions" section of the label for proper mixing/application instructions. Ensure that DRAFT is completely in suspension in the spray tank before adding LONTREL* 360 or CT MIX 360.

Apply the tankmix from the full 3 leaf stage to the flag leaf stage of spring wheat and barley. Consult the LONTREL* 360 or CT MIX 360 label for additional application instructions and use precautions.

WEEDS CONTROLLED:

Canada Thistle (seasonal control) †

Wild Buckwheat††

PLUS: All other broadleaf weeds controlled or suppressed by DRAFT alone at 20 grams/ha.

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

†Canada Thistle (seasonal control): Apply DRAFT plus LONTREL* 360 or CT MIX 360 plus a recommended surfactant postemergent, after all the thistles have emerged, and are actively growing. For best control, apply when thistles are no larger than 15 cm tall, before bud stage and before crop canopy prevents thorough coverage of thistles. Later emerging thistles will not be controlled.

†† Wild Buckwheat: Apply DRAFT plus LONTREL* 360 or CT MIX 360 plus a recommended surfactant to actively growing wild buckwheat in the cotyledon to 5-leaf stage. Under stress conditions such as dry weather, control may be reduced. Large plants may regrow after treatment.

DRAFT + BANVEL II* HERBICIDE

For Use on Spring Wheat (excluding Durum), and Spring Barley Only. Not Underseeded to Forages. For Sale for Use in the Prairie Provinces and Interior of British Columbia (Including the Peace River Region) Only

Apply from the 2-5 leaf stage of spring wheat or spring barley and seedling stage to 8 leaf stage of kochia.

DRAFT	20 grams/hectare
Banvel II* Herbicide	0.110 litres/hectare
Adjuvants: Agral*90, Ag-surf*, HiActivate Non-Ionic Liquid Spreader/Activator or Citowett* Plus	2 litres/1000 litres of spray solution

Mixing order is DRAFT then Banvel II* Herbicide then adjuvant. Refer to the MIXING DIRECTIONS section for general instructions. Use minimum volume of 55 L/ha.

Do not permit lactating dairy animals to graze fields within 7 days after application. Consult the Banvel II* Herbicide label for additional application instructions and use precautions.

WEEDS CONTROLLED:

Broadleaf weeds controlled or suppressed by DRAFT

Kochia, including Group 2 resistant kochia

DRAFT + 2,4-D AMINE OR ESTER + EVEREST 70 WDG

For Use on Spring Wheat (Including Durum) Only

For Sale for Use in the Prairie Provinces and Interior of British Columbia (Including the Peace River Region) Only

Apply one of the following tank mixes from the 1 to 6-leaf (1 to 4 leaves on main stem, plus 2 tillers) growth stage of the crop. For wild oats, green foxtail and volunteer tame oats, apply from the 1 to 6-leaf (1 to 4 leaves on main stem, plus 2 tillers):

TANK MIX COMPONENTS AND APPLICATION RATES				
Component 1	Component 2	Component 3	Component 4	WEEDS CONTROLLED - Broadleaf weeds controlled or suppressed by DRAFT alone plus:

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

DRAFT	20 g/ha	Everest* 70 WDG	43 g/ha	Agral * 90, HiActivate Non-Ionic Liquid Spreader/ Activator or Agsurf*	0.25% v/v			Wild oats, green foxtail, volunteer tame oats. Use 43 g/ha when wild oat infestations are greater than 100 plants/m ² or when plants are not actively growing due to poor environmental conditions.
DRAFT	20 g/ha	Everest* 70 WDG	43 g/ha	2,4-D Amine or Ester	up to 560 g ae/ha	Agral * 90, HiActivate Non-Ionic Liquid Spreader/A ctivator or Agsurf*	0.25 % v/v	Wild oats, green foxtail and volunteer tame oats. Use 43 g/ha when wild oat infestations are greater than 100 plants/m ² or when plants are not actively growing due to poor environmental conditions.
DRAFT	20 g/ha	Everest* 70 WDG	28.5 g/ha	2,4-D Amine or Ester	up to 560 g ae/ha	Agral * 90, HiActivate Non-Ionic Liquid Spreader/A ctivator or Agsurf*	0.25 % v/v	Wild oats and volunteer tame oats only. Use 28.5 g/ha when wild oat populations are below 100 plants/m ² and when plants are actively growing.
DRAFT	20 g/ha	Everest* 70 WDG	21.5 g/ha	2,4-D Amine or Ester	up to 560 g ae/ha	Agral * 90, HiActivate Non-Ionic Liquid Spreader/A ctivator or Agsurf*	0.25 % v/v	Green foxtail only

Mixing order is Everest 70 WDG then DRAFT then 2,4-D Amine or Ester then adjuvant. Refer to MIXING DIRECTIONS section for general instructions.

Observe minimum interval to harvest of 80 days after treatment. For black, dark brown and gray-wooded soil zones only, field peas may be grown the year following Everest* application in fields where precipitation has been normal or above normal (10 year average) during the growing season, and where organic matter content is above 4% and pH below 7.5. Re- crop spring wheat only in brown soil zone. Consult the 2,4-D Amine or Ester and Everest* 70 WDG labels for additional information on rotational cropping restrictions, application instructions, use precautions and other restrictions.

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

DRAFT PLUS SIMPLICITY* GoDRI* HERBICIDE

Spring Wheat (Including Durum) and Winter Wheat

APPLY:

DRAFT	at 20 grams per hectare
plus	
SIMPLICITY* GoDRI* Herbicide	at 70 grams per hectare
plus	
A non-ionic surfactant	at 0.25% v/v

WEEDS CONTROLLED:

barnyard grass (1-5 leaf)	hemp nettle (1-8 leaf)
brome, Japanese (1-6 leaf)	oats, wild (up to 4 leaf, 2 tiller)
canola, volunteer (1-6 leaf)	pigweed, redroot (1-8 leaf)
chickweed, common (up to 10 cm)	round-leaved mallow (up to 6 leaf stage, <10 cm in size)
cleavers (up to 6 whorl)	shepherd's purse (up to 30 cm tall)
corn spurry (up to 2 whorl stage,	smartweed (lady's-thumb, 1-5 leaf)
foxtail, yellow (1-5 leaf)	stinkweed (up to 30 cm tall)

WEEDS SUPPRESSED:

brome, downy (2-6 leaf, 4-tillers)	Persian darnel (1-leaf to 4-leaf, 2-tiller)
Canada thistle (up to 30 cm tall, prebud)	Russian thistle (up to 10 cm tall)
dandelion (seedlings and over-wintered rosettes less than or equal to 20 cm)	white cockle (up to the first flower stage, less than 20 cm in height)
foxtail, green (1-5 leaf)	wild buckwheat (1-4 leaf)

PLUS: all other broadleaf weeds controlled or suppressed by DRAFT alone at 20 grams per hectare.

DRAFT PLUS AXIAL* 100EC Herbicide

For Sale for Use in the Prairie Provinces and Interior of British Columbia (Including the Peace River Region) Only

For use on spring wheat and spring barley only

Post-emergent application from 2 to flag leaf. For annual grass weeds, apply from 2- to 6-leaf stage, up to emergence of 4th tiller.

TANK MIX COMPONENTS AND APPLICATION RATES:

Draft Herbicide at 20 grams/hectare with AXIAL* 100 EC Herbicide at 600 mL/hectare for wild oats, volunteer oats, green foxtail, yellow foxtail, volunteer canary seed, and proso millet and with ADIGOR* Adjuvant at 700 mL/hectare

MIXING: Ensure that Draft Herbicide is completely dissolved. Then add Axial* 100 EC Herbicide with reduced agitation, and finally Adigor adjuvant.

DO NOT USE ADDITIONAL NON-IONIC SURFACTANT.

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

Consult the Axial* 100EC Herbicide label for application instructions and use precautions.

WEEDS CONTROLLED:

Broadleaf weeds controlled or suppressed by Draft Herbicide alone, plus wild oats, volunteer oats, green foxtail, yellow foxtail, volunteer canary seed, and proso millet.

Do not harvest within 60 days for grain and straw and 30 days for hay.

Do not graze treated crop for 7 days after application.

Consult AXIAL* 100 EC Herbicide label for additional information on appropriate weed staging, application instructions, environmental hazards and use precautions.

DRAFT PLUS AUDIBLE B HERBICIDE

Apply post-emergence when weeds are actively growing, with best results on seedling weeds that are less than 10 cm in height. Only weeds emerged at the time of treatment will be controlled. Thorough covering of the target weeds is essential. The tank-mix can be applied to actively growing spring wheat, including durum, spring barley and oats from the 2-leaf crop growth stage up to and including initiation of stem elongation. **Do not apply to crops underseeded to legumes or grasses.**

Rates of Application: To control the following weeds apply a uniform spray containing 20 g/ha of Draft with 0.21 to 0.41 litres of Audible B Herbicide and 2 litres per 1000 L spray solution of non-ionic surfactant with a spray volume of 100 L/ha.

Weeds Controlled or Suppressed:

Weeds Controlled at 20 g/ha Draft and 0.21 L/ha of Audible B Herbicide:

annual smartweed (green smartweed, lady's thumb)	groundsel (common)	shepherd's purse
ball mustard	hemp-nettle	stinkweed
chickweed (1-6 leaf) ¹	kochia ²	tartary buckwheat
cleavers (1-4 whorls)	lamb's-quarters	volunteer rapeseed ³
corn spurry	narrow-leaved hawk's beard	volunteer sunflower
cow cockle	redroot pigweed	wild buckwheat ⁴
flixweed	Russian thistle	wild mustard

¹ Spray before canopy prevents thorough coverage of weeds. Chickweed emerging after application will not be controlled.

² Excluding Group 2 resistant kochia

³ The tank-mix will not control imazethapyr tolerant canola varieties (e.g. Clearfield varieties or other varieties with the Pursuit Smart trait)

⁴ Apply to actively growing wild buckwheat at the cotyledon to 3-leaf stage. Control may be reduced under environmental stress. Large plants may regrow after treatment.

Weeds Suppressed at 20 g/ha Draft and 0.21 L/ha of Audible B Herbicide:

Canada thistle ⁵	sow thistle ⁵
round leaved mallow (2-6 leaf) ⁶	stork's bill (2-6 leaf stage)
scentless chamomile	toadflax (up to 15 cm) ⁷

⁵ Apply when most of the thistles have emerged and are actively growing. Apply before bud stage and before crop canopy prevents thorough coverage of thistles, when thistles are no larger than 15 cm in

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

height for best top growth control. A single application will effectively inhibit the ability of emerged thistles to compete with the crop. Later emerging thistles will not be controlled.

⁶ Apply to actively growing round leaved mallow. Will keep mallow stunted, but may not be able to reduce overall populations.

⁷ A control program for this weed includes chemical application and frequent tillage.

Additional Weeds Controlled at 20 g/ha Draft and 0.31 L/ha of Audible B Herbicide:

kochia (2-8 leaf; including group 2 resistant biotypes)

volunteer flax (1-12 cm)

Additional Weeds Controlled and Suppressed at 20 g/Ha Draft and 0.41 L/ha Audible B Herbicide:

Weeds Controlled

round leaved mallow (1-6 leaf)

Weeds Suppressed

chickweed (up to 8 cm; including group 2 resistant biotypes)

MIXING INSTRUCTIONS:

1. Select a spray volume that will ensure thorough coverage and uniform spray pattern. Use a minimum of 55-L spray volume per hectare. If tankmixing with other herbicides, always consult the label of the tank mix partner(s) for minimum spray volume requirements and apply the tank mixture using a water volume recommended for all products.
2. Fill the spray tank 1/4 – 1/3 full with fresh water and start agitation.
3. Add the required amount of DRAFT to the tank. Make sure all of the DRAFT is in suspension before proceeding to the next step. Continuous agitation is recommended throughout the entire mixing procedure to ensure products are completely dissolved and/or in suspension.
4. If using DRAFT alone, a recommended surfactant such as AGRAL* 90, AG-SURF*, HIACTIVATE NON-IONIC LIQUID SPREADER/ACTIVATOR or CITOWETT® PLUS must be added to the spray mixture at a rate of 2 litres per 1000 litres of spray solution.
5. Specific Tank Mix Instructions:
If tankmixing with MCPA, 2,4-D, CURTAIL* M, LONTREL* 360 or CT Mix 360, add the required amount of MCPA, 2,4-D, CURTAIL* M, LONTREL* 360 or CT Mix 360 to the tank while maintaining agitation. Once the herbicide is completely in suspension, add a recommended surfactant such as AGRAL* 90, AG-SURF*, HIACTIVATE NON-IONIC LIQUID SPREADER/ACTIVATOR or CITOWETT® PLUS at 2 L per 1000 L spray solution. Anti-foaming agents may be required.

If tankmixing with PUMA* SUPER or MCPA + PUMA* SUPER, add the required amount of PUMA* SUPER and/or MCPA while maintaining agitation. DO NOT ADD A SURFACTANT.

If tank mixing with ASSERT* 300 Herbicide or ASSERT*300 Herbicide + MCPA: once DRAFT is completely in suspension, add the required amount of Spray Water pH Adjuster and agitate until dissolved. Add the required amount of ASSERT* 300 Herbicide while maintaining agitation. If adding MCPA, add it to the tank last. Addition of surfactant is not required with this mixture.

When mixing with LONTREL* 360 + 2,4-D Ester or MCPA Ester, mix DRAFT first in fresh water, then add the required amount of LONTREL* 360 or CT Mix 360 followed by MCPA or 2,4-D. Add the recommended non-ionic surfactant such as AGRAL* 90, AG-SURF*, HIACTIVATE NON-IONIC LIQUID SPREADER/ACTIVATOR or CITOWETT® PLUS last, at a rate of 2 litres per 1000 litres spray solution.

If tankmixing with YUMA GL LIQUID EC HERBICIDE add the required amount of YUMA GL LIQUID EC HERBICIDE after DRAFT is completely in suspension, and maintain agitation. Add the MERGE or AMIGO adjuvant at 0.5 %v/v. DO NOT ADD AN ADDITIONAL NON-IONIC

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04
SURFACTANT.

If tankmixing with ARROW 240 EC add the required amount of ARROW 240 EC after DRAFT is completely in suspension, and maintain agitation. Add the ADAMA X-ACT adjuvant at 0.5 %v/v. DO NOT ADD AN ADDITIONAL NON-IONIC SURFACTANT.

When tankmixing with ACCORD* DF Herbicide, add the required amount of DRAFT and ACCORD* to the tank. When these products have completely dissolved and are in suspension, add the MERGE* adjuvant at the appropriate rate last. Do not add an additional non-ionic surfactant.

6. Continue to fill tank and maintain agitation for at least 5 minutes to ensure thorough mixing.
7. On repeat loads, ensure that the amount of spray solution left in the tank from the previous load is less than 10% of the volume about to be mixed.
8. Use the spray preparation of DRAFT within 24 hours or product degradation may occur. Use spray preparation containing ASSERT* 300 Herbicide within 12 hours or product degradation may occur. If spray solution is left standing without agitation, thoroughly agitate and re-suspend any settled DRAFT before resuming the spraying operation.

The volumetric measuring guide to measure DRAFT is to be used only as a guide. For precise measurement use scales calibrated in grams.

APPLICATION INSTRUCTIONS:

Apply the spray mixture uniformly with properly calibrated ground equipment only. Ensure thorough coverage and a uniform spray pattern. Flat fan nozzles are recommended (minimum 55 litres spray volume per hectare). Use 50 mesh filter screens or larger (metal or nylon). A spray pressure of 210-275 kPa is recommended. Continuous agitation is required to keep DRAFT in suspension. Apply uniformly. Avoid overlapping and shut off spray boom while starting, turning, slowing or stopping to prevent crop injury from over application.

SPRAYER CLEANUP:

To avoid subsequent injury to crops other than wheat, barley or oats, immediately after spraying and prior to spraying other crops, thoroughly remove all traces of DRAFT from mixing and spray equipment as follows:

1. Drain tank; then flush tank, boom and hoses with clean water for a minimum of ten minutes. Visually inspect tank to ensure removal of all visible residues of DRAFT. If necessary, repeat Step 1. DO NOT CLEAN SPRAYER NEAR WELL OR WATER SOURCE OR NEAR TO DESIRABLE VEGETATION.
2. Fill the tank with clean water, then add 1 litre household AMMONIA (containing minimum of 3% ammonia) per 100 litres of water. Flush solution through boom and hoses, and then add more water to completely fill tank. Allow to sit for 15 minutes with agitation. Again flush the hoses, boom and nozzles with the cleaning solution and drain tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat Steps 2 and 3.
5. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing water through the hoses and boom.

CAUTION: Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odour which may cause eye, nose, throat, and lung irritation. Do not clean equipment in an enclosed area.

RESISTANCE-MANAGEMENT RECOMMENDATIONS:

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

For resistance management, DRAFT is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to DRAFT and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible rotate the used of DRAFT or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact your local Albaugh representative at 1-800-247-8013.

USE PRECAUTIONS:

- Applications to wheat, barley or oats that are stressed by severe weather conditions, frost, low fertility, drought, water logged soil (soils at or near field capacity), disease or insect damage may result in crop injury. Drought, disease or insect damage following application may also result in crop injury.
- Do not apply to irrigated land where tail water will be used to irrigate cropland.
- Do not contaminate irrigation water.
- Do not apply to crops underseeded to legumes or grasses as injury to forage may result.
- Do not remove soil from treated fields for use on lawns, in gardens or for back fill.
- DRAFT should be applied only with water as a carrier. Other carriers may accelerate breakdown of the product and reduce its effectiveness.
- Do not exceed a total of 20 g of DRAFT per hectare per crop year.
- Do not plant to any crop until two months after application.
- Wheat, barley or oats may not be grazed by or fed to livestock within 7 days of application.
- Under certain conditions (such as heavy rainfall, prolonged cool weather, frost conditions, wide fluctuations in day/night temperatures) temporary lightening in crop colour and occasionally a slight reduction in crop height may occur.
- Do not use on highly variable soils that have large gravelly or sandy areas, eroded knolls, or calcium deposits.
- Limit use of DRAFT to currently licensed varieties of wheat (spring, durum, winter), barley, oats and SU tolerant canola. Consult Albaugh prior to use on unlicensed varieties for crop tolerance information. **DO NOT USE ON LEGER BARLEY AND BELVEDERE WHEAT.**

2022-6356

2022-12-05

Last reviewed Sub. No. 2022-1981, 2022-06-01

With changes from Sub. No. 2022-3214, 2022-08-04

Overspray or drift to important wildlife habitats such as shelterbelts, wetlands, sloughs or dry slough borders and woodlots should be avoided.

STORAGE:

Store this product away from food or feed. Not for use or storage in or around the home. Keep container closed.

DISPOSAL:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

If this is a refillable container:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

If this is a returnable container:

Do not reuse this container for any other purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For all package types:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

*All other products mentioned are trademarks of their respective companies