

JM-IMAXI 70WG-LEAFLET-20181109

SIZE:180(W)X150(H)MM PJMI02M010181109

EMPTY CONTAINERS NEED TO BE WASHED THREE TIMES; WATER RESULTING FROM THE TRIPLE RINSE MUST BE ADDED TO THE SPRAY TANK.

USING THE EMPTY PACKING AND CONTAINERS FOR DIFFERENT USE AS INTENDED, REPRESENTS A RISK TO HUMANS AND THE ENVIRONMENT.

GUARANTEE NOTE

The manufacturer and registration owner guarantees the quality of the product in the original and sealed hermetic container. It also guarantees the efficacy of the product, if it is used as indicated in the pamphlet. Since the storage and handling is out of its control it is impossible to assume the responsibility of damage resulting from such factors. We also guarantee the chemical composition of the product as indicated in the label, when the bottle is originally sealed.

FORMULATOR:

JIANGSU ROTAM CHEMISTRY CO. LTD.,
N° 88 Rotam Road, ETDZ, Kunshan 215301, Jiangsu, CHINA.
TEL: 0512 – 5771 – 1988 FAX: 0512 – 5771 – 8697

COUNTRY
JAMAICA

REGISTRATION NUMBER
MHP1/019/18

DATE OF REGISTRATION
11-SEPTEMBER-2018

¡STOP! READ THE PAMPHLET BEFORE USING THE PRODUCT AND CONSULT A PROFESSIONAL IN AGRICULTURAL SCIENCES

IMAXI® 70 WG

NEONICOTINOID - INSECTICIDE
IMIDACLOPRID

CAUTION

ANTIDOTE: NO KNOWN ANTIDOTE

THIS PRODUCT IS MODERATELY TOXIC, IT DOES NOT IRRITATE EYES OR SKIN, HOWEVER NORMAL PRECAUTIONS FOR THE HANDLING OF AGROCHEMICALS SHOULD BE OBSERVED DURING HANDLING AND APPLICATION OF THIS PRODUCT

**DO NOT STORE IN HOMES
STORE AWAY FROM CHILDREN, MENTALLY ILL PEOPLE,
DOMESTIC ANIMALS, FOOD AND MEDICINE**



AGRONOMIC USES:

MODE OF ACTION:

IMAXI® 70 WG is an insecticide that acts systemically, by ingestion and by contact. Once absorbed in the plant it is very systemic, moving acropetally and reaching all parts of the plant.

APPLICATION EQUIPMENT:

IMAXI® 70 WG can be applied with manual or motorized equipments, as back sprayers or by tractor, using high volume, low volume or ULV nozzles. Make sure that the spraying equipment is in good condition. Calibrate with water before spraying. Do not apply trough irrigation systems. Spraying equipment used for the application of **IMAXI® 70 WG** requires cleaning according with the manufacturers of the equipment. This product does not have any specific recommendation in this respect. To realize the applications, it is recommended to use cone spraying tips, to give a good coverage of the foliage. Use the following personal protection equipment while handling, mixing and using the product: rubber boots, respirator, goggles, coverall and gloves.



PREPARING THE TANK MIX:

It is recommended to prepare a premix, mixing water with the required amount of product in a bucket. Then fill the spray tank to ½ the volume of the required clean water and start stirring. Add the premix of the product and complete the total volume with water. Add none ionic surfactant at the end in a proportion of 0,25% to the total volume of water. Use complete personal protection equipment to prepare the spray mix. The used equipments need to be washed carefully after preparing and applying the product.



JM-IMAXI 70WG-LEAFLET-20181109

SIZE: 180(W)X150(H)MM

AUTHORIZED USES: RECOMMENDATIONS FOR USE:

CROPS	PESTS	RATES	OBSERVATIONS
Rice (<i>Oryza sativa</i>)	Stinkbug (<i>Nezara viridula</i>) (<i>Oebalus</i> spp.); (<i>Tibraca</i> spp.)	0.140 Kg/Ha (0.100 Kg/Mz)	Apply in flowering stage, (if required).
	Hidrellia (<i>Hidrellia</i> spp.), Rice leafhopper (<i>Sogatodes oryzicola</i>)	0.035 - 0.070 Kg/Ha (0.025 - 0.050 Kg/Mz)	First application (ground or by air) Apply only in tank mix with herbicides, 8 to 15 days after seeding or transplant. Second application 30 to 45 days after seeding (if required)
Potato (<i>Solanum tuberosum</i>)	Trips (<i>Trips</i> spp.)	0.5 kg/ha 0.35 kg/Mz	Apply the complete rate of IMAXI on the potato seed in the open furrow.
Cucumber (<i>Cucumis sativus</i>)	Afidis (<i>Aphis</i> spp.), (<i>Myzus persicae</i>); White Fly (<i>Bemisia tabaci</i>)(Gennadi us)		Apply only with directed spray to the base of the plant 1 to 7 days after transplant. Also, can be applied by drip irrigation. Transplants should be treated with IMAXI in the seeder.
Sandía (<i>Citrullus lanatus</i>)			
Tobacco (<i>Nicotiana tabacum</i>)			
Tomato (<i>Solanum lycopersicum</i>)			
Peppers (<i>Capsicum</i> spp.)			
Melón (<i>Cucumis melo</i>)			
Papaya (<i>Carica papaya</i>)		White Fly (<i>Bemisia tabaci</i>)(Gennadi us)	2 cups/ 16 liter back sprayer.

APPLICATION TIMING:

Apply with appearance of the first symptoms of insects or preventively in plots where insect attack has been reported previously.

APPLICATION INTERVAL:

Apply in the seeder and one time at or 1 to 7 days after transplanting. Apply to the base of the plant or seedling.

PRE HARVEST INTERVAL: 21 days

REENTRY INTERVAL:

Reentry to treated areas is recommended when the spray residue has dried completely, if re entering is done before that it is recommended to use full personal protection equipment, same as during the spray.

PHYTOTOXICITY:

IMAXI® 70 WG is not phytotoxic to the crops when used at the recommended rates.

COMPATIBILITY:

When used as recommended by the manufacturer, IMAXI® 70 WG is compatible and can be applied in mixture with other Agrochemical products of common use in agriculture, however before preparing a tank mix for the first time it is recommended to do small scale test.

PRECAUTIONS AND INSTRUCTIONS OF USE

STORAGE AND TRANSPORT

Do not transport or store this product together with food, medicine, clothing, household utensils or forages. Store this product under lock, in a fresh, dry place, away from heat or fire. Keep and store this product only in its original container, with corresponding label and well sealed. Transport and store away from herbicides. IMAXI® 70 WG is not flammable or corrosive, but do not place near to sources of high heat.

**DO NOT STORE IN HOMES
STORE AWAY FROM CHILDREN, MENTALLY ILL PERSONS
DOMESTIC ANIMALS, FOOD AND MEDICINE**

**DO NOT EAT, SMOKE OR DRINK DURING THE HANDLING AND APPLICATION OF
THIS PRODUCT, TAKE A SHOWER AFTER WORK AND PUT ON CLEAN
CLOTHES.**

SYMPTOMS OF INTOXICATION:

There are no specific symptoms of intoxication. The product does not irritate skin or eyes.

FIRST AID

INGESTION: Induce vomiting by giving a mixture of warm water with salt.

INHALATION: Move the affected person to a place with fresh air and maintain at rest until recovered.

CONTACT WITH EYES: Clean the eye with abundant water for 15 minutes.

CONTACT WITH SKIN: If there is contact with skin, clean with abundant water and soap, remove all contaminated clothing and clean skin with water and soap.

**NEVER GIVE DRINK OR INDUCE VOMITING TO A PERSON WHICH IS NOT
CONSCIOUS**

ANTIDOTE AND MEDICAL TREATMENT:

There is no specific antidote, treat the patient symptomatically.

NATIONAL CENTERS OF INTOXICATIONS

COUNTRY	INSTITUTION	TELEPHONE
JAMAICA	POISON CONTROL CENTER CARPIN	1-888-764-7667 Or 188-POISONS

MEASURES TO PROTECT THE ENVIRONMENT

TOXIC TO FISH AND CRUSTACEA



TOXIC TO BIRDS AND BEES



**DO NOT CONTAMINATE RIVERS, LAKES, PONDS OR GROUND WATER WITH THIS
PRODUCT OR WITH CONTAINERS AND BOTTLES**

MANAGEMENT OF BOTTLES, PACKING, RESIDUES AND REMANENTS

In case of liquid spillage or residues of pesticides absorb with sand, wood chips or any absorbent material, for solids remove as much as possible, place the residues in a hermetic container and take to the distributor or eliminate them in a land fill authorized by the Ministry of Health.

Use the entire content of the bottles and when empty wash and triple rinse with clean water, adding the solution to the spray tank, destroy or puncture the empty containers. If the country has an official program for collection and empty container disposal, take these to the closest collecting center or dispose according to the instructions of the distributor of the product.