



# Euskatel<sup>®</sup> PLUS

EUSKATEL PLUS is an Emulsifiable Concentrate (EC) formulation containing 125 g/L Prothioconazole and 125 g/L Tebuconazole for use as a fungicide in winter and spring wheat, winter rye, winter and spring barley, winter and spring oats and oilseed rape.

F

MAPP 20147

## EUSKATEL PLUS

an Emulsifiable Concentrate (EC) formulation containing 125 g/L Prothioconazole and 125 g/L Tebuconazole

**Causes skin irritation.**

**May cause an allergic skin reaction.**

**Causes serious eye irritation.**

**May cause respiratory irritation.**

**Suspected of damaging the unborn child.**

**Very toxic to aquatic life with long lasting effects.**

Do not get in eyes, on skin or on clothing.

Avoid release to the environment.

Use personal protective equipment as required.

IF SWALLOWED: Call a POISON CENTER or doctor/physician if you feel unwell.

Collect spillage.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply with the instructions for use.**

WARNING



UFI: XTTW-EP89-U43V-VDTV

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work

### SAFETY PRECAUTIONS

#### Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION when handling the concentrate.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) when applying the product.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH ANY CONTAMINATION from eyes immediately.

WASH HANDS AND EXPOSED SKIN before meals and after work.

IF YOU FEEL UNWELL, seek medical advice (show the label where possible).

#### Environmental Protection

DO NOT CONTAMINATE SURFACE WATERS OR DITCHES with chemical or used container.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application.

Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

#### Storage and Disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

### Emergency Information:

CARECHEM: 01235 239 670 (24 hour)

Technical Enquiries: 0800 078 9649

For Batch Number and Manufacturing Date: see container

### Approval holder and Marketing company:

Albaugh UK Ltd., 1 Northumberland Avenue,  
Trafalgar Square, London WC2N 5BW, United Kingdom  
Tel: 020 3551 2580 - www.albaugh.eu



® TM trademark or trade name of Albaugh, LLC or an affiliated company



# Euskatel<sup>®</sup> PLUS

EUSKATELPLUS is an Emulsifiable Concentrate (EC) formulation containing 125 g/L Prothioconazole and 125 g/L Tebuconazole for use as a fungicide in winter and spring wheat, winter rye, winter and spring barley, winter and spring oats and oilseed rape.



MAPP 20147

**GROUP 3 FUNGICIDE**

## IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL FUNGICIDE

Crops/Situations	Maximum Individual Dose (l product/ha)	Maximum Number of Treatments	Latest Time of Application
Wheat, rye (winter)	1.0	Two per crop	End of flowering (BBCH 69)
Barley, oats	1.0	Two per crop	Beginning of flowering (BBCH 61)
Oilseed rape	1.0	Two per crop	56 days before harvest

### Other specific restrictions:

A minimum interval of 21 days must be observed between applications

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**



### Approval holder and Marketing company:

Albaugh UK Ltd., 1 Northumberland Avenue, Trafalgar Square, London WC2N 5BW, United Kingdom  
Tel: 020 3551 2580 - [www.albaugh.eu](http://www.albaugh.eu)

### Emergency Information:

CARECHEM: 01235 239 670 (24 hour)

Technical Enquiries: 0800 078 9649

For Batch Number and Manufacturing Date: see container

® TM trademark or trade name of Albaugh, LLC or an affiliated company

## DIRECTIONS FOR USE

**IMPORTANT:** This information is authorised as part of the product label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

## RESTRICTIONS/WARNINGS

EUSKATEL PLUS provides 'moderate control' of light leaf spot.

A minimum interval of 21 days must be observed between applications in all crops.

PROTECT FROM FROST

## CROP SPECIFIC INFORMATION

EUSKATEL PLUS may be used on all commercial varieties of winter and spring barley, winter and spring wheat, winter rye, winter and spring oats and oilseed rape.

### Cereals

#### Eyespot (*Tapesia* spp.)

Spray in the spring at the first sign of disease, from when the leaf sheaths begin to become erect until the 2<sup>nd</sup> node is detectable (GS 30-32).

#### Septoria Leaf Blotch and Glume Blotch (*Septoria tritici* and *Stagonospora nodorum*)

Apply before disease is established in the crop. To protect the upper leaves and ear apply EUSKATEL PLUS at full flag leaf emergence (GS 37) up to mid-flowering (GS 65). Where disease pressure remains high application may be repeated.

Applications to upper leaves where *S. tritici* symptoms are present are likely to be less effective. EUSKATEL PLUS contains two DMI fungicides. Resistance to some DMI fungicides has been identified in Septoria leaf blotch (*Mycosphaerella graminicola*) which may seriously affect the performance of some products. For further advice on resistance management in DMIs contact your agronomist or specialist advisor and visit the FRAG-UK website.

#### Powdery Mildew (*Blumeria graminis*)

Apply EUSKATEL PLUS at the first signs of disease. Where disease pressure remains high application may be repeated. EUSKATEL PLUS will only provide moderate control of powdery mildew in wheat.

#### Yellow Rust (*Puccinia striiformis*)

Apply EUSKATEL PLUS at the first signs of disease. A second application may be made later if re-infection occurs. Applications made to established infections are likely to be less effective.

#### Brown Rust (*Puccinia recondita* and *P. hordei*)

Apply EUSKATEL PLUS at the first signs of disease. A second application may be made later if re-infection occurs. Applications made to established infections are likely to be less effective.

#### Crown Rust (*Puccinia coronata*)

Apply EUSKATEL PLUS at the first signs of disease. EUSKATEL PLUS controls crown rust in winter and spring oats.

A second application may be made later if re-infection occurs. Applications made to established infections are likely to be less effective.

#### Tan Spot (*Pyrenophora tritici-repentis*)

Apply EUSKATEL PLUS at the first signs of disease in spring or early summer. Where disease pressure remains high application may be repeated.

### Ear Disease Complex

Apply EUSKATEL PLUS soon after ear emergence until the end of flowering (GS 59-69) for control of Fusarium ear blight in wheat (or moderate control in barley) and reduction of sooty moulds.

Control of ear diseases can result in cleaner, brighter ears.

Through the control of ear blight, EUSKATEL PLUS effectively reduces the level of the Fusarium mycotoxin deoxynivalenol (DON) in wheat grain. However, where Fusarium levels are high, the reduction achieved may not always be sufficient to ensure that DON levels fall below the statutory limit.

### Leaf Blotch (*Rhynchosporium secalis*)

Apply EUSKATEL PLUS in spring at the first signs of disease. For severe infections a second application may be necessary later.

### Net Blotch (*Pyrenophora teres*)

Apply EUSKATEL PLUS at the first signs of disease in spring/early summer. For severe infections, a second application later will give most effective control when conditions remain favourable for disease development.

### **Oilseed rape**

#### Light Leaf Spot

Apply EUSKATEL PLUS in autumn/winter (usually late October to early December) protectively. Follow up spray(s) may be required in early spring from the onset of stem elongation, depending on disease development.

Please note: EUSKATEL PLUS provides 'moderate control' of light leaf spot.

#### Phoma Leaf spot/Stem Canker

Apply EUSKATEL PLUS in autumn at the first sign of disease. Repeat application in late autumn/winter, if disease symptoms reoccur.

#### Sclerotinia stem rot (*Sclerotinia sclerotiorum*)

Apply EUSKATEL PLUS at early to full flower

### **MIXING AND SPRAYING**

Thoroughly shake the pack before use.

Add the required quantity of EUSKATEL PLUS to the half-filled spray tank with the agitation system in operation and then fill to the required level. Continue agitation at all times during spraying and stoppages until the tank is completely empty. Spray immediately after mixing.

Sprayers should be thoroughly cleaned with water and detergent after use, and filters and jets checked for damage and blockages. Boom height should be adjusted to ensure even coverage of the crop, particularly at later growth stages. The correct height is one at which the spray from alternate nozzles meets just above the crop, in dense crops, at later growth stages, higher water volumes should be used.

Apply EUSKATEL PLUS in 100-400 litres water per hectare. The higher spray volumes are recommended where the crop is dense or disease pressure / risk is high to ensure good penetration to the lower leaves and stem bases. Disease control may be compromised by reducing water volumes, where good spray coverage is difficult to achieve.

A spray pressure of 2-3 bar is recommended. Apply as a MEDIUM spray quality (as defined by BCPC).

## RESISTANCE MANAGEMENT

Repeated application of EUSKATEL PLUS alone should not be used on the same crop against a high risk pathogen such as cereal powdery mildew. Tank mixtures or alternation with fungicides having a different mode of action (e.g. morpholines) have been shown to protect against the development of resistant forms of disease.

**CAUTION:** The possible development of disease strains resistant to EUSKATEL PLUS cannot be excluded or predicted. Where such resistant strains occur, EUSKATEL PLUS is unlikely to give satisfactory control.

**Conditions of Supply:** all goods supplied by us are of high quality and we believe them to be correct but, as we cannot exercise control over their storage, handling, mixing or use, or weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or resellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.