

# MALIBU™

**CONTROL UNWANTED NOXIOUS WEEDS, WOODY PLANTS, AND BRUSH ACROSS A RANGE OF USE SITES.**



Malibu is used for control of annual and perennial broadleaf weeds including invasive and noxious weeds, certain annual grasses, and certain woody plants and vines on:

- Rangeland, permanent grass pastures (including grasses grown for hay\*), Conservation Reserve Program (CRP),
- Non-crop areas for example, airports, barrow ditches, communication transmission lines, electric power and utility rights-of-way, fencerows, gravel pits, industrial sites, military sites, mining and drilling areas, oil and gas pads, non-irrigation ditch banks, parking lots, petroleum tank farms, pipelines, roadsides, railroads, storage areas, dry storm water retention areas, substations, unimproved rough turf grasses,
- Natural areas (open space) for example, campgrounds, parks, prairie management, trailheads and trails, recreation areas, wildlife openings, and wildlife habitat and management areas including seasonally dry flood plains, deltas, marshes, prairie potholes, or vernal pools,
- Grazed areas in and around these sites.

Malibu is also used for control of annual and perennial broadleaf weeds in wheat (including spring wheat, winter wheat, and durum) and field corn.

\*Hay can only be used on the farm or ranch where product is applied.

## PRODUCT FEATURES

- Provides pre- and postemergence control of emerging seedlings of susceptible weeds and re-growth of certain perennial weeds following application.
- Important component of integrated vegetation management programs designed to renovate or restore desired plants.
- Provides long-term control of susceptible weeds, when applied according to labeled directions.
- Compatible in tank mix with other labeled brush-control herbicides.
- Excellent environmental profile that allows application up to the water's edge.
- Low use rate of 7 oz/A (maximum broadcast rate) and 14 oz/A (maximum spot treatment rate).

See label for complete application guidelines and restrictions.

### ACTIVE INGREDIENT

2.0 lb/gal aminopyralid

### HRAC GROUP

4

### FORMULATION

Soluble concentrate

### REI

48 hours

### COMPARE TO

Milestone®

### MODE OF ACTION

Synthetic auxin

### SIGNAL WORD

N/A

### RESTRICTED USE

No

### PACKAGING

2 x 2.5 gal, bulk

### EPA REG NUMBER

83100-66 | AD080223



# MALIBU™

## FREQUENTLY ASKED QUESTIONS

### How should I apply Malibu on pasture/rangeland for best results?

- Usage rate: 3.0-14.0 fl oz/A, depending on the weed species. Refer to the label for the appropriate rate range and any usage restrictions for the weed(s) needing control.
- Malibu may be applied as a broadcast spray by ground or aerial equipment or as a spot application to control weeds including, but not limited to, those listed on the product label.
- When a rate range is given, use the higher rate to control weeds at advanced growth stages or when under less-than-favorable growing conditions.
- For optimum uptake and translocation, avoid mowing, haying, shredding, burning, or soil disturbance in treated areas for at least 14 days following application.
- To maximize and extend the benefits of weed control provided by Malibu, it is important that other vegetation management practices, including proper grazing management, biological control agents, replanting, fertilization, prescribed fire, etc., be used in appropriate sequences and combinations to further alleviate the adverse effects of weeds on desirable plant species and to promote development of desired plant communities.
- Consult with appropriate state or local water authorities before applying this product around public waters. State or local public agencies may require permits.

### What restrictions do I need observe when using Malibu on pasture and rangeland?

- DO NOT use grasses treated with Malibu in the preceding 18 months for hay intended for export outside the United States.
- Hay from areas treated with Malibu in the preceding 18 months CANNOT be distributed or made available for sale off the farm or ranch where harvested unless allowed by supplemental labeling.
- Hay from areas treated with Malibu in the preceding 18 months CANNOT be used for silage, haylage, baleage, and green chop unless allowed by supplemental labeling.
- DO NOT move hay made from grass treated with Malibu within the preceding 18 months off farm unless allowed by supplemental labeling.
- DO NOT use hay or straw from areas treated with Malibu within the preceding 18 months or manure from animals feeding on hay treated with Malibu in compost.
- DO NOT use grasses treated with Malibu in the preceding 18 months for seed production.
- Refer to product label for additional restrictions for haying, grazing, and manure uses, and crop rotation restrictions.



◀ See label for complete application guidelines and restrictions.

©2025 Albaugh, LLC. All rights reserved. Albaugh and the Albaugh logo are registered trademarks and Malibu is a trademark of Albaugh, LLC. Milestone is a registered trademark of Corteva Agriscience. Always read and follow label directions, precautions and restrictions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status. All products are EPA registered.