

## INSECTICIDE

# DIAMID™ T&O

**DELIVERS SEASON-LONG CATERPILLAR AND GRUB CONTROL.**



Diamid™ T&O insecticide utilizes the active ingredient chlorantraniliprole to deliver season-long grub control with tremendous application flexibility and convenience. The highly concentrated formulation in Diamid T&O performs best when applied early. A single (4 - 8 fl oz/A) application, in early spring, will deliver control of all grub species and key surface feeding pests including several species of caterpillars as well as annual bluegrass weevil.

## PRODUCT FEATURES

- Season-long control of all white grub species and caterpillars including black cutworms, sod webworms, and fall armyworms.
- Excellent billbug and annual bluegrass weevil control.
- Flexible timing of rainfall or irrigation post application.
- No known adverse effects on beneficial and non-target organisms, including earthworms and honeybees.

## LABELED USES

- Commercial ornamental nurseries and greenhouses
- Commercial sod farms
- Golf courses
- Sports and recreational turf
- Lawns and landscaped areas on and around residential, commercial, and institutional properties
- Interior plantscapes

See label for complete application guidelines and restrictions.

### ACTIVE INGREDIENT

Chlorantraniliprole

### IRAC GROUP

28

### FORMULATION

Suspension concentrate

### CHEMICAL FAMILY

Anthranilic diamide

### COMPARE TO

Acelepryn®, Durentis™

### MODE OF ACTION

Ryanodine receptor modulator

### RESTRICTED USE

No

### SIGNAL WORD

Caution

### PACKAGING

4 x 1 qt

### EPA REG NUMBER

89442-70

# DIAMID™ T&O

DELIVERS SEASON-LONG CATERPILLAR AND GRUB CONTROL.

Diamid T&O insecticide is a member of the anthranilic diamide class of insecticides with a mode of action acting on insect ryanodine receptors. Diamid T&O is effective through ingestion of treated plant material. After exposure affected insects will rapidly stop feeding, become paralyzed, and typically die within 1 - 3 days. Time applications to the most susceptible insect pest stage, typically at egg lay, egg hatch and/or newly hatched larvae, before populations reach damaging levels. When pest populations are high, use the highest listed application rate for that pest.

## APPLICATION TIMING

PEST	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
White grubs				█	█	█	█	█				
Billbugs				█	█							
Adult Japanese beetles					█	█	█	█				
European crane fly									█	█	█	
Turf caterpillars	AS NEEDED											



◀ See label for complete application guidelines and restrictions.

©2024 Albaugh, LLC. All rights reserved. Albaugh and the A logo are registered trademarks and Diamid is a trademark of Albaugh, LLC. Acelepryn is a registered trademark of Syngenta. Durentis is a trademark of FMC. Always read and follow label restrictions, precautions, and directions for use. This product may not be registered for sale or use in all states or counties. Please check with your state agency responsible for pesticide registration to ensure registration status.