

FUNGICIDE

FAIRVIEW SELECT®

FOR DISEASE CONTROL OF TURFGRASSES
ON GOLF COURSES ONLY.



Fairview Select® is a broad-spectrum fungicide used to control difficult diseases of turfgrass on golf courses including anthracnose, dollar spot, brown patch, red thread, powdery mildew, and more. The combination of fungicides from 2 groups in **Fairview Select** supports disease-resistance management programs. **Fairview Select** works both systemically and on contact to control the toughest turf diseases on golf courses.

PRODUCT FEATURES

- Provides broad-spectrum control of tough to fight diseases in turfgrass.
- For golf course turf use only - including greens, tees, fairways, and aprons.
- Supports disease-resistance management programs with the combination of FRAC 29 and FRAC 3 active ingredients.
- Manages and controls diseases both systemically and on contact.
- For use on all golf course turf applications of warm- and cool-season grasses.

LABELED USES

- Golf course tees, greens, fairways, and rough.

See label for complete application guidelines and restrictions.

ACTIVE INGREDIENT

17.0% fluazinam
17.6% tebuconazole

FRAC GROUP

Fluazinam - Group 29
Tebuconazole - Group 3

FORMULATION

Soluble liquid concentrate

CHEMICAL FAMILY

FRAC 29 fungicides - Fluazinam
(2,6-dinitroanilines)

FRAC 3 fungicides - Tebuconazole
(Triazoles)

COMPARE TO

Traction®

MODE OF ACTION

Cell membrane disruptors, oxidative phosphorylation uncouplers

RESTRICTED USE

No

SIGNAL WORD

Caution

PACKAGING

2 x 2.5 gal

EPA REG NUMBER

89442-46

FAIRVIEW SELECT®

FOR DISEASE CONTROL OF TURFGRASSES ON GOLF COURSES ONLY.

APPLICATION INFORMATION

Resistance management

For resistance management, please note that **Fairview Select**® contains both fluazinam and tebuconazole. Fluazinam is classified in the 2,6-dinitro-anilines chemical class as a FRAC 29, respiration inhibitors. Tebuconazole is classified in the triazole chemical class as a FRAC 3, sterol synthesis. Any fungal population may contain individuals naturally resistant to **Fairview Select** and other FRAC 29 or FRAC 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same turf areas. Appropriate resistance management strategies should be followed.



Labeled Diseases	Use Rate	Retreatment Interval (days)
Dollar spot, Copper spot, Powdery mildew	1.3 fl oz /1,000 sq ft	14 days
Leaf spot (Melting out), Brown blight	1.3 fl oz /1,000 sq ft	14 days
Brown patch, Zoysia patch, Large patch	1.3 fl oz /1,000 sq ft	14 days
Red thread	1.3 fl oz /1,000 sq ft	14 days
Anthracnose	1.3 fl oz /1,000 sq ft	14 days
Gray snow mold, Pink snow mold	1.3 fl oz /1,000 sq ft	14 days
Pink patch	1.3 fl oz /1,000 sq ft	14 days
Stem rust	1.3 fl oz /1,000 sq ft	14 days
Algal scum* (filamentous blue-green algae/cyanobacteria)	1.3 fl oz /1,000 sq ft	14 days
Brown ring patch	1.3 fl oz /1,000 sq ft	21 days
<i>Fusarium</i> patch	1.3 fl oz /1,000 sq ft	21 days
Gray leaf spot	1.3 fl oz /1,000 sq ft	21 days
Stripe smut	1.3 fl oz /1,000 sq ft	One application only

* *Suppression*



◀ See label for complete application guidelines and restrictions.

©2023 Albaugh, LLC. Albaugh, the A logo, and Fairview Select are registered trademarks of Albaugh, LLC. Traction is a registered trademark of Nufarm Americas Inc. Always read and follow label restrictions, precautions, and directions for use. This product may not be registered for sale or use in all states or counties. Please check with your state agency responsible for pesticide registration to ensure registration status.