



MACHO[®] 4.0 FL

BROAD-SPECTRUM, SYSTEMIC INSECTICIDE THAT PROVIDES EFFECTIVE, ECONOMICAL CONTROL OF A RANGE OF INSECTS.

Macho[®] 4.0 FL is a systemic soil- and foliar-applied insecticide that provides economical control of a wide range of insects in a variety of crops. When applied to the soil, moves from the plant's root system to the upper vegetative parts via the xylem tissue. This results in extended insecticidal activity to control pests that can vector detrimental virus transmission. Macho[®] 4.0 FL can be applied to a variety of crops to provide selective control of many key piercing and sucking insects.

PRODUCT FEATURES

- Offers outstanding, long-lasting control of aphids, Colorado potato beetles, leafhoppers, mealybugs, whiteflies and many other piercing and sucking insects.
- May be applied by ground, aerial or chemigation systems for the reduction and prevention of virus transmission by insects.
- Systemic in nature which allows for continuous movement throughout all vegetative parts of the plant for an extended period.

See label for complete application guidelines and restrictions.



ACTIVE INGREDIENT

4.0 lbs/gal imidacloprid

IRAC GROUP

4A

FORMULATION

Liquid

REI

12 hours (refer to label for exceptions for soil injection and soil incorporation)

COMPARE TO

Admire[®] Pro

MODE OF ACTION

Nicotinic acetylcholine receptor competitive modulator

SIGNAL WORD

Caution

RESTRICTED USE

No

PACKAGING

4 x 1 gal

EPA REG NUMBER

42750-140 | AD07072016

MACHO[®] 4.0 FL

FREQUENTLY ASKED QUESTIONS

How should Macho 4.0 FL be used in soil applications for best results?

- Best results are achieved when applications are made to the root-zone of plants.
- The rate of Macho 4.0 applied affects the length of the plant protection. Use the specified higher rates when infestations occur later in crop development or where pest pressure is continuous.
- Pre-mix Macho 4.0 FL with water or other appropriate diluents prior to application. Maintain constant agitation to avoid settling.
- Do not apply more than 0.5 lb. active ingredient per acre per year regardless of formulation or method of application, unless specified within the crop-specific, application section for a given crop.

How should Macho 4.0 FL be used in foliar applications for best results?

- Thorough coverage of foliage without runoff is required for maximum insecticidal efficacy. Use of a spray adjuvant may enhance thorough coverage.
- Apply at a minimum of 10 gpa by ground or 5 gpa by air. Macho 4.0 may be applied by overhead chemigation. Refer to label pages 8-9 for chemigation instructions, if allowed in crop-specific application sections.
- Do not apply Macho 4.0 in enclosed structures such as planthouses or greenhouses except as specified in the TOBACCO, CUCURBIT VEGETABLES, FRUITING VEGETABLES and GREENHOUSE VEGETABLES, (Mature plants in production greenhouses): Cucumber, Tomato only sections of the product label.
- Do not apply more than 0.5 lb. active ingredient per acre per year regardless of formulation or method of application, unless specified within the crop-specific, application section for a given crop.

What rotational restrictions should be observed when using Macho 4.0 FL?

- Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application.
- All crops on the Macho 4.0 FL label, plus barley, canola, corn (field, pop, and sweet), rapeseed, sorghum, sugarbeet, and wheat have no plantback restrictions.
- For cereal crops including buckwheat, millet, oats, rice, rye, and triticale, as well as safflower, observe a 30-day plantback interval.
- For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established there is a 12-month plantback interval.

APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.



◀ See label for complete application guidelines and restrictions.

©2023 Albaugh, LLC. All rights reserved. Albaugh, the Albaugh logo, and Macho are registered trademarks of Albaugh, LLC. Admire is a registered trademark of Bayer CropScience LLP. Always read and follow label restrictions, precautions and directions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.