

OXYSTAR[®] 4L

RESIDUAL PRE- OR POSTEMERGENCE HERBICIDE FOR CONTROL OF GRASS AND BROADLEAF WEEDS.



OxyStar[®] 4L herbicide is a selective herbicide for postemergence and preemergence residual weed control in labeled crops, including artichokes (globe), blackberry/raspberry, broccoli/cabbage/cauliflower, cacao, citrus (nonbearing), coffee, conifer (seedbeds, transplants, container stock) and selected deciduous trees, corn, cotton, cottonwood, eucalyptus, fallow bed, cotton/soybeans, fallow land, garbanzo beans, garlic, grapes, guava (Hawaii only), horseradish, jojoba, mint, onions, onions grown for seed, papaya (Hawaii only), soybeans, taro, tree fruit/nut/ vine crops, and other listed crops.

PRODUCT FEATURES

- Provides long-lasting residual control by preventing the growth of weeds and eliminating weeds after sprouting.
- California VOC approved, low-odor formulation.
- May be applied by ground, air or through chemigation systems.
- Excellent tank-mix partner with other EPA-approved herbicides to expand weed control in labeled crops.

ACTIVE INGREDIENT

4.0 lbs/gal oxyfluorfen

HRAC GROUP

14

FORMULATION

Liquid

REI

48 hours to 6 days
(refer to label for exceptions)

COMPARE TO

GoalTender[®]

MODE OF ACTION

PPO inhibitor

SIGNAL WORD

Warning

RESTRICTED USE

No

PACKAGING

4 x 1 gal

EPA REG NUMBER

42750-199 | AD072821

See label for complete application guidelines and restrictions.

OXYSTAR® 4L



FREQUENTLY ASKED QUESTIONS

How should OxyStar 4L be applied to tree fruit, tree nut, and vine crops for best results?

Preemergence (dormant application)

- Apply 2.5-3.0 pt/A broadcast or 2.5-4.0 pt/A banded on tree fruit, tree nut, and vine crops listed on page 47 of the product label.
- Apply a minimum of 20 gallons of water per acre. Use higher spray volumes to ensure thorough coverage in high densities of emerged weeds or heavy trash. Sprays must be directed to the soil and the base of dormant trees or vines.
- In California, OxyStar 4L may be applied as an over-the-top or directed spray to dormant nonbearing grape plantings. The use of a low-pressure sprayer is suggested. DO NOT apply over-the-top to grape plantings that are under stress due to drought, flooding, excessive fertilizer or soil salts, storage conditions, wind injury, hail, injury from previously applied pesticides, or injury due to insects, nematodes, or diseases, as severe crop injury may result.

Postemergence (dormant application)

- Apply 1.0-3.0 pt/A broadcast or 1.0-4.0 pt/A banded on tree fruit, tree nut, and vine crops listed on page 47 of the product label.
- Apply in a spray volume of 40 or more gallons per acre. For optimum control, apply when weeds are at seedling stage of growth.
- The lower rate in the rate range (1 pint (0.5 lbs of active) per acre) is required for the control of susceptible seedling weeds in the early postemergence stage up to the 4-leaf stage. Higher rates (up to 3 pints (1.5 lbs of active) per acre) may be used for weeds up to the 6-leaf stage. Applications to weeds beyond the 6-leaf stage may result in partial control.

Refer to label page 47-48 for additional usage recommendations and restrictions. Refer to label page 49-51 for non-dormant applications to pistachios, walnuts, almonds and select stone fruits.

How should OxyStar 4L be applied to non-dormant grapes for best results?

- Apply at 1.0 pt/A for preemergence applications and 0.5-1.0 pt/A for postemergence applications to non-dormant grapes.
- OxyStar 4L may be applied preemergence or postemergence to weeds either as a directed spray in a minimum spray volume of 20 gallons per acre or through low-volume sprinkler (micro sprinkler) or drip irrigation systems. Repeat applications may be required. Applications may be made from completion of bloom up to 14 days before to harvest.
- When applied as a postemergence directed spray, add 1 quart 80% active nonionic surfactant cleared for application to growing crops per 100 gallons of spray. Direct sprays to the soil and the base of vines.
- *Refer to label page 32 for additional usage recommendations and restrictions.*



◀ See label for complete application guidelines and restrictions.

©2024 Albaugh, LLC. All rights reserved. Albaugh, the Albaugh logo and OxyStar are registered trademarks of Albaugh, LLC. GoalTender is a registered trademark of Nutrichem. Always read and follow label restrictions, precautions and directions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.