

PLOTTER[®] EXTRA

**SELECTIVE POSTEMERGENCE HERBICIDE FOR
USE ON WHEAT, BARLEY, TRITICALE AND FALLOW.**

Plotter[®] Extra herbicide is a dry flowable granule that is used for selective postemergence weed control in wheat (including durum), barley, triticale and fallow. The best control is obtained when applied to young, actively growing weeds. The degree and duration of control may depend on weed spectrum and infestation intensity, weed size at application and/or environmental conditions at and following treatment.

Provides effective control of a variety of weeds, including wild buckwheat, henbit, blue/purple mustard, Canada thistle, Russian thistle, and tansy mustard.

PRODUCT FEATURES

- Provides effective broadleaf control in wheat and barley crops.
- Outstanding tank-mix partner for added broad spectrum control.
- Long residual activity gives 4-6 weeks of control.
- Controls and/or suppresses over 70 broadleaf and brush weeds.
- Dry formulation makes application convenient and easy.

See label for complete application guidelines and restrictions.



ACTIVE INGREDIENTS

19% tribenuron-methyl +
38% thifensulfuron-methyl +
15% metsulfuron-methyl

HRAC GROUP

2

FORMULATION

Dry flowable granule

REI

12 hours

COMPARE TO

Ally[®] Extra

MODE OF ACTION

ALS inhibitor

SIGNAL WORD

Warning

RESTRICTED USE

No

PACKAGING

8 x 20 oz bottles (160 oz cases)

EPA REG NUMBER

83100-23-83979

PLOTTER[®] EXTRA

FREQUENTLY ASKED QUESTIONS

At what rate should I use Plotter Extra on my wheat/barley/triticale?

- Apply Plotter Extra at 0.2-0.4 oz/A. Use the high rate for heavy infestations of weeds listed on page 3 of the product label, or when application timing and environmental conditions are marginal. Use 0.2-0.3 oz/A for light infestations.

When should I apply Plotter Extra for best results?

- On wheat (except durum and wampum varieties of spring wheat), barley and triticale, make applications after the crop is in the two-leaf stage, but before the flag leaf is visible. Do not harvest within 45 days of the last application.
- On durum and wampum varieties of spring wheat, make applications after the crop is tillering, but before boot. Applications to durum and wampum varieties should be made in combination with 2,4-D.
- Weed control may be reduced if rainfall or snowfall occurs soon after application. Six hours of dry weather are needed to allow the herbicide to be sufficiently absorbed by weed foliage.
- Do not apply to stressed crops.

Can I tank mix Plotter Extra with other herbicides?

- Yes, Plotter Extra may be tank mixed with other suitable registered herbicides to control weeds listed as partially controlled, weeds resistant to Plotter Extra, or weeds not listed under WEEDS CONTROLLED on page 3 of the label. *Refer to TANK MIXTURES on pages 3-4 of the label for specific products and usage instructions.*

Do I need to use an adjuvant with Plotter Extra?

- Yes, include a spray adjuvant such as NIS, COC or MSO with applications of Plotter Extra. In addition, an ammonium nitrogen fertilizer may be used. *Refer to SPRAY ADJUVANTS on page 5 of the label for specific volumes and adjuvant composition.*

What are the rotational restrictions with Plotter Extra?

- Before using Plotter Extra herbicide, carefully consider your crop rotation plans and options. For rotational flexibility, do not treat all of your wheat, barley, triticale or fallow acres at the same time.
- Do not use Plotter Extra on soils having a pH above 7.9 as extended soil residual activity could extend crop rotation intervals beyond normal. Under certain conditions, Plotter could remain in the soil for 34 months or more, injuring wheat, barley or triticale. In addition, other crops planted in high pH soils can be extremely sensitive to low concentrations of Plotter.
- *Refer to label pages 6-9 for specific rotational restrictions by state and soil pH.*



◀ See label for complete application guidelines and restrictions.

©2023 Albaugh, LLC. All rights reserved. Albaugh, the Albaugh logo and Plotter are registered trademarks of Albaugh, LLC. Ally is a registered trademark of FMC Corporation. Always read and follow label restrictions, precautions and directions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.