

GROUP 11 FUNGICIDE

SPECIMEN LABEL

AZOXY 2SC SELECT™

FUNGICIDE

*Broad spectrum
fungicide for control
of plant diseases*

Manufactured For:
Prime Source,
a division of Albaugh, LLC
1525 NE 36th Street
Ankeny, IA 50021

ACTIVE INGREDIENT:

Azoxystrobin: methyl (E)-2-
{2-[6-(2-cyanophenoxy)pyrimidin-
4-yloxy]phenyl}-3-methoxyacrylate*22.9%

OTHER INGREDIENTS:77.1%

TOTAL: **100.0%**

*IUPAC

Contains 2.08 lbs. of active ingredient per gallon
Suspension Concentration

KEEP OUT OF REACH OF CHILDREN

CAUTION

Reformulation is prohibited. See individual
container labels for repackaging limitations.

**See complete First Aid, Precautionary
Statements, Directions For Use, and
Storage and Disposal inside booklet.**



EPA Reg. No. 89442-21

AD052815

FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies involving this product, call CHEMTREC® toll free at 1-800-424-9300.</p>	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemically resistant to this product are listed below. If you want more options follow the instructions for Category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks

USER SAFETY REQUIREMENTS

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash thoroughly with soap and water after handling.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Do not apply directly to water except as specified on this label. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory

Azoxystrobin and a degradate of azoxystrobin are known to leach through soil to ground water under certain conditions as a result of label use. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of azoxystrobin and a degradate of azoxystrobin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Notify State and/or Federal authorities and Prime Source, a division of Albaugh, LLC immediately if you observe any adverse environmental effects due to use of this product.

Physical or Chemical Hazards

Do not mix or allow coming into contact with oxidizing agent. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Use of AZOXY 2SC SELECT™ through air blast application equipment on grapes is prohibited in the following townships and boroughs of Erie County, Pennsylvania: North East, Harborcreek, Lawrence Park, Erie, Presque Isle, Millcreek, Fairview, Girard, and Springfield.

This prohibition is intended to help eliminate phytotoxicity problems with apples observed in this geographic location.

FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. The area being treated must be vacated by unprotected persons.

Do not treat areas while unprotected humans or domestic animals are present in the treatment areas. Because certain states may require more restrictive reentry intervals, consult your State Department of Agriculture for further information.

Do not allow entry into treatment area until area that was treated with this product is dry.

TURF

Golf course turf (not for use in California). Commercial turf farms (not for use in California).

AZOXY 2SC SELECT™ is recommended for control of anthracnose, brown patch, cool weather brown patch (yellow patch), Fusarium patch, gray leaf spot, gray snow mold (Typhula blight), leaf spot, melting out, necrotic ring spot, pink patch, pink snow mold, Pythium blight, Pythium root rot, red thread, Rhizoctonia large patch, southern blight, spring dead spot, summer patch, take-all patch, and Zoysia patch on golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

Integrated Pest (Disease) Management:

Sound turf management resulting in healthy, vigorous turf is the foundation of a good IPM program. Cultural practices such as proper choice of turf variety, nutrient management, proper cutting height, thatch management, and proper watering, drainage, and moisture stress management should be integrated with the use of fungicides to increase turf vigor and reduce the susceptibility to disease. Immunoassay detection kits and extension service diagnostic services can assist in the early and accurate identification of causal organisms and corresponding selection of the proper fungicide when required.

Resistance Management:

Some turf disease pathogens are known to have developed resistance to products used repeatedly for their control. AZOXY 2SC SELECT™ should be applied in a tank mix or alternation program with other registered fungicides that have a different mode of action and to which pathogen resistance has not developed. Do not apply more than two sequential AZOXY 2SC SELECT™ applications for *Pythium* spp. control. For all other diseases when *Pythium* spp. is not present, do not apply more than three sequential applications of AZOXY 2SC SELECT™.

Application Directions:

AZOXY 2SC SELECT™ should be applied prior to disease development. Mix AZOXY 2SC SELECT™ with the required amount of water and apply as a dilute spray application in 2-4 gallons of water per 1,000 square feet (87-174 gallons per acre). Repeat applications at specified intervals for as long as required. For spot treatments, use 0.4 fl. oz. AZOXY 2SC SELECT™ per 1 to 2 gallons of water. Do not apply more than 9.6 quarts product/acre/year (7.1 fl. oz. product/1,000 square feet/year). Apply by ground only.

Rate Ranges:

Use the shortest specified application interval and/or use the higher specified rate when prolonged favorable disease conditions exist.

Dollar Spot:

AZOXY 2SC SELECT™ does not control dollar spot. AZOXY 2SC SELECT™ is compatible in tank mixes with many other fungicides that control dollar spot. Always tank mix AZOXY 2SC SELECT™ with another fungicide that controls dollar spot when this disease is present.

Follow directions under TANK MIXES/COMPATIBILITY above.

DIRECTIONS FOR APPLICATION FOR TURF DISEASES

Target Diseases	Use Rate (fl. oz. product per 1,000 sq. ft.)	Application Interval (days)	Remarks*
Anthracnose (<i>Colletotrichum graminicola</i>)	0.38-0.77	14-28	Apply when conditions are favorable for disease development.
Brown patch (<i>Rhizoctonia solani</i>)	0.38-0.77	14-28	Apply when conditions are favorable for disease development.
Cool weather brown patch Yellow patch (<i>Rhizoctonia cerealis</i>)	0.38-0.77	28	Make one or two applications in fall or when conditions are favorable for disease development.

*Do not apply more than two sequential applications of AZOXY 2SC SELECT™ for control of *Pythium* spp. For all other diseases, do not apply more than four sequential applications of AZOXY 2SC SELECT™.

(continued)

DIRECTIONS FOR APPLICATION FOR TURF DISEASES *(cont.)*

Target Diseases	Use Rate (fl. oz. product per 1,000 sq. ft.)	Application Interval (days)	Remarks*
Fusarium patch <i>(Microdochium nivale)</i>	0.38-0.77	14-28	Apply when conditions are favorable for disease development.
Gray leaf spot <i>(Pyricularia grisea)</i>	0.38-0.77	14-28	Begin applications before disease is present and continue applications while conditions are favorable for disease development.
Gray snow mold Typhula blight <i>(Typhula incarnata, T. ishikariensis)</i>	1.35-0.77	Single application 14	Make a single application of 1.35 fl. oz. or two applications of 0.77 spaced 14 days apart in late fall just before snow cover. Tank mixing with another snow mold fungicide may enhance control under severe disease pressure.

*Do not apply more than two sequential applications of AZOXY 2SC SELECT™ for control of *Pythium* spp. For all other diseases, do not apply more than four sequential applications of AZOXY 2SC SELECT™.

(continued)

DIRECTIONS FOR APPLICATION FOR TURF DISEASES *(cont.)*

Target Diseases	Use Rate (fl. oz. product per 1,000 sq. ft.)	Application Interval (days)	Remarks*
Leaf spot <i>(Bipolaris sorokiniana)</i>	0.38-0.77	14-21	Apply when conditions are favorable for disease development.
Melting out <i>(Drechslera poae)</i>	0.38-0.77	14-21	Apply when conditions are favorable for disease development.
Necrotic ring spot <i>(Leptosphaeria korrae)</i>	0.38-0.77	14-28	Apply when conditions are favorable for disease development.
Pink patch <i>(Limonomyces roseipellis)</i>	0.38-0.77	14-28	Apply when conditions are favorable for disease development.

*Do not apply more than two sequential applications of AZOXY 2SC SELECT™ for control of *Pythium* spp. For all other diseases, do not apply more than four sequential applications of AZOXY 2SC SELECT™.

(continued)

DIRECTIONS FOR APPLICATION FOR TURF DISEASES *(cont.)*

Target Diseases	Use Rate (fl. oz. product per 1,000 sq. ft.)	Application Interval (days)	Remarks*
Pink snow mold <i>(Microdochium nivale)</i>	1.35-0.77	Single application 14	Make a single application of 1.35 fl. oz. or two applications of 0.77 spaced 14 days apart in late fall just before snow cover. Tank mixing with another snow mold fungicide may enhance control under severe disease pressure.
Pythium blight Pythium root rot <i>(Pythium aphanidermatum, Pythium spp.)</i>	0.38-0.77	10-14	Begin applications before disease is present. During periods of prolonged favorable conditions, treat on the 10 day application interval. For use on newly seeded as well as established turf.
Red thread <i>(Laetisaria fuciformis)</i>	0.38-0.77	14-28	Apply when conditions are favorable for disease development.

*Do not apply more than two sequential applications of AZOXY 2SC SELECT™ for control of *Pythium* spp. For all other diseases, do not apply more than four sequential applications of AZOXY 2SC SELECT™.

(continued)

DIRECTIONS FOR APPLICATION FOR TURF DISEASES *(cont.)*

Target Diseases	Use Rate (fl. oz. product per 1,000 sq. ft.)	Application Interval (days)	Remarks*
Rhizoctonia large patch (<i>Rhizoctonia solani</i>)	0.38-0.77	28	Make one or two applications in fall or when conditions are favorable for disease development.
Southern blight (<i>Sclerotium rolfsii</i>)	0.38-0.77	14-28	Apply when conditions are favorable for disease development.
Spring dead spot (<i>Leptosphaeria korrae</i>) or (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>) or (<i>Ophiosphaerella herpotricha</i>)	0.38-0.77	28	Make one or two applications in fall or when conditions are favorable for disease development.
Summer patch (<i>Magnaporthe poae</i>)	0.38-0.77	14-28	Apply when conditions are favorable for disease development.

*Do not apply more than two sequential applications of AZOXY 2SC SELECT™ for control of *Pythium* spp. For all other diseases, do not apply more than four sequential applications of AZOXY 2SC SELECT™.

(continued)

DIRECTIONS FOR APPLICATION FOR TURF DISEASES *(cont.)*

Target Diseases	Use Rate (fl. oz. product per 1,000 sq. ft.)	Application Interval (days)	Remarks*
Take-all patch <i>(Gaeumannomyces graminis var. avenae)</i>	0.38-0.77	28	Make two applications 28 days apart in the spring and two applications 28 days apart in the fall.
Zoysia patch <i>(Rhizoctonia solani and/or Gaeumannomyces incrustana)</i>	0.38-0.77	28	Make one or two applications in late fall before snow cover or when conditions are favorable for disease development. Do not apply on top of snow.

*Do not apply more than two sequential applications of AZOXY 2SC SELECT™ for control of *Pythium* spp. For all other diseases, do not apply more than four sequential applications of AZOXY 2SC SELECT™.

AZOXY 2SC SELECT™ Rate Conversion Chart for Turf

Fluid Ounces Product Per 1,000 Sq. Ft.	Ounces A.I. Per 1,000 Sq. Ft.	Fluid Ounces Product Per Acre	Pints of Product Per Acre
0.4	0.104	17.4	1.1
0.5	0.130	21.8	1.4
0.6	0.156	26.1	1.6
0.7	0.182	30.5	1.9
0.77	0.200	33.5	2.1
1.35	0.35	58.8	3.7

Amount of AZOXY 2SC SELECT™ to Mix 100 Gallons for Turf Applications

Spray Volume (gallons/1,000 square feet)			
AZOXY 2SC SELECT™ Use Rate (fl. oz.)	2.0 gals. (fl. oz.)	3.0 gals. (fl. oz.)	4.0 gals. (fl. oz.)
0.4	20	13	10
0.5	25	17	13
0.6	30	20	15
0.7	35	23	18
0.77	38.5	25.7	19.3
1.35	67.5	45	33.75

ORNAMENTALS

AZOXY 2SC SELECT™ controls certain pathogens causing foliar, aerial, and root diseases, including leaf, tip, and flower blights, leaf spots, downy mildew, powdery mildew, anthracnose, and rusts of ornamental plants. AZOXY 2SC SELECT™ controls certain diseases of container, bench, fiat, plug, bed or field-grown ornamentals in greenhouses, shade-houses, outdoor nurseries, retail nurseries, and other landscape areas.

INTEGRATED PEST (DISEASE) MANAGEMENT

Integrate AZOXY 2SC SELECT™ into an overall disease management strategy that includes selection of varieties with disease tolerance, optimum plant populations, proper fertilization, winter and/or spring pruning, plant residue management and proper timing and placement of irrigation. Immunoassay detection kits and diagnostic services can assist in the early and accurate identification of causal organisms and corresponding selection of the proper fungicide when required.

RESISTANCE MANAGEMENT

Some ornamental disease pathogens are known to have developed resistance to fungicides used repeatedly for their control. Apply AZOXY 2SC SELECT™ in an alternation or tank mix program with other registered fungicides that have a different mode of action and to which pathogen resistance has not developed. Do not make more than three (3) sequential applications of AZOXY 2SC SELECT™ before alternating with a fungicide of a different mode of action. A sound resistance management program includes blocks of three AZOXY 2SC SELECT™ applications separated by blocks of two alternate fungicide applications. Do not alternate AZOXY 2SC SELECT™ with other strobilurin fungicides.

APPLICATION DIRECTIONS

Apply AZOXY 2SC SELECT™ as a broadcast or banded spray targeted at the foliage or crown of the plant. Apply to runoff in sufficient water to ensure complete coverage of the target plant. Good coverage and wetting of foliage

is necessary for best control. Refer to the label for specific use directions for control of certain diseases. Repeat applications at specified intervals (plus alternations for resistance management) for as long as required. Applications may be made by ground only.

Start AZOXY 2SC SELECT™ applications prior to disease development and continue throughout the season at specified intervals following resistance management guidelines. AZOXY 2SC SELECT™ works best when used as part of a preventative disease management program.

Use only surfactants approved for ornamental plants in combination with AZOXY 2SC SELECT™. Do not use silicone based products with AZOXY 2SC SELECT™ due to possible phytotoxicity. Always test tank mixes on a small group of representative plants prior to broadscale use.

Apply 1.9 - 7.7 fl. oz./100 gallons (0.95 - 3.85 fl. oz./50 gallons) AZOXY 2SC SELECT™ every 7-28 days (or as otherwise specified for a specific plant or disease). The addition of a non-silicone based wetter-sticker at the specified use rate may enhance coverage on hard-to-wet plant foliage.

Under most conditions and for most diseases, apply 3.85 - 7.7 fl. oz./100 gallons (1.9 - 3.85 fl. oz./50 gallons) on a 7-14 day interval.

Under light to moderate disease pressure, use the lower rates within the specified rate range (1.9 - 3.85 fl. oz./100 gallons, or 0.95 - 1.9 fl. oz./50 gallons) on a 7-14 day interval or the higher rates within the specified rate range (5.75 - 7.7 fl. oz./100 gallons or 2.85 - 3.85 fl. oz./50 gallons) on a 14-28 day interval.

Under environmental conditions which promote severe disease development, use the higher rates within the specified rate range (5.75 - 7.7 oz./100 gallons or 2.85 - 3.85 fl. oz./50 gallons) on a 7-14 day interval.

Using AZOXY 2SC SELECT™ as a “rescue” (late curative or eradicator) treatment will not always result in satisfactory disease control.

Do not exceed 2.4 gallons of product/crop acre/year or 8 applications/crop/year.

Do not exceed 600 gallons spray volume per acre for foliar applications. For drench and crown applications, do not exceed 2 pints volume per square foot.

In addition, do not tank mix AZOXY 2SC SELECT™ with other fungicides, insecticides, herbicides, fertilizers, adjuvants, etc., unless local experience indicates that the tank mix is safe to ornamental plants.

DRENCH APPLICATION

Apply AZOXY 2SC SELECT™ to control soilborne, seedling, and crown diseases of production ornamentals (greenhouse, shadehouse, and container grown) as a preventative, drench treatment prior to infection. Good coverage of the pre-infection area (root zone, root ball, crown, etc.) is necessary for satisfactory control. Drench apply AZOXY 2SC SELECT™ to container grown ornamentals using 0.38 - 1.75 fl. oz./100 gallons of water. Apply 1-2 pints of the solution per square foot surface area on a 7-28 day interval. Apply drench prior to infection as healthy roots are necessary to optimize product uptake, systemic translocation and disease protection.

For resistance management do not make more than three sequential drench applications of AZOXY 2SC SELECT™ before alternating with a fungicide of a different mode of action.

Caution should be taken before making application of AZOXY 2SC SELECT™ as a drench to small bedding plants in the seedling/plug stage due to possible phytotoxicity. A limited quantity of plants should be tested prior to full-scale application.

DRIP IRRIGATION

Apply AZOXY 2SC SELECT™ through drip irrigation systems to potted ornamentals or to bedded, field grown ornamentals for soil-borne disease control. Apply 3.85 - 30.75 fl. oz. AZOXY 2SC SELECT™ per acre as a preventative

disease application. The soil or potting media should have adequate moisture capacity prior to drip application.

Terminate drip irrigation at fungicide depletion from the main feed supply tank or after 6 hours from start, whichever is shorter. For maximum efficacy, subsequent irrigation (water only) should be delayed for at least for 24 hours following drip application.

ORNAMENTAL USE RESTRICTIONS

- Do not apply AZOXY 2SC SELECT™ to apple or cherry trees (Flowering, Yoshino variety) due to possible phytotoxicity.
- Do not use spray equipment that has applied AZOXY 2SC SELECT™ for use in these sensitive crops due to possible phytotoxicity from residue remaining in the sprayer.

Apply AZOXY 2SC SELECT™ to certain varieties of crabapple for control of apple scab. AZOXY 2SC SELECT™ is safer when applied to the species and varieties listed in Table 4. However, due to the large number of genera, species, and varieties of crabapple, it is impossible to test every one for tolerance to AZOXY 2SC SELECT™. The professional user should conduct small scale testing to insure plant safety prior to broadscale commercial use on plant genera and species not listed on this label.

TABLE 1: DISEASES CONTROLLED

When used in accordance with the label directions, AZOXY 2SC SELECT™ will provide control of the following diseases of ornamental plants:

DISEASE (Pathogen)	Use Rates and Remarks	
	8 oz. and larger containers (fl. oz. product per 100 gallons)	4 oz. containers (fl. oz. product per 50 gallons)
1. CONIFER BLIGHTS		
a. Phomopsis Blight (<i>Phomopsis juniperovora</i>)	Apply 1.9 - 7.7 fl. oz. every 7-28 days.	Apply 0.95 - 3.85 fl. oz. every 7-28 days.
b. Tip Blight (<i>Sirococcus strobilinus</i>)	Apply 1.9 - 7.7 fl. oz. every 7-28 days.	Apply 0.95 - 3.85 fl. oz. every 7-28 days.
2. LEAF BLIGHTS/LEAF SPOTS		
a. Alternaria Leaf Spot (<i>Alternaria</i> spp.)	Apply 1.9 - 7.7 fl. oz. every 7-28 days.	Apply 0.95 - 3.85 fl. oz. every 7-28 days.
b. Anthracnose (<i>Colletotrichum</i> spp., <i>Elsinoe</i> spp.)	Apply 1.9 - 7.7 fl. oz. every 7-28 days.	Apply 0.95 - 3.85 fl. oz. every 7-28 days.
c. Downy Mildew of Rose (<i>Peronospora sparsa</i>)	Apply 3.85 - 7.7 fl. oz. every 7-21 days during periods of active plant growth and prior to dormancy or severe infection.	Apply 1.9 - 3.85 fl. oz. every 7-21 days during periods of active plant growth and prior to dormancy or severe infection.

(continued)

TABLE 1: DISEASES CONTROLLED (cont.)

DISEASE (Pathogen)	Use Rates and Remarks	
	8 oz. and larger containers (fl. oz. product per 100 gallons)	4 oz. containers (fl. oz. product per 50 gallons)
2. LEAF BLIGHTS/LEAF SPOTS (cont.)		
d. Entomosporium Leaf Spot (<i>Entomosporium mespili</i>)	Apply 1.9 - 7.7 fl. oz. every 7-28 days.	Apply 0.95 - 3.85 fl. oz. every 7-28 days.
e. Iris Leaf Spot (<i>Mycosphaerella macrospora</i>)	Apply 3.85 - 7.7 fl. oz. every 7-21 days.	Apply 1.9 - 3.85 fl. oz. every 7-21 days.
f. Leaf Spot (<i>Cladosporium echinulatum</i>)	Apply 1.9 - 7.7 fl. oz. every 7-28 days.	Apply 0.95 - 3.85 fl. oz. every 7-28 days.
g. Rose Blackspot (<i>Diplocarpon rosea</i>)	Apply 7.7 - 15.4 fl. oz. every 7-14 days. Apply AZOXY 2SC SELECT™ on a 7 day interval unless disease pressure is light. Under severe disease conditions or if disease is already present, AZOXY 2SC SELECT™ may be tank mixed with another rose blackspot fungicide. Do not exceed 46 fl. oz./acre application.	Apply 3.85 - 7.7 fl. oz. every 7-14 days. Apply AZOXY 2SC SELECT™ on a 7 day interval unless disease pressure is light. Under severe disease conditions or if disease is already present, AZOXY 2SC SELECT™ may be tank mixed with another rose blackspot fungicide. Do not exceed 46 fl. oz./acre/ application.

(continued)

TABLE 1: DISEASES CONTROLLED (cont.)

DISEASE (Pathogen)	Use Rates and Remarks	
	8 oz. and larger containers (fl. oz. product per 100 gallons)	4 oz. containers (fl. oz. product per 50 gallons)
2. LEAF BLIGHTS/LEAF SPOTS (cont.)		
h. Myrothecium Leaf Spot (<i>Myrothecium</i> spp.)	Apply 3.85 - 7.7 fl. oz. every 7-21 days.	Apply 1.9 - 3.85 fl. oz. every 7-21 days.
i. Downy Mildew of Bedding Plants (<i>Peronospora</i> spp.)	Apply 1.9 - 7.7 fl. oz. every 7-28 days.	Apply 0.95 - 3.85 fl. oz. every 7-28 days.
j. Scab (<i>Venturia inaequalis</i>)	Apply 1.9 - 7.7 fl. oz. every 10-28 days. Do not apply to apple trees. For crabapples only, see Table 4 for tolerant species.	Apply 0.95 - 3.85 fl. oz. every 10-28 days. Do not apply to apple trees. For crabapples only, see Table 4 for tolerant species.
k. Marssonina Leaf Spot (<i>Marssonina</i> spp.)	Apply 1.9 - 7.7 fl. oz./100 gals. every 14-28 days.	Apply 0.95 - 3.85 fl. oz. every 14-28 days.
l. Cercospora Leaf Spot	Apply 1.9 - 7.7 fl. oz./100 gals. every 7-28 days.	Apply 0.95 - 3.85 fl. oz. every 7-28 days.

(continued)

TABLE 1: DISEASES CONTROLLED (cont.)

DISEASE (Pathogen)	Use Rates and Remarks	
	8 oz. and larger containers (fl. oz. product per 100 gallons)	4 oz. containers (fl. oz. product per 50 gallons)
3. POWDERY MILDEW Preventative applications only. Do not make more than 2 sequential applications before rotating to another class of fungicide.		
a. <i>Erysiphe pannosa</i> , <i>E. spp.</i>	Apply 1.9 - 7.7 fl. oz. every 7-28 days.	Apply 0.95 - 3.85 fl. oz. every 7-28 days.
b. <i>Microsphaera azaleae</i>	Apply 1.9 - 7.7 fl. oz. every 7-28 days.	Apply 0.95 - 3.85 fl. oz. every 7-28 days.
c. <i>Sphaerotheca pannosa</i>	Apply 1.9 - 7.7 fl. oz. every 7-28 days.	Apply 0.95 - 3.85 fl. oz. every 7-28 days.
4. RUSTS		
a. Needle Rust (<i>Melampsora occidentalis</i>)	Apply 1.9 - 7.7 fl. oz. every 7-28 days.	Apply 0.95 - 3.85 fl. oz. every 7-28 days.
b. <i>Phragmidium spp.</i>	Apply 1.9 - 7.7 fl. oz. every 7-28 days.	Apply 0.95 - 3.85 fl. oz. every 7-28 days.
c. <i>Puccinia spp.</i>	Apply 1.9 - 7.7 fl. oz. every 7-28 days.	Apply 0.95 - 3.85 fl. oz. every 7-28 days.
d. <i>Gymnosporangium spp.</i>	Apply 1.9 - 7.7 fl. oz. every 7-28 days.	Apply 0.95 - 3.85 fl. oz. every 7-28 days.

(continued)

TABLE 1: DISEASES CONTROLLED (cont.)

DISEASE (Pathogen)	Use Rates and Remarks	
	8 oz. and larger containers (fl. oz. product per 100 gallons)	4 oz. containers (fl. oz. product per 50 gallons)
5. FLOWER BLIGHTS		
a. Anthracnose (<i>Colletotrichum</i> spp., <i>Elsinoe</i> spp.)	Apply 1.9 - 7.7 fl. oz. every 7-28 days.	Apply 0.95 - 3.85 fl. oz. every 7-28 days.
b. Botrytis Blight (<i>Botrytis cinerea</i>)	Apply 7.7 - 15.4 fl. oz. every 7-21 days. For suppression only. Do not exceed 46 fl. oz./acre.	Apply 3.85 - 7.7 fl. oz. every 7-21 days. For suppression only. Do not exceed 46 fl. oz./acre.
6. SHOOT/STEM DISEASES		
a. Aerial/Shoot Blight (<i>Phytophthora</i> spp.)	Apply 1.9 - 3.85 fl. oz. every 7-28 days.	Apply 0.95 - 1.9 fl. oz. every 7-28 days.
7. SOILBORNE DISEASES (Directed Spray) For directed spray applications utilize the following rates below.		
a. <i>Rhizoctonia solani</i>	Apply 1.9 - 7.7 fl. oz. every 7-21 days.	Apply 0.95 - 3.85 fl. oz. every 7-21 days.
b. <i>Sclerotium rolfsii</i>	Apply 1.9 - 7.7 fl. oz. every 7-21 days.	Apply 0.95 - 3.85 fl. oz. every 7-21 days.
c. <i>Rosarium</i> spp.	Apply 1.9 - 7.7 fl. oz. every 7-21 days.	Apply 0.95 - 3.85 fl. oz. every 7-21 days.

(continued)

TABLE 1: DISEASES CONTROLLED (cont.)

DISEASE (Pathogen)	Use Rates and Remarks	
	8 oz. and larger containers (fl. oz. product per 100 gallons)	4 oz. containers (fl. oz. product per 50 gallons)
8. SOILBORNE DISEASES (Drench) See Ornamentals section for additional drench directions.		
a. <i>Rhizoctonia solani</i>	Apply 0.35 - 1.75 fl. oz., 1-2 pints of the solution per square foot surface area, every 7-28 days.	Apply 0.19 - 0.95 fl. oz., 1-2 pints of the solution per square foot surface area, every 7-28 days.
b. <i>Sclerotium rolfsii</i>	Apply 0.35 - 1.75 fl. oz., 1-2 pints of the solution per square foot surface area, every 7-28 days.	Apply 0.19 - 0.95 fl. oz., 1-2 pints of the solution per square foot surface area, every 7-28 days.
c. <i>Fusarium</i> spp.	Apply 0.35 - 1.75 fl. oz., 1-2 pints of the solution per square foot surface area, every 7-28 days.	Apply 0.19 - 0.95 fl. oz., 1-2 pints of the solution per square foot surface area, every 7-28 days.

PLANT SAFETY

AZOXY 2SC SELECT™ is safe when applied to the ornamental plants listed in Tables 2, 3, and 4; however, due to the large number of genera, species and varieties of ornamental and nursery plants, it is impossible to test every one for tolerance to AZOXY 2SC SELECT™. Neither the manufacturer nor the seller has determined whether or not AZOXY 2SC SELECT™ can be used safely on genera, species, or varieties of ornamental and nursery plants not specified on this label. The professional user should conduct small scale

testing to insure plant safety prior to broadscale commercial use on plant genera and species not listed in this label.

Do not tank mix AZOXY 2SC SELECT™ with other fungicides, insecticides, herbicides, fertilizer, adjuvants, etc., unless local experience indicates that the tank mix is safe to ornamental plants.

Do not apply AZOXY 2SC SELECT™ to certain apple, crabapple or cherry trees due to possible phytotoxicity. Further, do not use spray equipment that has applied AZOXY 2SC SELECT™ for use in these sensitive crops due to possible phytotoxicity from residue remaining in the sprayer.

Tolerant Ornamental Plants

AZOXY 2SC SELECT™ is safe when applied to the plants listed in Tables 2, 3, and 4 when applied according to specified application methods, rates, and timings:

TABLE 2: Tolerant Plants Listed by Botanical Name

BOTANICAL NAME	COMMON NAME	DISEASES
<i>Abelia</i> spp.	Abelia	2
<i>Abies fraseri</i>	Fraser fir	1, 4
<i>Abies procera</i>	Noble fir	1, 4
<i>Acer palmatum</i>	Japanese maple	2
<i>Acer saccharum</i>	Sugar maple	2
<i>Ageratum</i> spp.	Floss-Flower	3, 4
<i>Ageratum</i> spp.	Pussy's-Foot	3, 4
<i>Aglaonema</i> spp.	Chinese evergreen	2, 4
<i>Ajuga reptans</i>	Bugle, Bugleweed	3

(continued)

TABLE 2: Tolerant Plants Listed by Botanical Name (cont.)

BOTANICAL NAME	COMMON NAME	DISEASES
<i>Antirrhinum</i> spp.	Snap-Dragon	2i, 3, 4
<i>Aphelandra</i> spp.	Zebra-Plant	2
<i>Artemisia</i> spp.	Mugwort, Sagebrush	2
<i>Artemisia</i> spp.	Wormwood	2
<i>Aster</i> spp.	Aster, Starwort	4
<i>Aucuba japonica</i>	Japanese aucuba, Japanese laurel	7
<i>Begonia</i> spp. (except <i>Rieger begonia</i>)	Begonia	2, 3
<i>Berberis thunbergii</i>	Barberry	3, 4
<i>Betula nigra</i>	River birch	3, 4
<i>Bougainvillea</i> spp.	Bougainvillea	2
<i>Brassaia actinophyllia</i>	Rubber-tree, Umbrella-tree	2, 7
<i>Buddleia davidii</i>	Buddleia, Butterfly bush	2
<i>Buxus sempervirens</i>	Boxwood	2, 7a
<i>Caladium</i> spp.	Caladium	7
<i>Camellia japonica</i>	Camellia	2
<i>Caryota urens</i>	Sago palm	2, 7
<i>Catharanthus roseus</i>	Vinca	2
<i>Ceanothus sanguineus</i>	Wild lilac	3

(continued)

TABLE 2: Tolerant Plants Listed by Botanical Name (cont.)

BOTANICAL NAME	COMMON NAME	DISEASES
<i>Ceanothus</i> spp.	Ceanothus, California lilac, Snowball	3
<i>Cedrus Atlantica</i>	Atlas cedar	2, 4
<i>Cedrus</i> spp.	White cedar	2, 4
<i>Cercis occidentalis</i>	Western redbud	2
<i>Chamaecyparis</i> spp.	Cypress, Leyland cypress	1
<i>Chamaecyparis pisifera</i> spp.	Sawara cypress	1
<i>Chamaedora elegans</i>	Parlor palm	7
<i>Chrysanthemum</i> spp.	Chrysanthemums	2, 7c
<i>Clethra alnifolia</i>	Clethra, White alder	2
<i>Cornus</i> spp.	Dogwood, Pink Dogwood, Flowering Dogwood	2b, 3
<i>Cornus florida</i>	Dogwood	2b, 3
<i>Cortaderia selloana</i>	Pampas grass	3
<i>Cotoneaster adpressus</i>	Creeping cotoneaster	7
<i>Cotoneaster horizontalis</i>	Cotoneaster - variegated rockspray	7
<i>Cyclamen</i> spp.	Cyclamen	7c
<i>Cyperus</i> spp.	Cyperus	1

(continued)

TABLE 2: Tolerant Plants Listed by Botanical Name (cont.)

BOTANICAL NAME	COMMON NAME	DISEASES
<i>Delphinium</i> spp.	Larkspur	2
<i>Dianthus caryophyllus</i>	Carnation	3, 4
<i>Dianthus</i> spp.	Pink	3, 4
<i>Dieffenbachia</i> spp.	Dumb-Cane	2
<i>Dietes iridioides</i>	African iris, Butterfly iris	4c
<i>Digitalis</i> spp.	Foxglove	2, 3
<i>Epipremnum</i> spp.	Pothos	2
<i>Erica darleyensis</i>	Heather	2
<i>Euonymus alata</i>	Dwarf winged euonymus	2
<i>Euonymus alatus</i>	Burning bush	2
<i>Euonymus japonicus</i>	Evergreen euonymus	2
<i>Euphorbia</i> spp.	Poinsettia	2a
<i>Fatsia japonica</i>	Japanese fatsia, Paper-plant	2
<i>Ficus</i> spp.	Fig	2
<i>Forsythia viridissima</i>	Forsythia	2
<i>Gaillardia</i> spp.	Blanket flower	2
<i>Gardenia jasminoides</i>	Gardenia	3
<i>Geranium</i> spp.	Cranesbill	5b

(continued)

TABLE 2: Tolerant Plants Listed by Botanical Name (cont.)

BOTANICAL NAME	COMMON NAME	DISEASES
<i>Gerbera jamesonii</i>	Gerber daisy, Transvaal daisy	3
<i>Hedera algeriensis</i>	Algerian ivy	2
<i>Hedera helix</i>	English ivy	2
<i>Hibiscus moscheutos</i>	Hibiscus	2, 3
<i>Hibiscus rosa-sinensis</i>	Hibiscus	2, 3
<i>Hibiscus syriacus</i>	Rose of Sharon	2, 3
<i>Hosta</i> spp.	Hosta	2
<i>Hydrangea macrophylla</i>	French hydrangea	2, 3
<i>Hydrangea</i> spp.	Hydrangea	2, 3
<i>Ilex</i> spp.	Holly, Winterberry, Yaupon	3
<i>Impatiens</i> spp. ¹	Balsam, Impatiens ¹	2a, 7a
<i>Iris xiphium</i>	Iris (bulbous, Spanish, Dutch)	2e
<i>Itea virginica</i>	Virginia willow	3, 4
<i>Juniperus procumbens</i>	Juniper	1a, 4
<i>Juniperus scopulorum</i>	Juniper	1a, 4
<i>Juniperus</i> spp.	Juniper	1a, 4
<i>Juniperus virginiana</i>	Red cedar	1a, 4

¹Do not exceed 3.85 fl. oz./100 gallons on these species.

(continued)

TABLE 2: Tolerant Plants Listed by Botanical Name (cont.)

BOTANICAL NAME	COMMON NAME	DISEASES
<i>Lagerstroemia indica</i>	Crapemyrtle	2, 3
<i>Laurus nobilis</i>	Laurel	3
<i>Lilium</i> spp.	Asiatic lily	2
<i>Liriope muscari</i>	Lily-turf	2
<i>Lobularia maritima</i>	Sweet alyssum	7
<i>Magnolia grandiflora</i>	Southern magnolia	2
<i>Magnolia soulangiana</i>	Saucer magnolia	2
<i>Magnolia</i> spp.	Magnolia	2
<i>Malus</i> spp.	Crabapple (See Table 4 for variety list)	2i
<i>Nandina domestica</i>	Nandina	2
<i>Nerium oleander</i>	Oleander, Rose-bay	2
<i>Pelargonium</i> spp.	Geranium	3, 4, 5b
<i>Pennisetum alopecuroides</i>	Grass	2
<i>Peperomia</i> spp.	Baby rubber-plant	2, 7
<i>Petunia</i> spp.	Petunia	6a
<i>Phalaris</i> spp.	Dwarf pampas grass	3
<i>Philodendron</i> spp.	Philodendron	2j

(continued)

TABLE 2: Tolerant Plants Listed by Botanical Name (cont.)

BOTANICAL NAME	COMMON NAME	DISEASES
<i>Phlox</i> spp.	Phlox	3
<i>Phoenix dactylifera</i>	Date palm	2, 7
<i>Phoenix roebelenii</i>	Roebelin's palm	2, 7
<i>Photinia glabra</i>	Red tip photinia	2, 3, 4
<i>Picea abies</i>	Norway spruce	1
<i>Picea glauca</i>	White spruce	1
<i>Picea pungens</i>	Blue spruce	1
<i>Pieris japonica</i>	Japanese andromeda	2, 7
<i>Pinus muhgo</i>	Muhgo pine	1b, 4
<i>Pinus nigra</i>	Black pine	1b, 4
<i>Pinus silvestris</i>	Scotch pine	1, 4
<i>Pinus</i> spp.	Pine	1b, 4
<i>Pinus strobus</i>	Eastern white pine	1b, 4
<i>Pittosporum</i> spp.	Australian laurel	3, 4
<i>Pittosporum tobira</i>	Mock-orange	3, 4
<i>Plectranthus</i> spp.	Swedish ivy, Coleus	2
<i>Populus trichocarpa</i>	Poplar	4
<i>Populus</i> spp.	Aspen Trees	2
<i>Potentilla</i> spp.	Cinquefoil	2

(continued)

TABLE 2: Tolerant Plants Listed by Botanical Name (cont.)

BOTANICAL NAME	COMMON NAME	DISEASES
<i>Primula</i> spp.	Primrose	2
<i>Prunus pumila</i>	Cherry	2, 5
<i>Prunus</i> spp.	Flowering plum, Purple-leaf plum	2, 5
<i>Pseudotsuga</i> spp.	Douglas fir	1, 4
<i>Pyrus calleryana</i>	Bradford's pear	3
<i>Quercus falcata</i>	Red oak	2, 3
<i>Quercus palustris</i>	Pin oak	2, 3
<i>Rhaphiolepis indica</i>	Indian hawthorn	2, 3, 4
<i>Rhododendron</i> spp.	Azaleas, Rhododendron	2b, 3, 6, 7
<i>Rhododendron</i> spp.	Glacier Azalea	2b, 3, 6, 7
<i>Rosa</i> spp.	Rose	2a, 2c, 3c, 4b
<i>Rosmarinus</i> spp.	Rosemary (prostrate)	2
<i>Rudbeckia hirta</i>	Black-eyed Susan	2j
<i>Salvia</i> spp.	Sage	3, 4j
<i>Schlumbergera</i>	Holiday cactus	2, 7
<i>Sedum</i> spp.	Orpine, Stonecrop	2
<i>Sempervivum</i> spp.	Live-forever, House-Leek	2
<i>Setaria</i> spp.	Ribbon Grass	2, 3

(continued)

TABLE 2: Tolerant Plants Listed by Botanical Name (cont.)

BOTANICAL NAME	COMMON NAME	DISEASES
<i>Spathiphyllum floribundum</i>	Peace lily	2, 7
<i>Spiraea bumalda</i>	Spirea	3
<i>Spiraea japonica</i>	Spirea	3
<i>Syagrus romanzoffianum</i>	Queen palm	2
<i>Tagetes</i> spp.	Marigold	2a
<i>Taxus baccata</i>	Spreading yew	7
<i>Thuja plicata</i>	Western red cedar	4
<i>Thujopsis</i> spp.	Arborvitae	2
<i>Thymus serpyllum</i>	Creeping thyme	2
<i>Tsuga heterophylla</i>	Western hemlock	4
<i>Tsuga</i> spp.	Hemlock	4
<i>Verbena</i> spp.	Verbena, Vervain	3
<i>Viburnum</i> spp.	Viburnum	2, 3, 4
<i>Vinca</i> spp.	Periwinkle	2, 6a
<i>Viola</i> spp. ¹	Viola, Pansy ¹	2
<i>Weigela Florida</i>	Pink Weigela	2
<i>Yucca</i> spp.	Yucca	7
<i>Zinnia</i> spp.	Zinnia	2a, 3

¹Do not exceed 3.85 fl. oz./100 gallons on these species.

TABLE 3: Tolerant Plants Listed by Common Name

COMMON NAME	BOTANICAL NAME
Abelia	<i>Abelia</i> spp.
Andromeda Japanese	<i>Pieris japonica</i>
Arborvitae	<i>Thuja</i> spp.
Aspen Trees	<i>Populus</i> spp.
Aster	<i>Aster</i> spp.
Aucuba, Japanese	<i>Aucuba japonica</i>
Azalea, Glacier	<i>Rhododendron</i> spp.
Azaleas	<i>Rhododendron</i> spp.
Balsam	<i>Impatiens</i> spp.
Barberry	<i>Berberis thunbergii</i>
Begonia (except Rieger begonia)	<i>Begonia</i> spp.
Birch, River	<i>Betula nigra</i>
Black-eyed Susan	<i>Rudbeckia hirta</i>
Blanket Flower	<i>Gaillardia</i> spp.
Bougainvillea	<i>Bougainvillea</i> spp.
Boxwood	<i>Buxus sempervirens</i>
Buddleia	<i>Buddleia davidii</i>
Bugle	<i>Ajuga reptans</i>
Bugleweed	<i>Ajuga reptans</i>

(continued)

TABLE 3: Tolerant Plants Listed by Common Name (cont.)

COMMON NAME	BOTANICAL NAME
Burning Bush	<i>Euonymus alatus</i>
Butterfly Bush	<i>Buddleia davidii</i>
Cactus, Holiday	<i>Schlumbergera</i>
Caladium	<i>Caladium</i> spp.
Camellia	<i>Camellia japonica</i>
Carnation	<i>Dianthus caryophyllus</i>
Ceanothus	<i>Ceanothus</i> spp.
Cedar, Atlas	<i>Cedrus atlantica</i>
Cedar, Red	<i>Juniperus virginiana</i>
Cedar, Western Red	<i>Thuja plicata</i>
Cedar, White	<i>Cedrus</i> spp.
Cherry	<i>Prunus pumila</i>
Christmas Tree	See Fraser fir, Scotch pine, and Douglas fir
Chrysanthemum	<i>Chrysanthemum</i> spp.
Cinquefoil	<i>Potentilla</i> spp.
Clethra	<i>Clethra alnifolia</i>
Coleus	<i>Plectranthus</i> spp.
Cotoneaster, Creeping	<i>Cotoneaster adpressus</i>

(continued)

TABLE 3: Tolerant Plants Listed by Common Name (cont.)

COMMON NAME	BOTANICAL NAME
Cotoneaster, Variegated Rockspray	<i>Cotoneaster horizontalis</i>
Crabapple (See Table 4 for variety list)	<i>Malus</i> spp.
Cranesbill	<i>Geranium</i> spp.
Crapemyrtle	<i>Lagerstroemia indica</i>
Cyclamen	<i>Cyclamen</i> spp.
Cyperus	<i>Cyperus</i> spp.
Cypress, Sawara	<i>Chamaecyparis pisifera</i>
Cypress, Leyland	<i>Chamaecyparis</i> spp.
Daisy, Gerber	<i>Gerbera jamesonii</i>
Daisy, Transvaal	<i>Gerbera jamesonii</i>
Dogwood	<i>Cornus</i> spp.
Dogwood	<i>Cornus florida</i>
Dogwood, Pink	<i>Cornus</i> spp.
Dumb-Cane	<i>Dieffenbachia</i> spp.
Euonymus, Dwarf Winged	<i>Euonymus alata</i>
Euonymus, Evergreen	<i>Euonymus japonicus</i>
Evergreen, Chinese	<i>Aglaonema</i> spp.
Fatsia, Japanese	<i>Fatsia japonica</i>

(continued)

TABLE 3: Tolerant Plants Listed by Common Name (cont.)

COMMON NAME	BOTANICAL NAME
Fig	<i>Ficus</i> spp.
Fir, Douglas	<i>Pseudotsuga</i> spp.
Fir, Fraser	<i>Abies fraseri</i>
Fir, Noble	<i>Abies procera</i>
Floss-Flower	<i>Ageratum</i> spp.
Forsythia	<i>Forsythia viridissima</i>
Foxglove	<i>Digitalis</i> spp.
Gardenia	<i>Gardenia jasminoides</i>
Geranium	<i>Pelargonium</i> spp.
Grass	<i>Pennisetum alopecuroides</i>
Grass, Dwarf Pampas	<i>Phalaris</i> spp.
Grass, Pampas	<i>Cortaderia selloana</i>
Hawthorn, Indian	<i>Rhaphiolepis indica</i>
Heather	<i>Erica darleyensis</i>
Hemlock	<i>Tsuga</i> spp.
Hemlock, Western	<i>Tsuga heterophylla</i>
Hibiscus	<i>Hibiscus moscheutos</i>
Hibiscus	<i>Hibiscus rosa-sinensis</i>
Holly	<i>Ilex</i> spp.

(continued)

TABLE 3: Tolerant Plants Listed by Common Name (cont.)

COMMON NAME	BOTANICAL NAME
Hosta	<i>Hosta</i> spp.
House-Leek	<i>Sempervivum</i> spp.
Hydrangea	<i>Hydrangea</i> spp.
Hydrangea, French	<i>Hydrangea macrophylla</i>
Impatiens ¹	<i>Impatiens</i> spp. ¹
Iris (Bulbous, Spanish, Dutch)	<i>Iris xiphium</i>
Iris, African	<i>Dietes iridioides</i>
Iris, Butterfly	<i>Dietes iridioides</i>
Ivy, Algerian	<i>Hedera algeriensis</i>
Ivy, English	<i>Hedera helix</i>
Ivy, Swedish	<i>Plectranthus</i> spp.
Juniper	<i>Juniperus procumbens</i>
Juniper	<i>Juniperus scopulorum</i>
Juniper	<i>Juniperus</i> spp.
Larkspur	<i>Delphinium</i> spp.
Laurel	<i>Laurus nobilis</i>
Laurel, Australian	<i>Pittosporum</i> spp.
Laurel, Japanese	<i>Aucuba japonica</i>

¹Do Not Exceed 3.85 fl. oz./100 gallons on these species.

(continued)

TABLE 3: Tolerant Plants Listed by Common Name (cont.)

COMMON NAME	BOTANICAL NAME
Lilac, California	<i>Ceanothus</i> spp.
Lilac, Wild	<i>Ceanothus sanguineus</i>
Lily, Asiatic	<i>Lilium</i> spp.
Lily, Peace	<i>Spathiphyllum floribundum</i>
Lily-Turf	<i>Liriope muscari</i>
Live-Forever	<i>Sempervivum</i> spp.
Magnolia	<i>Magnolia</i> spp.
Magnolia, Saucer	<i>Magnolia soulangiana</i>
Magnolia, Southern	<i>Magnolia grandiflora</i>
Maple, Japanese	<i>Acer palmatum</i>
Maple Sugar	<i>Acer saccharum</i>
Marigold	<i>Tagetes</i> spp.
Mock-Orange	<i>Pittosporum tobira</i>
Mugwort	<i>Artemisia</i> spp.
Nandina	<i>Nandina domestica</i>
Oak, Pin	<i>Quercus palustris</i>
Oak, Red	<i>Quercus falcata</i>
Oleander	<i>Nerium oleander</i>
Orpine	<i>Sedum</i> spp.

(continued)

TABLE 3: Tolerant Plants Listed by Common Name (cont.)

COMMON NAME	BOTANICAL NAME
Palm, Date	<i>Phoenix dactylifera</i>
Palm, Parlor	<i>Chamaedorea elegans</i>
Palm, Queen	<i>Syagrus romanzoffianum</i>
Palm, Roebelin's	<i>Phoenix roebelenii</i>
Palm, Sago	<i>Caryota urens</i>
Pansy ¹	<i>Viola</i> spp. ¹
Paper Plant	<i>Fatsia japonica</i>
Pear Bradford's	<i>Pyrus calleryana</i>
Periwinkle	<i>Vinca</i> spp.
Petunia	<i>Petunia</i> spp.
Philodendron	<i>Philodendron</i> spp.
Phlox	<i>Phlox</i> spp.
Photinia, Red-Tip	<i>Photinia glabra</i>
Pine	<i>Pinus</i> spp.
Pine, Black	<i>Pinus nigra</i>
Pine, Eastern White	<i>Pinus strobus</i>
Pine, Muhgo	<i>Pinus Muhgo</i>
Pine Scotch	<i>Pinus Sylvestris</i>

¹Do Not Exceed 3.85 fl. oz./100 gallons on these species.

(continued)

TABLE 3: Tolerant Plants Listed by Common Name (cont.)

COMMON NAME	BOTANICAL NAME
Pink	<i>Dianthus</i> spp.
Plum, Flowering	<i>Prunus</i> spp.
Plum, Purple-Leaf	<i>Prunus</i> spp.
Poinsettia	<i>Euphorbia</i> spp.
Poplar	<i>Populus trichocarpa</i>
Pothos	<i>Epipremnum</i> spp.
Primrose	<i>Primula</i> spp.
Pussy's-Foot	<i>Ageratum</i> spp.
Redbud, Western	<i>Cercis occidentalis</i>
Rhododendron	<i>Rhododendron</i> spp.
Ribbon-Grass	<i>Setaria</i> spp.
Rose of Sharon	<i>Hibiscus syriacus</i>
Rose	<i>Rosa</i> spp.
Rose-Bay	<i>Nerium oleander</i>
Rosemary (Prostrate)	<i>Rosmarinus</i> spp.
Rubber-Plant, Baby	<i>Peperomia</i> spp.
Rubber Tree	<i>Brassaia actinophylla</i>
Sage	<i>Salvia</i> spp.
Sagebrush	<i>Artemisia</i> spp.

(continued)

TABLE 3: Tolerant Plants Listed by Common Name (cont.)

COMMON NAME	BOTANICAL NAME
Snap-Dragon	<i>Antirrhinum</i> spp.
Snowball	<i>Ceanothus</i> spp.
Spirea	<i>Spiraea bumalda</i>
Spirea	<i>Spiraea japonica</i>
Spruce, Blue	<i>Picea pungens</i>
Spruce, Norway	<i>Picea abies</i>
Spruce, White	<i>Picea glauca</i>
Starwort	<i>Aster</i> spp.
Stonecrop	<i>Sedum</i> spp.
Sweet Alyssum	<i>Lobularia maritima</i>
Thymes, Creeping	<i>Thymus serpyllum</i>
Umbrella, Tree	<i>Brassaia actinophylla</i>
Verbena	<i>Verbena</i> spp.
Vervain	<i>Verbena</i> spp.
Viburnum	<i>Viburnum</i> spp.
Vinca	<i>Catharanthus roseus</i>
Viola	<i>Viola</i> spp.
White alder	<i>Clethra</i> spp.
Weigela, Pink	<i>Weigela Florida</i>

(continued)

TABLE 3: Tolerant Plants Listed by Common Name (*cont.*)

COMMON NAME	BOTANICAL NAME
Willow, Virginia	<i>Itea virginica</i>
Winterberry	<i>Ilex</i> spp.
Wormwood	<i>Artemisia</i> spp.
Yaupon	<i>Ilex</i> spp.
Yew, Spreading	<i>Taxus baccata</i>
Yucca	<i>Yucca</i> spp.
Zebra-Plant	<i>Aphelandra</i> spp.
Zinnia	<i>Zinnia</i> spp.

TABLE 4: Tolerant Varieties of Crabapple Species (Genus *Malus*)
Tolerant Varieties of *Malus*

Arkansas Black	Candymint Sargent	Dorothea	<i>floribunda</i>
<i>atrosanguinea</i>	Christmas Holly	Doubloons	Gloriosa
<i>baccata</i>	<i>coronaria</i>	Eleyi	Golden Delicious
<i>baccata</i> var. <i>jackii</i>	David	Enterprise	Golden Raindrops
<i>baccata</i> var. <i>mandshurica</i>	Dolgo	Evereste	Hopa
Callaway	Donald Wyman	Eyelynn	Indian Magic
Island	Pink Satin	Sargent	<i>spectabilis</i>
Katherine	Prairie Maid	<i>sargentii</i>	Sugar Tyme

(continued)

TABLE 4: Tolerant Varieties of Crabapple Species (Genus *Malus*)
Tolerant Varieties of *Malus* (cont.)

Lancelot	Prairifire	<i>sieboldii</i>	Van Eseltine
Louisa	Profusion	Selkirk	White Angel
Mary Potter	<i>pumila</i>	Sentinel	Williams Pride
Molten Lava	Ralph Shay	Silver Moon	Winter Gold
New Centennial	Red Jade	Silver Drift	Yellow Delicious
Ormiston Roy	Red Baron	Sinai Fire	<i>zumi</i> Calocarpa

TABLE 5. Intolerant Plants
(DO NOT apply AZOXY 2SC SELECT™ to these species or varieties)

COMMON NAME	BOTANICAL NAME
Apple	<i>Malus domestics</i>
Crabapple - Flame variety	<i>Malus</i> spp.
Crabapple - Brandywine variety	<i>Malus</i> spp.
Crabapple - Novamac variety	<i>Malus</i> spp.
Cherry, Flowering - Yoshino variety	<i>Prunus yedoensis</i>
Leatherleaf Fern and Other Ferns for cut foliage	<i>Rumohra adianformis</i> and other species for cut foliage
Privet	<i>Ligustrum</i> spp.

CONIFERS INCLUDING CHRISTMAS TREES, COMMERCIAL PRODUCTION ROSES

AZOXY 2SC SELECT™ controls certain diseases on conifers in production (indoor and outdoor) and landscape situations.

Please see the Ornamental section above for more detailed directions for use in landscape situations.

Crop	Target Diseases	Use Rate fl. oz. product/ Acre (lb. a.i./A)	Remarks
Conifers including Christmas Trees	Diplodia Tip Blight <i>(Diplodia pinea)</i> Lophodermium Needlecast <i>(Lophodermium pinastri)</i> Swiss Needlecast <i>(Phaeocryptopus gaumannii)</i>	6.1 - 15.3 (0.10 - 0.25)	<p>Integrated Pest (Disease) Management: Integrate AZOXY 2SC SELECT™ into an overall disease management strategy that includes selection of varieties with disease tolerance and removal of plant debris in which inoculum may overwinter.</p> <p>Resistance Management: Do not apply more than four sequential applications of AZOXY 2SC SELECT™ before alternating with a fungicide that is not in Group 11. Do not make more than eight applications of AZOXY 2SC SELECT™ per acre per year.</p>

**CONIFERS INCLUDING CHRISTMAS TREES,
COMMERCIAL PRODUCTION ROSES (cont.)**

Crop	Target Diseases	Use Rate fl. oz. product/ Acre (lb. a.i./A)	Remarks
Conifers including Christmas Trees (cont.)			<p>Application Directions: Begin AZOXY 2SC SELECT™ applications prior to disease development and continue throughout the season at 7-21 day intervals following the resistance management guidelines. Make applications by ground, air or chemigation. An adjuvant may be added at specified rates.</p>

(continued)

**CONIFERS INCLUDING CHRISTMAS TREES,
COMMERCIAL PRODUCTION ROSES (cont.)**

Crop	Target Diseases	Use Rate fl. oz. product/ Acre (lb. a.i./A)	Remarks
Roses (Commercial Rose Production)	Downy Mildew (<i>Peronospora sparsa</i>) Powdery Mildew (<i>Sphaerotheca pannosa</i>) Rust (<i>Phragmidium mucronatum</i> , <i>P. tuberculatum</i> , and other <i>Phragmidium spp.</i>) Septoria Leaf Spot (<i>Septoria rosea</i>)	3.0 - 15.3 (0.05 - 0.25)	Integrated Pest (Disease) Management: Integrate AZOXY 2SC SELECT™ into an overall disease manage- ment strategy that includes selection of varieties with disease tolerance, optimum plant populations, proper fertilization, winter and/or spring pruning, plant residue management and proper timing and placement of irrigation.

(continued)

**CONIFERS INCLUDING CHRISTMAS TREES,
COMMERCIAL PRODUCTION ROSES (cont.)**

Crop	Target Diseases	Use Rate fl. oz. product/ Acre (lb. a.i./A)	Remarks
Roses (Commercial Rose Production) (cont.)	(cont.) Alternaria Leaf Spot (<i>Alternaria alternata</i>)		<p>Resistance Management: Do not make more than four sequential applications of AZOXY 2SC SELECT™ before alternating with a fungicide that is not in Group 11. Do not make more than eight applications per acre per year.</p> <p>Application Directions: Begin AZOXY 2SC SELECT™ application prior to disease development and continue throughout the season on 7-21 day intervals following the resistance management guidelines. Make applications by ground, air or chemigation. An adjuvant may be added at specified rates.</p>

(continued)

**CONIFERS INCLUDING CHRISTMAS TREES,
COMMERCIAL PRODUCTION ROSES (cont.)**

Crop	Target Diseases	Use Rate fl. oz. product/ Acre (lb. a.i./A)	Remarks
Roses (Commercial Rose Production) (cont.)			Plant Safety: AZOXY 2SC SELECT™ is safe when applied to roses. However, all varieties of roses have not been evaluated for safety. Small scale variety safety testing must be conducted to insure plant safety prior to large scale application. In addition, do not tank mix AZOXY 2SC SELECT™ with other fungicides, insecticides, herbicides, fertilizer, etc. unless local experience indicates that the tank mix is safe to roses.

Specific Use Restrictions: Do not apply more than 123 fluid ounces of product/acre/season (2.0 lbs. a.i./A).

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

CONTAINER HANDLING [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of PRIME SOURCE, A DIVISION OF ALBAUGH, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold PRIME SOURCE, A DIVISION OF ALBAUGH, LLC and Seller harmless for any claims relating to such factors.

PRIME SOURCE, A DIVISION OF ALBAUGH, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or PRIME SOURCE, A DIVISION OF ALBAUGH, LLC, and, (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, PRIME SOURCE, A DIVISION OF ALBAUGH, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall PRIME SOURCE, A DIVISION OF ALBAUGH, LLC be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF PRIME SOURCE, A DIVISION OF ALBAUGH, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF PRIME SOURCE, A DIVISION OF ALBAUGH, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PRIME SOURCE, A DIVISION OF ALBAUGH, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of PRIME SOURCE, A DIVISION OF ALBAUGH, LLC.

Ambush[®], Callisto[®], Halex[®], Plant Performance[™], Warrior with Zeon Technology[®], the ALLIANCE FRAME, the SYNGENTA Logo and the PURPOSE ICON, are Trademarks of a Syngenta Group Company.

Acrobat[®] is a trademark of BASF Corporation.

Aliette[®] and Phaser[®] are trademarks of Bayer CropScience.

Botran[®] is a trademark of Gowan Company.

Lorsban[®] and Kelthane[®] are trademarks of Dow AgroSciences, LLC.

Lannate[®] is a trademark of DuPont Crop Protection.

M-Pede[®] is a trademark of Mycogen Corporation.

Pounce[®] is a trademark of FMC Corporation and Agrilience, LLC.