

# SPECIMEN LABEL

# BUTYRAC<sup>®</sup> 175

*Controls Broadleaf Weeds in Soybeans and Peanuts*

**ACTIVE INGREDIENT:**

4-(2,4-Dichlorophenoxy)butyric acid, dimethylamine salt\*.....23.0%

**OTHER INGREDIENTS** .....77.0%

**TOTAL**..... 100.00%

\*4-(2,4-Dichlorophenoxy)butyric acid equivalent 19.5% by weight or 1.75 pounds per gallon.

**KEEP OUT OF REACH OF CHILDREN  
DANGER/PELIGRO**

**Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)**

See inside booklet for complete Precautionary Statements, Directions For Use, Storage and Disposal, and Conditions of Sale and Warranty.

For 24-hour chemical spill, leak, fire, exposure or accident response information, call CHEMTREC toll free at 1-800-424-9300.

**FIRST AID**

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. **HOT LINE NUMBER:** Have the product container or label with you when calling a poison control center (1-800-222-1222) or doctor or going for treatment For non-emergency exposure information on this product, call 1-888-347-6732 (7 days/week, 24-hr) For medical emergencies, dial 911 **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

Manufactured by:  
**ALBAUGH, LLC**

1525 NE 36th Street, Ankeny, IA 50021



EPA Reg. No. 42750-39  
AD081810A

---

## PRECAUTIONARY STATEMENTS

---

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

# DANGER

**DANGER.** Corrosive. Causes irreversible eye damage. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin.

---

## PERSONAL PROTECTIVE EQUIPMENT (PPE)

---

Some materials that are chemical-resistant to this product are butyl rubber, natural rubber, neoprene or nitrile. If you want more options, follow the instructions for Category A on an EPA chemical-resistance category selection chart.

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves
- Protective eyewear
- Chemical-resistant apron when mixing/loading, cleaning up spills, cleaning equipment, or otherwise exposed to concentrate.

See “Engineering Controls” for additional requirements.

---

## USER SAFETY REQUIREMENTS

---

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product’s concentrate. Do not reuse them. Follow manufacturer’s instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

---

## ENGINEERING CONTROL STATEMENTS

---

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use an enclosed cockpit in a manner that is consistent with the WPS for Agricultural Pesticides [40 CFR 170.240(d)(6)]. Pilots must wear the PPE required on this labeling for applicators.

---

## USER SAFETY RECOMMENDATIONS

**Users should:**

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

---

## ENVIRONMENTAL HAZARDS

---

This product chemical is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring issues. Do not contaminate water when disposing of equipment washwaters. Do not contaminate water intended for irrigation or domestic purposes. Do not apply when weather conditions favor drift from target area.

**Groundwater Contamination:** Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-DB have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-DB pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Do not apply BUTYRAC® 175 Broadleaf Herbicide directly to or otherwise permit it to come in contact with cotton, okra, grapes, tomatoes, fruit trees, vegetables, flowers, or other desirable crop or ornamental plants. Do not permit spray mist to drift onto susceptible plants since very small quantities of the BUTYRAC® 175 Broadleaf Herbicide can cause severe injury during the growing or dormant periods.

---

## DIRECTIONS FOR USE

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks and
- Protective eyewear

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container in a dry secured storage area. Keep container tightly closed when not in use.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL: Non-refillable containers (1, 2.5, 30 & 55 gallons):** Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

**(Non-refillable <5 gallons):** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**(Non-refillable >5 gallons):** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable container (250 gallon & bulk):** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing process two more times.

---

## PRODUCT RESTRICTIONS

- Do not apply BUTYRAC® 175 Broadleaf Herbicide through any type of irrigation system.
- Do not use in or near a greenhouse.
- Do not feed/graze soybean forage or harvest hay for 60 days following any 2,4-DB application.

---

## APPLICATION PROCEDURES

BUTYRAC® 175 can be applied to registered use areas by ground and aerial application equipment. The following provides recommended methods of application for each crop.

---

## SPRAY DRIFT MANAGEMENT

Do not make applications into temperature inversions.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2–10 mph at the application site.

## GROUND APPLICATION

Use a standard herbicide boom sprayer that provides uniform and accurate application. Sprayer should be equipped with screens no finer than 50 mesh in the nozzle tips and in-line strainers.

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Select a spray volume and delivery system that will ensure thorough and uniform spray coverage. For optimum spray distribution and thorough coverage, use of flat fan nozzles (maximum tip size 8008) with a minimum spray pressure of 30 psi at the nozzle tips is recommended. Other nozzle types that produce coarse spray droplets may not provide adequate coverage of the weeds to ensure optimum control. Raindrop nozzles are not recommended as weed control with BUTYRAC® may be reduced. In general, a minimum spray volume of 10 gallons per acre (GPA) is recommended for optimum spray coverage. When using higher speed equipment, a maximum speed of 10 mph is suggested if field conditions cause excessive boom movement during application and subsequent poor spray coverage. Ground applications made when dry dusty field conditions exist may provide reduced weed control in wheel track areas.

## AERIAL APPLICATION

Use orifice discs, cores and nozzle types and arrangements that will provide for optimum spray distribution and maximum coverage. In general, a minimum spray volume of 5 GPA and a maximum pressure of 40 psi are recommended.

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

---

## MIXING INSTRUCTIONS

---

**BUTYRAC® 175 ALONE:** Fill the spray tank 1/2 to 3/4 full with clean water. Begin agitation and add the recommended amount of BUTYRAC® 175. Add water to the spray tank to the desired level. Maintain sufficient agitation to ensure a uniform spray mixture during application.

**TANK MIXTURES:** BUTYRAC® 175 can be applied in tank mixture with other herbicides registered for use on approved crops. Refer to the specific crop section for rate recommendations and other restrictions. To apply BUTYRAC® 175 in mixture with another product, fill the spray tank 1/2 to 3/4 full with clean water and begin agitation. If tank mixing with wettable powder, soluble powder, flowable or dry flowable products, add the powder or flowable product first. After the other herbicide is thoroughly mixed with water, add the specified amount of BUTYRAC® 175 and add water to the spray tank to the desired level. If tank mixing with other product types, add the BUTYRAC® 175 first before adding the other product. Always mix one product in water thoroughly before adding another product or compatibility problems may occur. Never mix two products together without first mixing in water.

Maintain sufficient agitation while mixing and during application to ensure a uniform spray mixture. If spray mixture is allowed to remain without agitation for short periods of time, be sure to agitate until uniformly mixed before application.

This product can be mixed with Gramoxone® Extra, GlyStar® Original or Roundup®, Honcho®, Lorox®, Prowl®, Pursuit® Plus, Scepter®, Scepter® 70DG, Sencor®, and Squadron® for use on soybeans in accordance with the most restrictive of label limitations and precautions. Do not exceed label dosage rates. This product cannot be mixed with any product containing a label prohibition against such mixing. Refer to the crop specific information section of this label for additional information.

---

## COMPATIBILITY

---

Evaluate tank mixtures not listed on this label for compatibility and crop safety on a small area before applying to the entire field.

BUTYRAC® 175 Broadleaf Herbicide may form an insoluble precipitate in very hard water. If you expect to mix BUTYRAC® 175 with very hard water, test compatibility by mixing a small amount of BUTYRAC® 175 in the proposed dilution ratios, shake and observe. A compatibility agent approved for use on growing crops such as UNITE or E-Z MIX may be tested to reduce precipitation. Whenever hard water is used to dilute BUTYRAC® 175, spray immediately and do not allow spray mixture to sit overnight.

## PRODUCT INFORMATION

Spray tank residues of 2,4-D or MCPA mixed with BUTYRAC® 175 Broadleaf Herbicide can cause serious crop or ornamental plant injury. A sprayer previously used to apply these chemicals must be thoroughly cleaned with alkali and water before applying BUTYRAC® 175. Be sure sprayer is clean before applying BUTYRAC® 175 Broadleaf Herbicide.

Local conditions may affect the use of herbicides. Consult your State Agricultural Experiment Station, Farm Advisors, or Extension Weed Specialists for advice in selecting treatment from this label to best fit local conditions.

**INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, DO NOT USE THIS PRODUCT.**

Crop varieties vary in response to 2,4-DB and some are easily injured. Apply BUTYRAC® 175 only to varieties known to be tolerant to 2,4-DB. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to 2,4-DB, consult your seed company, State Agricultural Extension Service or qualified crop consultant for advice.

Be sure that use of this product conforms to all applicable laws, rules, and regulations. Certain states have restrictions pertaining to application distances from susceptible crops. The applicator must become familiar with these laws, rules, or regulations and follow them exactly.

## WEED LIST

**BUTYRAC® 175 Broadleaf Herbicide** will control or suppress the following weeds depending on weed size.

COMMON NAME	BOTANICAL NAME	MAXIMUM SIZE CONTROLLED*
pigweed (carelessweed)	<i>Amaranthus</i> spp.	3 inches
morningglory	<i>Ipomoea</i> spp.	36 inches
cocklebur	<i>Xanthium</i> spp.	36 inches
velvetleaf	<i>Abutilon theophrasti</i>	1 inch
Virginia copperleaf	<i>Acalypha virginica</i>	1 inch
prickly sida (teaweed)	<i>Sida spinosa</i>	1 inch
common ragweed	<i>Ambrosia artemisiifolia</i>	1 inch
jimsonweed	<i>Datura stramonium</i>	1.5 inches
devilsclaw	<i>Proboscidea louisianica</i>	2 inches
thersal croton	<i>Croton lindheimeri</i>	1 inch
lambquarters	<i>Chenopodium album</i>	1 inch
wild mustard	<i>Sinapis arvensis</i>	1 inch
field pennycress	<i>Thlaspi arvense</i>	1 inch

\*Growth of larger weeds will be only suppressed.

# SOYBEANS

## BUTYRAC® 175 Broadleaf Herbicide Directions

PRODUCT	RATE	APPLICATION TIMING AND SPECIFIC COMMENTS	
		CROP	WEEDS
Preplant through pre-emergence  BUTYRAC® 175 + Non-ionic surfactant	0.8–1.0 pint/A + 0.5% V/V	Apply to soybeans before planting or before crop emergence.	For control of emerged cocklebur, annual morningglories and other susceptible broadleaf weeds, apply when weeds are small and actively growing (see “WEED LIST”). BUTYRAC® 175 may not give complete control of larger overwintered mustards. Best results will be achieved by adding a non-ionic surfactant at a 0.5% V/V to the spray tank when making a preplant or preemergence application of BUTYRAC®175.
Postemergence broadcast (over the top)  BUTYRAC® 175	0.8–1.0 pint/A	Apply to soybeans grown in the Southern states only from 7 to 10 days before bloom up to mid-bloom when soybeans are about knee-high and growing actively. Soybean foliage should be dark green indicating that nodulation and nitrification are under way. Postemergence broadcast application at these rates prior to or after this application timing is not recommended as reduced flowering and yield may result. DO NOT APPLY BUTYRAC® 175 post-emergence broadcast to soybeans grown in the midwest states of IA, IL, IN, KS, KY (except the Purchase area), MO (except the MO bootheel), MI, MN, NE, ND, OH, SD and WI.	For control of emerged cocklebur, annual morningglories and other susceptible broadleaf weeds, apply when weeds are small and actively growing (see “WEED LIST”).
Postemergence directed band  BUTYRAC® 175	0.8–1.8 pints/A	Apply when soybeans are 8 or more inches tall with sprayer nozzles mounted to ensure proper placement of spray on only the lower 1/3 of the soybean plants. Do not allow spray to contact growing terminals of beans as excessive crop injury will result.  Do not mount nozzles on booms with drop pipes or on cultivators without gauge wheels. Use flat fan type nozzles, 8001 or larger or the equivalent with nozzle pressure less than 30 psi and at least 10 gallons of spray volume per acre.	To control emerged cockleburs and annual morningglories up to 3 inches tall, apply 0.8 to 1.0 pint/A per broadcast acre as a directed band treatment. To control other susceptible broadleaf weeds up to 1 inch tall, apply 1.6 to 1.8 pints per broadcast acre as a directed band treatment. Apply no more than 2 higher rate applications per season to reduce the potential for crop stunting.

## TANK MIXTURES OF BUTYRAC® 175 BROADLEAF HERBICIDE AND OTHER HERBICIDES IN SOYBEANS

Applying tank mixtures of BUTYRAC® 175 Broadleaf Herbicide preplant/preemergence or postemergence with other soybean herbicides as directed below can reduce competition from early weed populations and can improve weed control or control mid-to-late-season weed populations, thus minimizing the likelihood of yield reduction in soybean fields with heavy broadleaf weed infestations. However, treating soybeans under stress (as from drought or disease such as Phytophthora root rot) or in any other manner not directed on this label, can cause crop injury and yield reduction. The advantage of treating weeds with an appropriate tank mixture should be balanced against any potential for loss. Base your decision on an evaluation of each factor. Follow the local recommendations of your State Cooperative Extension Service or other agricultural weed control authority. Always follow the recommendations of any product used in tank mixture with BUTYRAC® 175.

PRODUCT	RATE	APPLICATION TIMING AND SPECIFIC COMMENTS	
		CROP	WEEDS
BUTYRAC® 175 + Gramoxone® Extra + Non-ionic surfactant	0.6–0.8 pint/A + 1.5–5 pints/A + 0.5% V/V	Apply before planting or before soybean emergence. Do not apply this tank mixture after crop emergence.	For improved control of emerged annual morningglories (ivyleaf, tall), common cocklebur, marehail and other problem weeds, apply to small actively growing weeds. Refer to the Gramoxone® Extra label for full list of weed species controlled and specific application stage and rate recommendations.
BUTYRAC® 175 + Prowl® + Non-ionic surfactant	0.8–1.0 pint/A + 1.0–3.0 pints/A + 0.5% V/V	Apply up to 45 days prior to soybean planting. Do not apply a BUTYRAC® 175 + Prowl® tank mixture at or after planting North of Interstate 80. Surface applications of Prowl® tank mixtures North of Interstate 80 require at least 1 inch rainfall or mechanical incorporation prior to planting or crop injury may result. Do not apply this tank mixture after crop emergence.	For control of emerged cocklebur, annual morningglories and other susceptible broadleaf weeds, apply when broadleaf weeds are actively growing and small (see “WEED LIST”). BUTYRAC® 175 mixtures may not give complete control of larger overwintered mustards. Best results will be achieved by adding a non-ionic surfactant to the spray tank when making a preplant application.
BUTYRAC® 175 + GlyStar® Original, Roundup® or Honcho® + Non-ionic surfactant	0.6–0.8 pint/A + 1–1.5 pints/A + 0.5–1.0% V/V	Apply before planting or before soybean emergence. Do not apply this tank mixture after crop emergence.	For improved control of emerged annual morningglories (ivyleaf, tall), common cocklebur and other problem weeds, apply to small actively growing weeds. Refer to the GlyStar® Original, Roundup® or Honcho® labels for specific rate recommendations, application stage and weed species controlled.
BUTYRAC® 175 + Pursuit® Plus + Non-ionic surfactant	0.6–0.8 pint/A + 2.5 pints/A + 0.25% V/V	Apply before planting or before soybean emergence. Do not apply this tank mixture after crop emergence.	For improved control of emerged mustards, field pennycress and other problem weeds, apply when broadleaf weeds are actively growing and small (see “WEED LIST”).
BUTYRAC® 175 + Scepter® –OR– Scepter® 70DG + Non-ionic surfactant	0.6–0.8 pint/A + 0.66 pint/A –OR– 2.8 oz./A + 0.25% V/V	Apply before planting or before soybean emergence. Do not apply this tank mixture after crop emergence.	For improved control of emerged mustards, field pennycress and other problem weeds, apply when broadleaf weeds are actively growing and small (see “WEED LIST”).

(continued)



**TANK MIXTURES OF BUTYRAC® 175 BROADLEAF HERBICIDE  
AND OTHER HERBICIDES IN SOYBEANS (cont.)**

PRODUCT	RATE	APPLICATION TIMING AND SPECIFIC COMMENTS	
		CROP	WEEDS
BUTYRAC® 175 + Squadron® + Non-ionic surfactant	0.6–0.8 pint/A + 3 pints/A + 0.25% V/V	Apply before planting or before soybean emergence. Do not apply this tank mixture after crop emergence.	For improved control of emerged mustards, field pennycress and other problem weeds, apply when broadleaf weeds are actively growing and small (see “WEED LIST”).
BUTYRAC® 175 + Lorox®	0.9 pint/A + 1 pint/A	Apply to soybeans as a directed band treatment only when soybeans are at least 8 inches high. Do not spray higher than 3 inches on the soybean stem or unacceptable crop injury may result. Do not spray over the top of soybean plants.	For postemergence control of annual morningglories, sicklepod, teaweed, cocklebur and sesbania, apply when weeds do not exceed 4 inches in height. A second application may be necessary, but do not make more than 2 applications per season.
BUTYRAC® 175 + Sencor® DF	1.0 pint/A + 0.33–0.66 lb./A	Apply to soybeans as a directed band treatment only when soybeans are at least 8 inches high, with spray or nozzles mounted to ensure proper placement of spray on no more than the lower 1/3 of the soybean plants. Do not apply directly to soybean plants or serious crop injury will occur. Soybean leaves contacted by spray will be killed. Follow all variety restrictions on the full Sencor® DF label.	For improved control of cocklebur, annual morningglories and other broadleaf weeds, apply before weeds are 3 inches tall. A non-ionic surfactant may be added to improve broadleaf weed control.

**RESTRICTIONS AND LIMITATIONS FOR USE ON SOYBEANS:**

- Beans stressed by drought or other influences should not be sprayed.
- Do not use this product on soybeans that show symptoms of disease such as Phytophthora root rot.
- Do not graze or feed soybean hay within 60 days after application of BUTYRAC® 175 tank-mix application.
- Do not harvest beans within 60 days after spray application.
- Do not treat soybeans with a tank mixture of BUTYRAC® 175 Broadleaf Herbicide and SEVIN® Carbaryl Insecticide as severe injury may result.
- When preplant through preemergence treatment is followed with a BUTYRAC® 175 postemergence application, the cumulative rate should not exceed 1.8 pints per acre per season.
- Follow all restrictions and limitations of any product used in tank mixture with BUTYRAC® 175.
- Do not use BUTYRAC® 175 alone or in tank mixture as a preplant through preemergence application to soybeans in California.

# PEANUTS

## BUTYRAC® 175 Broadleaf Herbicide Directions

PRODUCT	RATE	APPLICATION TIMING AND SPECIFIC COMMENTS	
		CROP	WEEDS
BUTYRAC® 175	0.9–1.1 pints/A	Apply to peanuts 2 to 12 weeks after planting in the states of AL, AR, FL, GA, LA, NC, SC, TN and VA. In NC, SC and VA, do not apply later than 45 days before harvest.	For control of annual morningglories, cocklebur and other broadleaf weeds, apply when weeds are small and actively growing (see “WEED LIST”). For late germinating cocklebur and morningglory, apply a second application about 3 weeks after the first application.
	0.9–1.8 pints/A	Apply to peanuts 2 to 12 weeks after planting in the states of OK, TX and NM only. BUTYRAC® 175 may be applied twice during the season, but the second application should not be made later than 100 days after planting or 30 days before harvest.	Use the low rate on morningglory and cocklebur up to 12 inches in size. Use the higher rate on other weeds. For best suppression of prickly sida, space the treatments 14 days apart.

### RESTRICTIONS AND LIMITATIONS FOR USE ON PEANUTS

- Do not feed treated peanut vines or peanut hay to livestock.
- Do not apply BUTYRAC® 175 Broadleaf Herbicide if peanut plants are under stress from drought as injury may occur.
- Do not apply to peanuts within 60 days of harvest.

---

## CONDITIONS OF SALE AND WARRANTY

---

The DIRECTIONS FOR USE of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of ALBAUGH, LLC, its Supplemental Distributors, or the Seller. All such risks shall be assumed by the Buyer.

ALBAUGH, LLC, its Supplemental Distributors and the Seller warrant that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NEITHER ALBAUGH, LLC NOR ITS SUPPLEMENTAL DISTRIBUTORS MAKE ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS FOR ANY PARTICULAR PURPOSE OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND THE EXCLUSIVE LIABILITY OF ALBAUGH, LLC, ITS SUPPLEMENTAL DISTRIBUTORS AND THE SELLER FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED.** When Buyer suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), Buyer must promptly notify Seller in writing of any claims to be eligible to receive either remedy stated above. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL ALBAUGH, LLC, ITS SUPPLEMENTAL DISTRIBUTORS, OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.** ALBAUGH, LLC, its Supplemental Distributors, and the Seller offer this product, and the Buyer accepts it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of ALBAUGH, LLC.

No employee or agent of ALBAUGH, LLC, its Supplemental Distributor, or the Seller is authorized to vary or exceed the terms of this Warranty in any other manner.

Butyrac® and GlyStar® are registered trademarks of Albaugh, LLC.

Sencor® and Sevin® are registered trademarks of Bayer CropScience.

Prowl®, Pursuit®, Scepter®, and Squadron® are registered trademarks of BASF Ag Products.

Lorox® is a registered trademark of Griffin, LLC.

Roundup® and Honcho® are registered trademarks of the Monsanto Company.

Gramoxone® is a registered trademark of Syngenta Crop Protection, Inc.