

RESTRICTED USE PESTICIDE

DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

PESTICIDA DE USO RESTRINGIDO

Tóxico para peces y organismos acuáticos. Para la venta al por menor y uso por aplicadores certificados o personas que estén bajo su directa supervisión y únicamente para los usos detallados en la certificación del aplicador certificado.

LAMBDA-CYHALOTHRIN GROUP 3 INSECTICIDE

SPECIMEN LABEL

CRUSADER[®] 1EC

INSECTICIDE

ACTIVE INGREDIENT:

Lambda-cyhalothrin..... 12.7%

OTHER INGREDIENTS:..... 87.3%

TOTAL:..... 100.0%

Contains 1 pound of active ingredient per gallon

Contains petroleum distillate

**KEEP OUT OF REACH OF CHILDREN /
MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS
WARNING / AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you **DO NOT** understand this label, find someone to explain it to you)

See inside booklet for complete First Aid, Precautionary Statements, Directions For Use, Storage and Disposal, and Conditions of Sale and Warranty.

Manufactured for:
ALBAUGH, LLC
1525 NE 36th Street,
Ankeny, IA 50021

EPA Reg. No. 42750-373
AD120220S

 **ALBAUGH[®]**
your alternative

FIRST AID / PRIMEROS AUXILIOS	
IF SWALLOWED/ SI SE INGIERE:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • DO NOT induce vomiting unless told by a poison control center or doctor. • DO NOT give any liquid to the person. • DO NOT give anything by mouth to an unconscious person. <hr/> <ul style="list-style-type: none"> • Llame de inmediato a un centro de control de envenenamientos o a un médico para consejo de tratamiento. • No induzca el vómito a menos que así se lo indique un centro de control de envenenamientos o un médico. • No le dé ningún líquido a la persona. • No administre nada por boca a una persona que haya perdido el conocimiento.
IF ON SKIN OR CLOTHING / SI CAE EN LA PIEL:	<ul style="list-style-type: none"> • Takeoff contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes • Call a poison control center or doctor for treatment advice. <hr/> <ul style="list-style-type: none"> • Quítese la ropa contaminada. • Enjuague la piel inmediatamente con bastante agua por 15-20 minutos. • Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento.
IF INHALED/ SI ES INHALADO :	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. • Call a poison control center or doctor for further treatment advice. <hr/> <ul style="list-style-type: none"> • Traslade a la persona al aire fresco. • Si la persona no está respirando llame al 911 o a una ambulancia, luego dé respiración artificial, preferiblemente de boca a boca, si es posible. • Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento.
IF IN EYES / SI ENTRA EN CONTACTO CON LOS OJOS:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. <hr/> <ul style="list-style-type: none"> • Mantenga los ojos abiertos y enjuáguelos lenta y cuidadosamente con agua, durante 15 a 20 minutos. • Si utiliza lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos. • Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Emergency phone numbers (800) 424-9300 CHEMTREC (transportation and spills)</p> <p>Cuando llame a un centro de control de envenenamiento, o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto. Teléfonos de emergencia (800) 424-9300 CHEMTREC (transporte y derrames)</p>	
<p>NOTE TO PHYSICIAN: Contains pretroleum distillate. Vomiting may cause aspiratiuon pneumonia.</p> <p>NOTA PARA EL MÉDICO: Contiene destilado de pretroleo. Los vómitos pueden causar neumonía por aspiración.</p>	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING/AVISO: May be fatal if swallowed. Harmful if absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Skin exposure may also result in a sensation described as a tingling, itching, burning, or prickly feeling. Onset may occur immediately to 4 hours after exposure and may last 2 to 30 hours, without damage. Wash exposed areas once with soap and water. Relief from the skin sensation may be obtained by applying an oil-based cream.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

1. Long-sleeved shirt and long pants
2. Chemical resistant gloves, made of barrier laminate or viton (> 14 mils)
3. Shoes plus socks

Mixers/loaders supporting aerial applications to Wild rice at the rate of 0.04 lb. a.i. (5.12 fl. ozs.) per acre and treating 1,200 acres (or more) per day must wear: Wear a minimum of a NIOSH-approved particulate facepiece respirator with any R or P filter, OR a NIOSH approved elastometric particulate respiratory with any R, or P filter OR a NIOSH approved powered air-purifying respirator with a HE filter.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

DECLARACIONES PREVENTIVAS RIESGOS PARA LOS SERES HUMANOS Y ANIMALES DOMÉSTICOS

AVISO: Puede ser letal si se ingiere. Dañino si se absorbe por la piel. Dañino si se inhala. Causa irritación moderada en los ojos. Evite el contacto con la piel, los ojos o ropa. Evite respirar rociado o aerosol. El contacto prolongado y frecuente con la piel puede causar reacciones alérgicas en algunos individuos.

La exposición de la piel también puede provocar una sensación descrita como hormigueo, picazón, ardor o picazón. El inicio puede ocurrir inmediatamente hasta 4 horas después de la exposición y puede durar de 2 a 30 horas, sin daño. Lave las áreas expuestas una vez con agua y jabón. Se puede obtener alivio de la sensación de la piel aplicando una crema a base de aceite.

EQUIPO DE PROTECCIÓN PERSONAL (PPE)

Los aplicadores y otros manipuladores de pesticidas (plaguicidas) deberán usar:

1. Camisa de manga larga, pantalones largos
2. Guantes resistentes a productos químicos hechos de material resistente al agua- La Categoría A (p.ej. laminados impermeables, goma de butilo, goma de nitrilo, goma de neopreno, polietileno, cloruro de polivinilo (PVC, por sus siglas en inglés), o Viton®)
3. Zapatos y calcetines

Mezcladores/cargadores que soportan aplicaciones aéreas al arroz silvestre a razón de 0,04 lb. ai. (5,12 onzas líquidas) por acre y para tratar 1200 acres (o más) por día se debe usar: Usar como mínimo un respirador de pieza facial para partículas aprobado por NIOSH con cualquier filtro R o P, O un respirador elastométrico para partículas aprobado por NIOSH con cualquier filtro R, o filtro P O un respirador purificador de aire motorizado aprobado por NIOSH con filtro HE.

Deseche la ropa y otros materiales absorbentes que hayan sido empapados o muy contaminados con el concentrado de este producto. NO los reutilice. Siga las instrucciones del fabricante para limpiar y mantener el EPP. Si no existen instrucciones para los productos lavables, utilice detergente y agua caliente. Mantenga y lave el EPP por separado del resto de la ropa.

Cuando los manipuladores utilizan sistemas cerrados, cabinas cerradas o aeronaves de una manera que cumpla con los requisitos enumerados en la Norma de Protección al Trabajador (WPS) para pesticidas agrícolas [40 CFR 170.240(d)(4-6)], los requisitos del PPE del manipulador pueden ser reducido o modificado según lo especificado en la WPS.

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

RECOMENDACIONES DE SEGURIDAD PARA EL USUARIO

1. Lávese las manos antes de comer, beber, masticar goma o tabaco, o ir al baño. Quitarse la ropa/EPP inmediatamente si el pesticida entra dentro de ellos. Luego lávese abundantemente y póngase ropa limpia.
2. Quítese la ropa/EPP inmediatamente si el pesticida entra. Luego lávese bien y póngase ropa limpia.
3. Quítese el EPP inmediatamente después de manipular este producto. Lave el exterior de los guantes antes de quitárselos. Tan pronto como sea posible, lávese bien y cámbiese a ropa limpia.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT mix or allow to come into contact with oxidizing agent. Hazardous Chemical reaction may occur.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic organisms and toxic to wildlife. **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high-water mark. **DO NOT** apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when disposing of equipment wash water.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. **DO NOT** apply this product or allow it to drift to blooming crops or weeds if bees are foraging in the treatment area.

DIRECTIONS FOR USE

Restricted Use Pesticide

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

DIRECCIONES DE USO

PESTICIDA DE USO RESTRINGIDO

El uso de este producto de forma contraria a lo indicado en su etiqueta constituye una infracción de la ley federal.

No aplique este producto de manera que entre en contacto con trabajadores u otras personas, ya sea directamente o por deriva. Solo los manipuladores protegidos pueden estar en el área durante la aplicación. Para cualquier requisito específico de su estado o tribu, consulte a la agencia estatal o tribal responsable de la regulación de pesticidas.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

1. Coveralls
2. Chemical-resistant gloves (made of any waterproof material), made of barrier laminate or viton (> 14 mils)
3. Shoes plus socks

REQUISITOS DE USO AGRICOLA

Utilice este producto solo de acuerdo con su etiqueta y el Estándar para la Protección del Trabajador Agrícola (WPS, por sus siglas en inglés), 40 CFR Parte 170. Esta ley contiene los requisitos para la protección de los trabajadores agrícolas en granjas, bosques, viveros e invernaderos, y para las personas que manipulan pesticidas agrícolas. Contiene requisitos para la capacitación, descontaminación, notificación y asistencia de emergencia. También contiene instrucciones específicas y excepciones relacionadas con las indicaciones en esta etiqueta acerca del equipo de protección personal (PPE, por sus siglas en inglés), y el intervalo de ingreso restringido. Los requisitos en esta sección de la etiqueta (requisitos para uso agrícola) aplican únicamente a los usos de este producto que están cubiertos por el Estándar para la Protección del Trabajador Agrícola.

No ingrese o permita el ingreso de trabajadores a las áreas tratadas durante el intervalo de acceso restringido (REI, por sus siglas en inglés) de 24 horas.

El equipo de protección personal requerido para el acceso anticipado a áreas tratadas de acuerdo con el Estándar para la Protección del Trabajador Agrícola, y que involucra el contacto con material tratado, como plantas, tierra o agua, es:

1. Overol (mameluco)
2. Guantes resistentes a productos químicos
3. Zapatos y calcetines

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standards for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries and greenhouses.

Do not allow people or pets on treated surfaces until the spray has dried.

REQUISITOS DE USO NO AGRÍCOLA

Los requisitos en este recuadro se aplican a los usos de este producto que NO están dentro del alcance de la Norma de Protección del Trabajador para pesticidas agrícolas (40 CFR Parte 170). El WPS se aplica cuando este producto se usa para producir plantas agrícolas en granjas, bosques, viveros o invernaderos.

No permita que personas o mascotas entren en las superficies tratadas hasta que el aerosol se haya secado.

PRODUCT INFORMATION

Apply in sufficient water for thorough coverage of listed crops unless otherwise specifically noted. Base rate of application upon pest pressure, timing of sprays, and field scouting. Use higher rates under heavy pest pressure and lower rates under low to moderate pest pressure. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds and other local methods. For ground and air applications, unless otherwise noted, use the following spray volumes:

Row Crops:

By ground, apply in a minimum of 10 gallons of finished spray per acre.

By air, apply in a minimum of 2 gallons of finished spray per acre.

Orchard and Vine Crops:

By ground, apply in a minimum of 50 gallons of finished spray per acre.

By air, apply in a minimum of 10 gallons of finished spray per acre.

For cutworm control, CRUSADER® 1EC may be applied before, during, or after planting. For soil incorporated applications, use listed higher rates for improved control.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, this product contains a Group 3 insecticide. Any insect population may contain individuals naturally resistant to this product and other Group 3 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed. -Rotate the use of this product or other Group 3 insecticides within a growing season, or among growing seasons, with different groups that control the same pests. Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. **DO NOT** rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):

Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.

Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.

When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).

Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.

The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active. Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices. -Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area. For further information or to report suspected resistance contact Albaugh, LLC at 1-800-247-8013.

SPRAY DRIFT PRECAUTIONS

BUFFER ZONES

Vegetative Buffer Strip

Construct and maintain a minimum 10-foot-wide vegetative filter strip of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as but not limited to lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing lambda-cyhalothrin onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.

For guidance, refer to the following publication for information on constructing and maintaining effective buffers: *Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services. USDA, NRCS.2000. Fort Worth, Texas. 21 pp.* https://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs143_023819.pdf

Buffer Zone for Ground Application (ground boom, overhead chemigation, or airblast)

DO NOT apply within 25 feet of aquatic habitats (such as but not limited to lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

Buffer Zone for ULV Aerial Application

DO NOT apply within 450 feet of aquatic habitats (such as but not limited to lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

Buffer Zone for Non-ULV Aerial Application

DO NOT apply within 150 feet of aquatic habitats (such as but not limited to lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

In the state of New York, a 25-foot vegetated, non-cropped buffer strip untraversed by drainage tiles must be maintained between a treated field and a coastal salt marsh or stream that drains into a coastal salt marsh, for both aerial or ground application. For aerial applications, the 25-foot vegetated non-cropped buffer strip for runoff protection would be part of the larger 150-foot buffer strip (or 450-foot buffer strip for ULV application) required for spray drift.

SPRAY DRIFT REQUIREMENTS

Wind Direction and Speed

Only apply this product if the wind direction favors on-target deposition.

DO NOT apply when the wind velocity exceeds 15 mph.

Temperature Inversion

DO NOT make aerial or ground applications into temperature inversions.

Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Droplet Size

Use only medium or coarser spray nozzles (for ground and non-ULV aerial application) according to ASAE (S572) definition for standard nozzles. In conditions of low humidity and high temperatures, use a coarser droplet size.

Additional Requirements for Ground Applications

Wind speed must be measured adjacent to the application site on the upwind side, immediately prior to application.

For ground boom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy.

For airblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

Additional Requirements for Aerial Applications

Mount the spray boom on the aircraft to minimize drift caused by wingtip or rotor vortices. Use the minimum practical boom length; **DO NOT** exceed 75% of the wing span or 80% rotor diameter.

Flight speed and nozzle orientation must be considered in determining droplet size.

Spray must be released at the lowest height consistent with pest control and flight safety. **DO NOT** release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

TANK MIX APPLICATION

Fill the spray tank at least one-third full of clean water or diluent. With the pump and agitator running continuously, add the specified amount of each product in the tank mix to the spray tank and allow to fully disperse, adding this product last. Add the remainder of water or diluent to the spray tank. Follow the precautions and limitations of the most restricted product in the tank mixture.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Compatibility testing for tank mixing partners: Test compatibility of the intended tank mixture by adding proportionate amounts of each ingredient to a pint or quart jar, cap, shake, and let set 15 minutes. Formation of precipitates that **DO NOT** readily redisperse indicates an incompatible mixture that must not be used.

DO NOT use non-emulsifiable oils in combination with this product. If adjuvants are used, use only:

- Nonionic Surfactant (NIS) containing at least 75% surface agent, or
- Non-phytotoxic Crop Oil Concentrate (COC) including once refined Vegetable Oil concentrate (VOC), or
- Methylated Seed Oils (MSO) containing a minimum of 17% emulsifier.

Adjuvants other than NIS or COC may be used providing the product;

1. Contains only EPA exempt ingredients.
2. Is non-phytotoxic to the target crop.
3. Is compatible in mixture (may be established through a jar test).
4. Is supported locally for use with this product on the target crop through proven field trials and through university and extension specifications.

The following may be used as diluents:

Crop Oil Concentrate
Methylated Seed Oils
Urea-Ammonium Nitrate

DO NOT use the following in combination with this product as diluents or adjuvants:

Non-emulsifiable Oils
Diesel Fuel
Straight Mineral Oil

When an adjuvant is to be used with this product, Albaugh suggests the use of a Chemical Producers and Distributors Association certified adjuvant.

CHEMIGATION

Sprinkler Irrigation Application

Apply this product at rates and timing described elsewhere in this label. Consult your local State Extension Service or other local experts for specifications pertinent for your area.

Thorough, uniform coverage of foliage is required for good control. Maintain good agitation in the pesticide supply tank prior to and during the entire application period.

Apply by injecting the specified rate of this product into the irrigation system using a metering device that will introduce a constant flow and by distributing the product to the target area in 0.1-0.2 acre-inch of water. Use the least amount of water required for proper distribution and coverage. Inject the product into the main irrigation line ahead of a right angle turn in the line to insure adequate dispersion or mixing in the irrigation water. Once the application is completed, flush the entire irrigation and injection system with clean water before stopping the system.

Additionally, if application is being made during a normal irrigation set of a stationary sprinkler, inject the specified rate of this product for the area covered into the system only during the end of the irrigation set for sufficient time to provide adequate coverage and product distribution.

DO NOT apply this product through an irrigation system connected to a public water system. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves and average of at least 25 individuals daily at least 60 days out of the year.

Use Precautions: Sprinkler Irrigation Application

Apply this product only through sprinkler irrigation systems (including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move). **DO NOT** apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, contact state extension service specialist, equipment manufacturers, or other experts.

DO NOT connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and a low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

DO NOT apply when wind speed favors drift beyond the area intended for treatment.

**CROP USE DIRECTIONS
AGRICULTURAL USES**

ALFALFA AND ALFALFA GROWN FOR SEED			
TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Alfalfa Caterpillar Army cutworm Cutworm spp. Green Cloverworm Leafhopper spp. Looper spp. Threecornered Alfalfa Hopper Velvetbean Caterpillar Webworm spp.	0.015-0.025	1.92-3.20	<p>Ground application: Apply in a minimum of 10 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Make applications when pests appear. Apply in sufficient volume to ensure sufficient coverage of foliage.</p>
Alfalfa Seed Chalcid (Adult) Alfalfa Weevil Armyworm Bean Leaf Beetle (Adult) Blister Beetle spp. Blue Alfalfa Aphid Clover Leaf Weevil spp. Clover Root Borer (Adult) Clover Root Curculio spp. (Adult) Clover Stem Borer (Adult) Corn Earworm Cowpea Aphid Cowpea Curculio (Adult) Cowpea Weevil (Adult) Cucumber Beetle spp. (Adult) Egyptian Alfalfa Weevil Fall Armyworm ¹ Grape Colaspis (Adult) Grasshopper spp. Green June Beetle (Adult) Green Peach Aphid ³ Japanese Beetle (Adult) Meadow Spittlebug Mexican Bean Beetle Pea Aphid Pea Weevil (Adult) Plant Bug spp., including Lygus spp. ³ Spotted Alfalfa Aphid Stink Bug spp. Sweet Clover Weevil (Adult) Thrips spp. ⁴ Western Yellow-striped Armyworm Whitefringed Beetle spp. (Adult) Yellow-striped Armyworm	0.02-0.03	2.56-3.84	<p>When foliage is dense and/or pest populations are high, use 5-10 gals/A by air or 20 gals/A by ground and higher use rates. Use higher rates for increased residual control.</p> <p>DO NOT apply when bees are foraging. Apply during the early morning or evening hours. Be aware of bee hazard resulting from a cool evening and/or morning dew.</p> <p>Remove bee shelters during and for 2-3 days following application. DO NOT apply directly to bee shelters.</p> <p>Apply only to fields planted to pure stands of alfalfa.</p> <p>Apply as required by scouting.</p> <p>1 For control of first and second instars only. 2 Suppression only. 3 See resistance statement under PRODUCT INFORMATION. 4 Does not include Western Flower Thrips</p>
Beet Armyworm ^{1,3} Blotch Leafminer ³ Spider Mites ²	0.03	3.84	
Restrictions:			
<ul style="list-style-type: none"> • DO NOT apply more than 0.03 lb a.i. (0.24 pt) per acre per cutting. • DO NOT apply more than 0.12 lb a.i. (0.96 pt) per acre per season. • DO NOT apply within 1 day of harvest for forage or within 7 days of harvest for hay. 			

CANOLA			
TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Armyworm spp. Cabbage Seedpod Weevil Cutworm spp. Diamondback Moth Flea Beetle Grasshoppers Looper spp. Lygus Bug	0.015-0.03	1.92-3.84	Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area. Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area. Make applications when pests appear. If needed, make repeat applications after at least 5 or more days. Apply in sufficient volume to ensure sufficient coverage of foliage.
Cabbage Aphid	0.03	3.84	
Restrictions:			
<ul style="list-style-type: none"> • DO NOT apply within 7 days of harvest. • DO NOT apply more than 0.09 lb a.i. (0.72 pt) per acre per year. 			

CEREAL GRAINS:						
Corn (At-Plant): Field Corn, Popcorn, Seed Corn, Sweet Corn						
TARGET PESTS	RATE			REMARKS		
	lb a.i./A	fl oz/A				
Corn Rootworm Larvae (Western, Northern, Southern, Mexican) Cutworm spp. Seedcorn Maggot Seedcorn Beetle Lesser Cornstalk Borer White Grub spp. Wireworm spp. Red Imported Fire Ant ¹	0.005 lb a.i. per 1000 ft of row ²	0.66 fl oz per 1000 ft of row ²		Banded Applications: Apply at planting as a 5-7 inch T-band sprayed across the open seed furrow between the furrow openers and the press wheels or as a band application behind the press wheel. In-Furrow Applications: Apply into the seed furrow through spray nozzles or microtubes behind the planter furrow openers and in front of the press wheel. Apply a minimum of 3 gals of finished spray/A. ¹ Suppression only.		
² lbs ai and fl oz/A of CRUSADER® 1EC applied at 0.66 fl oz/1000 ft of row for various row spacings:						
Row Spacing	40"	38"	36"	34"	32"	30"
Linear Ft per acre	13,068	13,756	14,520	15,374	16,335	17,424
Lbs a.i. per acre	0.067	0.07	0.075	0.079	0.084	0.09
Fl oz per acre	8.6	9.1	9.6	10.1	10.8	11.5
Restrictions:						
<ul style="list-style-type: none"> • DO NOT harvest or graze livestock or cut treated crops for feed within 21 days of at-plant application. • DO NOT apply more than 0.09 lb a.i. (0.72 pt) per acre per crop at-plant. • DO NOT apply more than 0.12 lb a.i. per acre per crop from at-plant and foliar applications for field corn, popcorn, and seed corn. For sweet corn, DO NOT apply more than 0.48 lb a.i. per acre per crop from at-plant and foliar applications. 						

CEREAL GRAINS:			
Corn (Foliar): Field Corn, Popcorn, Seed Corn			
TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Corn Earworm ¹ Cutworm spp. Green Cloverworm Meadow Spittlebug Western Bean Cutworm ¹	0.015-0.025	1.92-3.20	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p>
Armyworm ² Bean Leaf Beetle Bird Cherry-Oat Aphid ³ Cereal Leaf Beetle Corn Leaf Aphid ³ English Grain Aphid ³ European Corn Borer ¹ Fall Armyworm ² Flea Beetle spp. Grasshopper spp. Hop Vine Borer ¹ Japanese Beetle (Adult) Lesser Cornstalk Borer Mexican Corn Rootworm Beetle (Adult) Northern Corn Rootworm Beetle (Adult) Sap Beetle (Adult) Seedcorn Beetle Southern Corn Rootworm Beetle (Adult) Southwestern Corn Borer ¹ Stalk Borer ¹ Stink Bug spp. Tobacco Budworm ^{1,4} Webworm spp. Western Corn Rootworm Beetle (Adult) Yellow-striped Armyworm ²	0.02-0.03	2.56-3.84	<p>Make applications when pests appear. If needed, make repeat applications after at least 7 days. Apply in sufficient volume to ensure sufficient coverage of foliage.</p> <p>Chinch bug control: Begin applications when bugs migrate from small grains or grass weeds to small corn. Direct spray to the base of corn plants. Repeat applications at 3-5 day intervals if needed. CRUSADER[®] 1EC may only suppress heavy infestations and/or subsequent migrations.</p> <p>Adult corn rootworm beetles (<i>Diabrotica</i> species): Use a minimum of 3.84 fl oz per acre (0.03 lb a.i. per acre) as part of an aerial-applied corn rootworm control program.</p> <p>¹ For control before the larva bores into the plant stalk or ear. ² For control of first and second instar only. ³ Suppression only. ⁴ See resistance statement under PRODUCT INFORMATION.</p>

(continued)

CEREAL GRAINS (cont.):			
Corn (Foliar): Field Corn, Popcorn, Seed Corn			
TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Beet Armyworm ⁴ Chinch Bug Green Bug ^{3,4} Southern Corn Leaf Beetle ³ Rice Stalk Borer ¹ Mexican Rice Borer ¹ Sugarcane Borer ¹	0.03	3.84	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Make applications when pests appear. If needed, make repeat applications after at least 7 days. Apply in sufficient volume to ensure sufficient coverage of foliage.</p> <p>Chinch bug control: Begin applications when bugs migrate from small grains or grass weeds to small corn. Direct spray to the base of corn plants. Repeat applications at 3-5 day intervals if needed. CRUSADER® 1EC may only suppress heavy infestations and/or subsequent migrations.</p> <p>Adult corn rootworm beetles (<i>Diabrotica</i> species): Use a minimum of 3.84 fl oz per acre (0.03 lb a.i. per acre) as part of an aerial-applied corn rootworm control program.</p> <p>¹ For control before the larva bores into the plant stalk or ear. ² For control of first and second instar only. ³ Suppression only. ⁴ See resistance statement under PRODUCT INFORMATION.</p>
<p>Restrictions:</p> <ul style="list-style-type: none"> • DO NOT apply within 21 days of harvest. • DO NOT allow livestock to graze in treated areas or harvest treat corn forage as feed for meat or dairy animals within 1 day after last treatment. • DO NOT feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment. • DO NOT apply more than 0.12 lb a.i. (0.96 pt) per acre per crop per season. • DO NOT apply more than 0.06 lb a.i. (0.48 pt) per acre after silk initiation. • DO NOT apply more than 0.03 lb a.i. (0.24 pt) after corn has reached the milk stage (yellow kernels with milky fluid). 			

CEREAL GRAINS:

Corn (Foliar): Sweet Corn

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Aphid spp. ^{2,3} Armyworm ¹ Aster Leafhopper Beet Armyworm ^{1,3} Chinch Bug Common Cornstalk Borer Corn Earworm Cutworm spp. European Corn Borer Fall Armyworm ¹ Flea Beetle spp. Grasshopper spp. J Japanese Beetle (Adult) Mexican Corn Rootworm Beetle (Adult) Northern Corn Rootworm Beetle (Adult) Sap Beetle (Adult) Southern Armyworm ¹ Southern Corn Rootworm Beetle (Adult) Southwestern Corn Borer Spider Mite spp. ² Stink Bug spp. Tarnished Plant Bug Webworm spp. Western Bean Cutworm Western Corn Rootworm Beetle (Adult) Yellow-Striped Armyworm ¹	0.02-0.03	2.56-3.84	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Make applications when pests appear. If needed, make repeat applications after at least 4 days and before insects enter the stalk or ear.</p> <p>Apply in sufficient volume to ensure sufficient coverage of foliage and ears (if present).</p> <p>Adult corn rootworm beetles (<i>Diabrotica</i> species): Use a minimum of 3.2 fl oz per acre (0.025 lb a.i. per acre) as part of an aerial-applied corn rootworm control program.</p> <p>¹For control of first and second instar only. ²Suppression only. ³See resistance statement under PRODUCT INFORMATION.</p>
Corn Silkfly (Adult) ²	0.03	3.84	

Restrictions:

- **DO NOT** apply within 1 day of harvest.
- **DO NOT** allow livestock to graze in treated areas or harvest treated corn forage as feed for meat or dairy animals within 1 day after last treatment.
- **DO NOT** feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment.
- **DO NOT** apply more than 0.48 lb a.i. (3.84 pts) per acre per crop from at plant and foliar applications.

CEREAL GRAINS:

Rice, Wild Rice

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl.oz./A	
Bird Cherry-Oat Aphid Chinch Bug Fall Armyworm Grasshopper spp. Greenbug Leafhopper spp. Rice Stink Bug Rice Water Weevil (Adult) Riceworm Sharpshooter spp. True Armyworm Yellow Sugarcane Aphid Yellow-striped Armyworm	0.025-0.04	3.20-5.12	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Mixers/loaders supporting aerial applications to wild rice at a rate of 0.04 a.i./A, and treating 1200 acres (or more) per day must wear dust/mist respirator. Apply in a minimum of 2 gals per acre in sufficient spray volume to obtain full coverage of the foliage or target area. Adding 1 pint per acre of an emulsifiable crop oil will help improve coverage, reduce evaporation, and improve efficacy. Monitor insect populations to determine timing and frequency of applications. Scout fields at a minimum of 5 day intervals.</p> <p>Make applications when pests appear. If needed, make repeat applications after at least 5-7 days. Apply in sufficient volume to ensure sufficient coverage of foliage. CRUSADER® 1EC can be safely used when propanil products are being used for weed control.</p> <p>Rice Water Weevil: In dry seeded rice, make a foliar application as indicated by scouting for the presence of adults and/or feeding scars usually within 0-5 days after permanent flood establishment. DO NOT exceed 10 days from starting permanent flood until insecticide application unless scouting indicates weevils have not been previously present. Adults may also be treated at later stages of rice development to reduce overwintering populations. In water seeded rice, make the first foliar application after pinpoint flood as indicated by scouting for the presence of adults and/or feeding scars usually when rice has emerged 0.5 inch above the waterline. Under conditions of prolonged migration into the field, start field scouting for rice water weevil adults and/or feeding scars 3-5 days after the initial treatment and, if needed, apply a second application within 7-10 days of the first application. Adults may also be treated at later stages of rice development to reduce overwintering populations.</p> <p>California: In addition to above directions, for control of rice water weevil in water seeded rice, CRUSADER® 1EC may be applied at the 1- to 3-leaf growth stage with the majority at the 2-leaf growth stage.</p> <p>Adults are vulnerable on levees and in the water. Larvae are vulnerable while feeding on the leaf prior to entering the soil. Monitor for adults based upon field history and density of population. Monitor field edges and levee areas for adults. Treat in the following manner: a) spray the inside perimeter of the field, or b) spray the entire field.</p> <p>Green Bug: Known to have many biotypes, CRUSADER® 1EC may only provide suppression. If satisfactory control is not achieved, a resistant biotype may be present. Use alternate chemistry for control. For control of stem borers, scout fields, when rice growth is near panicle differentiation, for early symptoms of damaging populations exhibited as discoloration (orange-tan) around the junction of the leaf sheath and leaf blade which is caused by feeding of young larvae within the sheath. Applications must be made before larvae bore into rice stems. Make the first application at panicle differentiation to 2 inch panicle for partial control. Make the second application at boot to heading for maximum control. All rice varieties are susceptible to stem borer damage, but Cocodrie and Priscilla are particularly susceptible.</p> <p>¹For control before the larvae bores into the plant stalk.</p>
European Corn Borer ¹ Mexican Rice Borer ¹ Rice Seed Midge ¹ Rice Stalk Borer ¹ Sugarcane Borer ¹	0.03-0.04	3.84-5.12	

Restrictions:

- **DO NOT** release floodwater within 7 days of an application.
- **DO NOT** apply more than 0.12 lb a.i. (0.96 pt) per acre per season.
- **DO NOT** apply more than 0.04 lb a.i. (0.32 pt) per acre within 21 to 27 days of harvest.
- **DO NOT** apply within 21 days of harvest.
- **DO NOT** use treated rice fields for the aquaculture of edible fish and crustacea.
- **DO NOT** apply as an ultra-low volume (ULV) spray.

CEREAL GRAINS:

Wet Sown Rice (CA Only)

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl.oz./A	
Rice Water Weevil	0.03-0.04	3.84-5.12	<p>Uniformly apply at 3.84-5.12 fl oz of product per acre as a pre-flood, pre-plant, broadcast soil application for control of Rice Water Weevil (<i>Lissorhoptus oryzophilus</i>) in wet- sown rice culture.</p> <p>Apply by air or ground equipment using sufficient water to obtain full coverage. Apply in a minimum of 2 gals of water (or a total carrier volume)/Acre by air or a minimum of 20 gals of water (or a total carrier volume)/Acre by ground.</p> <p>For improved efficacy, light incorporation of this product into the upper 1-2 inches of soil following application is recommended – a "roller" may be used for this incorporation. Apply pinpoint flood not more than 5 days after the soil application of this product, or weevil control may be reduced. Scout for feeding scars after plant emergence and apply a second foliar treatment if needed.</p>

Restrictions:

- Restricted Reentry Interval (REI) is 24 hours.
- **DO NOT** apply more than 0.04 lb a.i. (5.12 fl oz) per acre per season.
- **DO NOT** release floodwater within 7 days of application.
- **DO NOT** use treated rice fields for aquaculture of edible fish and crustacea.
- **DO NOT** apply as an ultra-low volume (ULV) spray.
- **DO NOT** apply by chemigation.

CEREAL GRAINS:

Sorghum (Grain)

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Cutworm spp. Sorghum Midge	0.015-0.02	1.92-2.56	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Make applications when pests appear. If needed, make repeat applications after at least 5 days. Apply in sufficient volume to ensure sufficient coverage of foliage. Sorghum Midge: Begin applications when 25% of the sorghum heads have emerged and are in tip bloom. Repeat applications at 5 day intervals if needed.</p> <p>Chinch Bug: Begin applications when bugs migrate from small grains or grass weeds to small sorghum. Direct spray to the base of sorghum plants. Repeat applications at 3 to 5 day intervals if needed. CRUSADER® 1EC may only suppress heavy infestations and/or subsequent migrations.</p> <p>¹For control of first and second instar only. ²For control before the larva bores into the plant stalk. ³See resistance statement under PRODUCT INFORMATION.</p>
Armyworm Beet Armyworm ³ Corn Earworm European Corn Borer ² Fall Armyworm ¹ Flea Beetle spp. Grasshopper spp. Lesser Cornstalk Borer ² Southwestern Corn Borer ² Stink Bug spp. Webworm spp. Yellow-striped Armyworm ¹	0.02-0.03	2.56-3.84	
Chinch Bug Mexican Rice Borer ² Rice Stalk Borer ² Sugarcane Borer ²	0.03	3.84	

Restrictions:

- **DO NOT** apply within 30 days of harvest.
- **DO NOT** apply more than 0.08 lb a.i. (0.64 pt) per acre per season.
- **DO NOT** apply more than 0.06 lb a.i. (0.48 pt) per acre per season after crop emergence.
- **DO NOT** apply more than 0.02 lb a.i. (0.16 pt) per acre per season once crop is in soft dough stage.
- **DO NOT** graze livestock in treated areas or harvest for fodder, silage, or hay.

CEREAL GRAINS:

Barley, Buckwheat, Oats, Rye, Triticale, Wheat, Wheat Hay

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Army Cutworm Cutworm spp.	0.015-0.025	1.92-3.20	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Make applications when pests appear. If needed, make repeat applications after at least 5 days. Apply in sufficient volume to ensure sufficient coverage of foliage.</p> <p>Chinch Bug: Repeat applications at 3 to 5 day intervals if needed. CRUSADER® 1EC may only suppress heavy infestations and/or migrations.</p> <p>Green Bug: Known to have many biotypes, CRUSADER® 1EC may only provide suppression. If satisfactory control is not achieved, a resistant biotype may be present. Use alternate chemistry for control.</p>
Armyworm Bird Cherry-Oat Aphid ¹ Cereal Leaf Beetle English Grain Aphid ¹ Fall Armyworm Flea Beetle spp. Grasshopper spp. Hessian fly ⁴ Orange Blossom Wheat Midge Russian Wheat Aphid ¹ Stink Bug spp. Yellow-striped Armyworm	0.02-0.03	2.56-3.84	
Grass Sawfly	0.025-0.03	3.20-3.84	
Chinch Bug Corn Leaf Aphid ² Greenbug ^{1,3} Mite spp. ²	0.03	3.84	

Restrictions:

- **DO NOT** apply within 30 days of harvest.
- **DO NOT** allow livestock to graze in treated areas or harvest treated wheat forage as feed for meat or dairy animals within 7 days after treatment.
- **DO NOT** feed treated straw to meat or dairy animals within 30 days after the last treatment.
- **DO NOT** apply more than 0.06 lb a.i. (0.48 pt) per acre per season.

COLE CROPS

Head and stem brassica crop group : Broccoli, Brussels Sprouts, Cabbage, Cavalo Broccolo, Cauliflower, Chinese Broccoli (gai lon), Chinese Cabbage (napa), Chinese Mustard Cabbage (gai choy), Kohlrabi

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Alfalfa Looper Cabbage Looper Cabbage Webworm Cutworm spp. Imported Cabbageworm Southern Cabbageworm	0.015-0.025	1.92-3.20	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Make applications when pests appear. If needed, make repeat applications after at least 5 days. Apply in sufficient volume to ensure sufficient coverage of foliage.</p>
Aphid spp. ^{2,3} Armyworm Beet Armyworm ^{1,3} Corn Earworm Diamondback Moth ³ Fall Armyworm ¹ Flea Beetle spp. Grasshopper spp. Japanese Beetle (Adult) Leafhopper spp. Meadow Spittlebug Plant Bug spp. including Lygus spp. ³ Spider Mite spp. ² Stink Bug spp. Thrips spp. ² Vegetable Weevil (Adult) Whitefly spp. ^{2,3} Yellow-striped Armyworm	0.02-0.03	2.56-3.84	<p>¹For control of first and second instar only.</p> <p>²Suppression only.</p> <p>³See resistance statement under PRODUCT INFORMATION.</p>

Restrictions:

- **DO NOT** apply within 1 day of harvest.
- **DO NOT** apply more than 0.24 lb a.i. (1.92 pts) per acre per season.

COTTON

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Cutworm spp. Soybean Thrips Tobacco Thrips	0.015-0.02	1.92-2.56	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>ULV application: CRUSADER® 1EC may be mixed with once-refined vegetable oil and applied in a minimum of at least 1 qt. of finished spray per acre.</p> <p>Make applications when pests appear. If needed, make repeat applications after at least 5 to 7 days. Apply in sufficient volume to ensure sufficient coverage of foliage.</p> <p>Under light bollworm/budworm infestation levels, 0.02 lb a.i. per acre may be applied in conjunction with intense field monitoring.</p> <p>Boll Weevil: Spray on a 3- to 5-day schedule.</p> <p>Cotton Bollworm, Tobacco Budworm: CRUSADER® 1EC also provides ovicidal control of unhatched <i>Heliothine</i> spp. eggs.</p> <p>¹For control of first and second instar only. ²Suppression only. ³See resistance statement under PRODUCT INFORMATION.</p>
Cabbage Looper Cotton Fleahopper Cotton Leafperforator Cotton Leafworm Lygus Bug spp. ³ Pink Bollworm Saltmarsh Caterpillar	0.02-0.03	2.56-3.84	
Bandedwing Whitefly ^{2,3} Beet Armyworm ^{1,3} Boll Weevil Brown Stink Bug Cotton Aphid ^{2,3} Cotton Bollworm European Corn Borer Fall Armyworm Green Stink Bug Southern Green Sting Bug Sweetpotato Whitefly ^{2,3} Tobacco Budworm ³ Two-spotted Spider Mite ²	0.025-0.04	3.20-5.12	

Restrictions:

- **DO NOT** apply within 21 days of harvest.
- **DO NOT** graze livestock in treated areas.
- **DO NOT** apply more than 1.6 pts (0.2 lb a.i.) per acre per season.
- **DO NOT** make more than a total of 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season.

Synthetic pyrethroid products include but are not limited to permethrin, esfenvalerate, cyfluthrin, bifenthrin, fenpropathrin, or other lambda-cyhalothrin products

CUCURBIT VEGETABLES CROP GROUP

Including:

Chayote (fruit), Chinese Waxgourd (Chinese preserving melon), Citron Melon, Cucumber, Gherkin, Gourd (edible) *Lagenaria* spp. – (includes: hyotan, cucuzza),
 Luffa acutangula, (includes: hechima, Chinese okra),
 Momordica spp. – (includes: balsam apple, balsam pear, bitter melon, Chinese cucumber),
 Muskmelon (hybrids and/or cultivars of *Cucumis melo*) – (includes: true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon honeydew melon, honey balls, mango melon Persian melon, pineapple melon, Santa Claus melon, snake melon),
 Pumpkin,
 Squash, summer (*Cucurbita pepo* var. *meloepo*) – includes: crookneck squash, straightneck squash, vegetable marrow, zucchini),
 Squash, winter (*Cucurbita maxima*, *C. moschata*) – includes: butternut squash, calabaza, hubbard squash (*C. mixta*; *C. pepo*) – includes: acorn squash, spaghetti squash
 Watermelon – includes: Hybrids and/or varieties of *Citrullus lanatus*

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Armyworm spp. ¹ Blister Beetle spp. Cabbage Looper Corn Earworm Cricket spp. Cucumber Beetle spp. (adults) Cutworm spp. Flea Beetle spp. Grasshopper spp. June Beetle spp. Leaffooted Bug Leafhopper spp. Lygus Bug spp. ¹ Melonworm Pickleworm Plant Bug spp. Rindworm spp. complex Saltmarsh Caterpillar Squash Beetle Squash Bug spp. Squash Vine Borer spp. Stink Bug spp. Thrips spp. ^{1,2} Tobacco Budworm ¹ Webworm spp.	0.02-0.03	2.56-3.84	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area. When applied by ground, use a minimum of 10 gal solution per acre.</p> <p>Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Monitor insect populations to determine timing and frequency of applications. Scout fields at a minimum of 5 day intervals.</p> <p>Apply in sufficient volume to ensure sufficient coverage of foliage.</p> <p>Insects that bore or tunnel into leaves, vines, stems or fruit must be controlled before penetration. Only exposed insects (larvae and/or adults) can be controlled with foliar applications of CRUSADER® 1EC.</p> <p>¹See resistance statement under PRODUCT INFORMATION. ²Does not include Western Flower Thrips. ³Suppression only.</p>
Aphid spp. ¹ Leafminer spp. ^{1,3} Spider Mite spp. ³ Whitefly spp. ^{1,3}	0.03	3.84	

Restrictions:

- **DO NOT** apply more than 0.18 lb a.i. (1.44 pts of product) per acre per season.
- **DO NOT** apply within 1 day of harvest.

FRUITING VEGETABLES (EXCEPT CUCURBITS) CROP GROUP

Including: Eggplant, Ground Cherry, Pepino, Peppers (bell and nonbell), Tomatillo, Tomato

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Cabbage Looper Cutworm spp. Hornworm spp.	0.015-0.025	1.92-3.20	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Make applications when pests appear. If needed, make repeat applications after at least 5 days. Apply in sufficient volume to ensure sufficient coverage of foliage.</p> <p>¹For control of first and second instar only.</p> <p>²Suppression only.</p> <p>³See resistance statement under PRODUCT INFORMATION.</p> <p>⁴For control before the larva bores into the plant stalk or fruit.</p> <p>⁵Does not include Western Flower Thrips.</p>
Aphid spp. ^{2,3} Beet Armyworm ^{1,3} Blister Beetle spp. Colorado Potato Beetle ³ Cucumber Beetle spp. (Adult) European Corn Borer ⁴ Fall Armyworm ¹ Flea Beetle spp. Grasshopper spp. Japanese Beetle (Adult) Leafhopper spp. Leafminer spp. ² Meadow Spittlebug Pepper Weevil (Adult) ² Plant Bug spp. Southern Armyworm ¹ Spider Mite spp. ² Stalk Borer ⁴ Stink Bug spp. Thrips ⁵ Tobacco Budworm ³ Tomato Fruitworm Tomato Pinworm Tomato Psyllid ^{2,3} Vegetable Weevil (Adult) Whitefly spp. ^{2,3} Yellow-striped Armyworm ¹	0.02-0.03	2.56-3.84	

Restrictions:

- **DO NOT** apply within 5 days of harvest.
- **DO NOT** apply more than 0.36 lb a.i. (2.88 pts) per acre per season.

GRASS FORAGE, FODDER, AND HAY

Pasture and Rangeland Grass, Grass Grown for Hay or Silage, Grass Grown for Seed

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Army Cutworm Cutworm spp. Essex Skipper Range Caterpillar Striped Grass Looper	0.015-0.025	1.92-3.20	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p>
Beet Armyworm Billbug spp. ³ Bird Cherry-Oat Aphid ¹ Black Grass Bug Black Turfgrass Beetle (Adult) Blue Stem Midge Cereal Leaf Beetle Chinch Bug Crane Fly spp. Cricket spp. English Grain Aphid ¹ Fall Armyworm Flea Beetle spp. Grass Mealybug Grass Sawfly (Adult) Grasshopper spp. Green June Beetle (Adult) Greenbug ^{1, 2} Japanese Beetle (Adult) Katydid spp. Leafhopper spp. Mite spp. Russian Wheat Aphid ¹ Southern Armyworm Spittlebug spp. Stink Bug spp. Sugarcane Aphid Thrips spp. Tick spp. True Armyworm Webworm spp. Yellowstriped Armyworm	0.02-0.03	2.56-3.84	<p>Monitor insect populations to determine timing and frequency of applications. Scout fields at a minimum of 5 day intervals.</p> <p>Apply in sufficient volume to ensure sufficient coverage of foliage.</p> <p>Chinch bugs: CRUSADER® 1EC may only suppress heavy infestations and/or migrations. In this situation, a second application using an alternative chemistry may be needed.</p> <p>Greenbug: Greenbug is known to have many biotypes. CRUSADER® 1EC may provide suppression only. In this situation, a second application using an alternative chemistry may be needed.</p> <p>Pasture and rangeland grass: May be used for grazing or cut for forage 0 days after application. Grass grown for seed: Straw and mature seed (seed screenings) may be used as feed 7 days after the last application.</p> <p>¹ Best control is obtained before insects begin to roll leaves.</p> <p>² See resistance statement under PRODUCT INFORMATION.</p> <p>³ Suppression only.</p>

Restrictions:

- **DO NOT** apply more than 0.03 lb a.i. (0.24 pts of product) per acre per cutting for pastures, rangeland and grasses grown for seed.
- A minimum re-treatment interval (RTI) of 30 days is required for pastures and rangeland receiving 0.03 lb a.i./A which have not been cut between applications.
- **DO NOT** apply more than 0.09 lb a.i. (0.72 pts of product) per acre per season.
- **DO NOT** cut grass to be dried and harvested for hay until 7 days after the last application.

LEGUME VEGETABLES (SUCCULENT OR DRIED) CROP GROUP

(BEANS AND PEAS) Edible Podded (only)

Canavalia gladiata- sword bean
Canavalia ensiformis – jackbean
Glycine max – Soybean immature seed

Dried Shelled (only)

Lupinus spp. includes: grain, sweet, white and sweet white lupines
Cicer arietinum- chickpea (garbanzo bean)
Cyamopsis tetragonoloba - guar
Lablab purpureus - Lablab bean (hyacinth bean) *Lens esculata* - Lentils

Edible Podded, Succulent Shelled, or Dried Shelled

Phaseolus spp. includes: black, field, kidney, lima, navy, pinto, runner, snap, tepary, and wax beans
Vigna spp. includes: adzuki, asparagus, moth, mung, rice, urd and yardlong beans, black-eyed pea, catjang, Chinese longbean, cowpea, Crowder pea, and Southern pea
Pisum spp. includes: dwarf, edible-pod, English, field, garden, green, snow and sugar snap peas
Cajanus cajan - Pigeon pea Succulent Shelled or Dried Shelled
Vicia faba.- broadbean (favabean)

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Cutworm spp. Green Cloverworm Imported Cabbageworm Mexican Bean Beetle Saltmarsh Caterpillar Velvetleaf Caterpillar	0.015-0.025	1.92-3.20	Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area. Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage Make applications when pests appear. If needed, make repeat applications after at least 5 days. Apply in sufficient volume to ensure sufficient coverage of foliage.
Alfalfa Caterpillar Aphid spp. ⁴ Armyworm ² Bean Leaf Beetle Bean Leafskeletonizer Blister Beetle spp. Corn Earworm Corn Rootworm Beetle spp. (Adult) Cucumber Beetle spp. (Adult) Curculio and Weevil spp. ¹ (foliage and pod feeding adults and larvae) European Corn Borer Fall Armyworm ² Flea Beetle spp. (Adult) Flea Hopper spp. Grasshopper spp. Japanese Beetle (Adult) Leafhopper spp. Leaf-tier spp. Looper spp. Meadow Spittlebug (continued)	0.02-0.03	2.56-3.84	¹ For control before the larva bores into the plant stalk or pods. ² For control of the first and second instar only. ³ For suppression only. ⁴ See resistance statement under PRODUCT INFORMATION . ⁵ Does not include Western Flower Thrips.

(continued)

LEGUME VEGETABLES (SUCCULENT OR DRIED) CROP GROUP (cont.)

(BEANS AND PEAS) Edible Podded (only)

Canavalia gladiata- sword bean
Canavalia ensiformis – jackbean
Glycine max – Soybean immature seed

Dried Shelled (only)

Lupinus spp. includes: grain, sweet, white and sweet white lupines
Cicer arietinum- chickpea (garbanzo bean)
Cyamopsis tetragonoloba - guar
Lablab purpureus - Lablab bean (hyacinth bean) *Lens esculata* - Lentils

Edible Podded, Succulent Shelled, or Dried Shelled

Phaseolus spp. includes: black, field, kidney, lima, navy, pinto, runner, snap, tepary, and wax beans
Vigna spp. includes: adzuki, asparagus, moth, mung, rice, urd and yardlong beans, black-eyed pea, catjang, Chinese longbean, cowpea, Crowder pea, and Southern pea
Pisum spp. includes: dwarf, edible-pod, English, field, garden, green, snow and sugar snap peas
Cajanus cajan - Pigeon pea Succulent Shelled or Dried Shelled
Vicia faba.- broadbean (favabean)

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
(cont.) Painted Lady Butterfly (larvae) Plant Bug spp. including Lygus spp. ⁴ Stalk Borer ¹ Stink Bug spp. Three-cornered Alfalfa Hopper Thrips spp. ^{4,5} Tobacco Budworm ⁴ Webworm spp. Western Bean Cutworm Western Yellow-striped Armyworm ² Yellow-striped Armyworm ²	0.02-0.03	2.56-3.84	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage</p> <p>Make applications when pests appear. If needed, make repeat applications after at least 5 days. Apply in sufficient volume to ensure sufficient coverage of foliage.</p> <p>¹For control before the larva bores into the plant stalk or pods. ²For control of the first and second instar only. ³For suppression only. ⁴See resistance statement under PRODUCT INFORMATION. ⁵Does not include Western Flower Thrips.</p>
Beet Armyworm ^{3,4} Leafminer spp. ^{3,4} Lesser Cornstalk Borer ³ Soybean Looper ^{3,4} Spider Mite spp. ³ Whitefly spp. ^{3,4}	0.03	3.84	

Restrictions:

- For edible podded and succulent shelled legume vegetables, **DO NOT** apply within 7 days of harvest.
- For dried shelled legume vegetables, **DO NOT** apply within 21 days of harvest.
- **DO NOT** apply more than 0.12 lb a.i. (0.96 pt) per acre per season.
- For succulent and dried shelled peas and beans, **DO NOT** graze livestock in treated areas or harvest vines for forage or hay.

LEGUME VEGETABLES

Soybean

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Bean Leaf Beetle Cabbage Looper Corn Earworm Cutworm spp. Green Cloverworm Mexican Bean Beetle Mexican Corn Rootworm Beetle (Adult) Northern Corn Rootworm Beetle (Adult) Painted Lady (Thistle) Caterpillar Potato Leafhopper Saltmarsh Caterpillar Southern Corn Rootworm Beetle (Adult) Soybean Aphid ⁴ Three-Cornered Alfalfa Hopper Thrips spp. ⁵ Velvetbean Caterpillar Western Corn Rootworm Beetle (Adult) Woollybear Caterpillar	0.015-0.02	1.92-3.20	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Make applications when pests appear. If needed, make repeat applications after at least 5 or more days. Apply in sufficient volume to ensure sufficient coverage of foliage.</p> <p>Adult corn rootworm beetles (<i>Diabrotica</i> .species): Use a minimum of 2.56 fl oz per acre (0.02 lb a.i. per acre) as part of an aerial-applied corn rootworm control program.</p> <p>¹Use higher rates for large larvae. ²Suppression only. ³See resistance statement under PRODUCT INFORMATION. ⁴Use lower rates for early season applications and/or lighter populations. ⁵Does not include Western Flower Thrips.</p>
Armyworm ¹ Blister Beetle spp. European Corn Borer Fall Armyworm ¹ Grasshopper spp. Japanese Beetle (Adult) Plant Bug spp. Silverspotted Skipper Stink Bug spp. Tobacco Budworm ³ Webworm spp. Yellow-striped Armyworm ¹	0.025-0.03	3.20-3.84	
Beet Armyworm ^{2,3} Lesser Cornstalk Borer ² Soybean Looper ^{2,3} Spider Mite spp. ²	0.03	3.84	

Restrictions:

- **DO NOT** apply within 30 days of harvest.
- **DO NOT** apply more than 0.06 lb a.i. (0.48 pt) per acre per season.
- **DO NOT** graze or harvest treated soybean forage, straw, or hay for livestock feed.

LETTUCE (HEAD AND LEAF)			
TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Alfalfa Looper Cabbage Looper Cutworm spp. Green Cloverworm Imported Cabbageworm Saltmarsh Caterpillar	0.015-0.025	1.92-3.20	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Make applications when pests appear. If needed, make repeat applications after at least 5 or more days. Apply in sufficient volume to ensure sufficient coverage of foliage.</p> <p>¹For control of first and second instar only. ²Suppression only. ³See resistance statement under PRODUCT INFORMATION.</p>
Aphid spp. ^{2,3} Armyworm Beet Armyworm ^{1,3} Corn Earworm Diamondback Moth ³ European Corn Borer Fall Armyworm ¹ Flea Beetle spp. Grasshopper spp. Japanese Beetle (Adult) Leafhopper spp. Meadow Spittlebug Plant Bug spp. including Lygus spp. ³ Southern Armyworm Spider Mite spp. ² Stink Bug spp. Tobacco Budworm ³ Vegetable Weevil (Adult) Whitefly spp. ^{2,3}	0.02-0.03	2.56-3.84	
Restrictions:			
<ul style="list-style-type: none"> • DO NOT apply within 1 day of harvest. • DO NOT apply more than 0.3 lb a.i. (2.4 pts) per acre per season. 			

ONION (BULB) AND GARLIC			
TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Cutworm spp. Leafminer spp. (Adult) Onion Maggot (Adult) Seedcorn Maggot (Adult)	0.015-0.025	1.92-3.20	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Make applications when pests appear. If needed, make repeat applications after at least 5 or more days. Apply in sufficient volume to ensure sufficient coverage of foliage. Use the higher label rates as thrips population increases and avoid rescue situations.</p> <p>For thrips control by aerial application, the addition of 1% COC v/v, ¼% NIS v/v, or a silicone adjuvant (follow manufacturer's use directions) may enhance the deposition of the spray and increase plant coverage.</p> <p>¹For control of the first and second instars only. ²Suppression only. ³See resistance statement under PRODUCT INFORMATION.</p>
Aphid spp. ² Armyworm spp. ¹ Flower Thrips ^{2,3} Onion Thrips ³ Plant Bug spp. Stink Bug spp. Tobacco Thrips ³ Western Flower Thrips ^{2,3}	0.02-0.03	2.56-3.84	
Restrictions:			
<ul style="list-style-type: none"> • DO NOT apply within 14 days of harvest. • DO NOT apply more than 0.24 lb a.i.(1.92 pts) per acre per season. 			

PEANUT

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Cutworm spp. Green Cloverworm Potato Leafhopper Red-necked Peanut Worm Threecornered Alfalfa Hopper Velvetbean Caterpillar	0.015-0.025	1.92-3.20	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Make applications when pests appear. If needed, make repeat applications after at least 7 or more days. Apply in sufficient volume to ensure sufficient coverage of foliage.</p> <p>¹Use higher rates for large larvae. ²Suppression only. ³See resistance statement under PRODUCT INFORMATION.</p>
Bean Leaf Beetle Corn Earworm Fall Armyworm ¹ Grasshopper spp. Southern Corn Rootworm (Adult) Stink Bug spp. Tobacco Thrips Vegetable Weevil Whitefringed Beetle (Adult)	0.02-0.03	2.56-3.84	
Aphid spp. ² Beet Armyworm ^{2,3} Lesser Cornstalk Borer ² Soybean Looper ^{2,3} Spider Mite spp. ²	0.03	3.84	

Restrictions:

- **DO NOT** apply within 14 days of harvest.
- **DO NOT** apply more than 0.12 lb a.i (0.96 pt) per acre per season.
- **DO NOT** graze livestock in treated area.
- **DO NOT** use treated vines or hay for animal feed.

POME FRUITS CROP GROUP

Apple, Crabapple, Loquat, Mayhaw, Oriental Pear, Pear, Quince

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Apple Aphid Apple Maggot (Adult) Cherry Fruit Fly spp. (Adult) Codling Moth Green Fruitworm Japanese Beetle Leafhopper spp. Leafroller spp. Lesser Appleworm Omnivorous leafroller Orange Tortrix Oriental Fruit Moth Pear Psylla ¹ Pear Sawfly Periodical Cicada Plant Bug spp. Plum Curculio Rosy Apple Aphid San Jose Scale (fruit infestations only) Spirea Aphid ¹ Stink Bug spp. Tent Caterpillar spp. Tentiform Leaf Miner spp. Tree Borer spp. Tufted Apple Budworm Webworm spp.	0.02-0.04	2.56-5.12	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 5 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Make applications when pests appear. If needed, make repeat applications after at least 5 or more days. Apply in sufficient volume to ensure sufficient coverage of foliage.</p> <p>¹Suppression only.</p>

Restrictions:

- **DO NOT** apply within 21 days of harvest.
- **DO NOT** apply more than 0.2 lb a.i. (1.6 pts) per acre per year.
- **DO NOT** apply more than 0.16 lb a.i. (1.28 pts) per acre per year post bloom.

STONE FRUITS CROP GROUP Apricot, Sweet and Tart Cherry, Nectarine, Peach, Plum, Chickasaw, Plum, Damson Plum, Japanese Plum, Plumcot, Prune

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
American Plum Borer Apple Maggot (Adult) Black Cherry Aphid Cherry Fruit Fly spp. (Adult) Codling Moth Green Fruitworm Japanese Beetle June Beetle Leafhopper spp. Leafroller spp. Oriental Fruit Moth Peach Twig Borer Peachtree Borer spp. Pear Sawfly Periodical Cicada Plant Bug spp. Plum Curculio Rose Chafer Stink Bug spp. Tent Caterpillar spp. Thrips spp.	0.02-0.04	2.56-5.12	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 5 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Make applications when pests appear. If needed, make repeat applications after at least 5 or more days. Apply in sufficient volume to ensure sufficient coverage of foliage.</p>

Restrictions:

- **DO NOT** apply within 14 days of harvest.
- **DO NOT** apply more than 0.2 lb a.i. (1.6 pts) per acre per year.
- **DO NOT** apply more than 0.16 lb a.i.(1.28 pts) per acre per year post bloom.

SUGARCANE

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Mexican Rice Borer ¹ Pygmy Mole Cricket Rice Stalk Borer ¹ Sugarcane Aphid ³ Sugarcane Beetle (Adult) ² Sugarcane Borer ¹ Western Indian Crane-fly Yellow Sugarcane Aphid ³	0.025-0.04	3.20-5.12	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Make applications when pests appear. If needed, make repeat applications after at least 7 or more days. Apply in sufficient volume to ensure sufficient coverage of foliage.</p> <p>¹For control before the larva bores into the plant stalk. ²Suppression only of beetles active above ground. ³See resistance statement under PRODUCT INFORMATION.</p>

Restrictions:

- **DO NOT** apply within 21 days of harvest.
- **DO NOT** apply more than 0.16 lb a.i. (1.28 pts) per acre per season.

SUNFLOWER			
TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Cutworm spp. Sunflower Beetle	0.015-0.025	1.92-3.20	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Make applications when pests appear. If needed, make repeat applications after at least 5 or more days. Apply in sufficient volume to ensure sufficient coverage of foliage.</p> <p>1For control of first and second instar only. 2Suppression only. 3See resistance statement under PRODUCT INFORMATION.</p>
Banded Sunflower Moth Fall Armyworm ¹ Grasshopper spp. Head-Clipper Weevil (Adult) Japanese Beetle (Adult) Leafhopper spp. Meadow Spittlebug Painted Lady (Thistle) Caterpillar Seed Weevil (Adult) Spotted Cabbage Looper Stem Weevil (Adult) Stink Bug spp. Sunflower Maggot (Adult) Sunflower Moth Woollybear Caterpillar	0.02-0.03	2.56-3.84	
Beet Armyworm ^{2,3} Spider Mite spp. ²	0.03	3.84	
<p>Restrictions:</p> <ul style="list-style-type: none"> • DO NOT apply within 45 days of harvest. • DO NOT apply more than 0.12 lb a.i. (0.96 pt) per acre per season. DO NOT apply more than 0.09 lb a.i. (0.72 pt) per acre per season after bloom initiation. • DO NOT apply as an ultra-low volume (ULV) spray. 			

TOBACCO

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Armyworm spp. ¹ Blister Beetle spp. Cabbage Looper Corn Earworm Cucumber Beetle spp. (Adult) Cutworm spp. Grasshopper spp. Japanese Beetle (Adult) Katydid spp. Plant Bug spp. ³ Potato Tuberworm Salt Marsh Caterpillar Stinkbug spp. Tobacco Aphid spp. ^{2,3} Tobacco Budworm ² Tobacco Flea Beetle (Adult) Tobacco Hornworm Tobacco Thrips spp. ² Tomato Hornworm Tree Cricket spp. Vegetable Weevil (Adult) Webworm spp.	0.015-0.03	1.92-3.84	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Make applications when pests appear. If needed, make repeat applications after at least 7 or more days. Apply in sufficient volume to ensure sufficient coverage of foliage.</p> <p>1For control of first and second instar only. 2Suppression only. 3See resistance statement under PRODUCT INFORMATION.</p>

Restrictions:

- **DO NOT** apply within 40 days of harvest.
- **DO NOT** apply more than 0.09 lb a.i. (0.72 pt) per acre per year.

TREE NUTS CROP GROUP

Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory Nut, Macadamia Nut (Bush Nut), Pistachio, Walnut, Black Walnut, English (Persian)

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Ants Chinch Bug Codling Moth Filbertworm Leaffooted Bug Leafroller spp. Navel Orangeworm Peach Twig Borer Plant Bug spp. Stink Bug spp. Walnut Aphid Walnut Husk Fly spp. (Adult)	0.02-0.04	2.56-5.12	<p>Ground application: Apply in or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 5 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Make applications when pests appear. If needed, make repeat applications after at least 5 or more days. Apply in sufficient volume to ensure sufficient coverage of foliage.</p>
<p>For Pecan</p> Hickory Shuckworm Pecan Casebearer spp. Pecan Weevil Pecan Aphid spp. Pecan Spittlebug Stink bug spp. Pecan Phylloxera spp.	0.02-0.04	2.56-5.12	

Restrictions:

- **DO NOT** apply within 14 days of harvest.
- **DO NOT** apply more than 0.16 lb a.i. (1.28 pts) per acre per year.
- **DO NOT** apply more than 0.12 lb a.i. (0.96 pt) per acre per year post bloom.

TUBEROUS AND CORM VEGETABLES CROP GROUP

Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem only), Canna (edible), Cassava (bitter and sweet), Chayote (root), Chufa, Dasheen, Ginger, Leren, Potato, Sweet Potato, Tanier, Turmeric, Yam (bean and true)

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Cutworm spp. Leafhopper spp. Saltmarsh Caterpillar Sweet Potato Hornworm Woollybear Caterpillar spp.	0.015-0.025	1.92-3.20	Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area. Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.
Aphid species ¹ Armyworm spp. ¹ Blister Beetle spp. Colorado Potato Beetle ¹ Corn Earworm Cricket spp. Cucumber Beetle spp. (adults) European Corn Borer Flea Beetle spp. (adults) Grasshopper spp. Looper spp. ¹ Lygus Bug spp. ¹ Plant Bug spp. Potato Psyllid Potato Tuberworm Stink Bug spp. Sweet Potato Leaf Beetle (adults) Sweet Potato Vine Borer Thrips spp. ^{1,2} Tortoise Beetle spp. Webworm spp. Weevil spp. (adults)	0.02-0.03	2.56-3.84	Make applications when pests appear. If needed, make repeat applications after at least 7 or more days. Apply in sufficient volume to ensure sufficient coverage of foliage. Insects that bore or tunnel into leaves, vines, stems, tubers or corms must be controlled before penetration. Only exposed insects (larvae and/or adults) can be controlled with foliar applications of CRUSADER® 1EC. ¹ See resistance statement under PRODUCT INFORMATION . ² Does not include Western Flower Thrips. ³ Suppression only.
Leafminer spp. ^{1,3} Whitefly spp. ^{1,3} Spider Mite spp. ³	0.03	3.84	

Restrictions:

- **DO NOT** apply more than 0.12 lb a.i. (0.96 pt) per acre per year per season.
- **DO NOT** apply within 7 days of harvest.

CONIFER AND DECIDUOUS TREES

Plantations and Nurseries

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Bagworm Balsam Twig Aphid Birch Leafminer Black Pine Weevil Elm Leaf Beetle European Elm Bark Beetle Gypsy Moth Japanese Beetle June Beetle spp. Leaf Beetle spp. Leafroller spp. May Beetle spp. Mealybug spp. ¹ Pales Weevil Pine Chafer Pine Colaspis Beetle Pine Conelet Bug Pine Leaf Chermid Balsam Wooly Aphid Pine Needle Scale Pine Sawfly spp. Pine Tip Moth spp. Pine Tortoise Scale Pine Weevil spp. Poplar Aphid spp. Sawfly spp. Spittlebug spp. Spruce Budworm Tent Caterpillar spp. Tussock Moth spp. Webworm spp.	0.02-0.04	2.56-5.12	<p>Ground application: Apply in sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Air application: Apply in a minimum of 2 gals per acre or sufficient spray volume to obtain full coverage of the foliage or target area.</p> <p>Make applications when pests appear. Apply in sufficient volume to ensure sufficient coverage of foliage.</p> <p>To control exposed foliage, flower, cone, seed, and bark feeding insects, apply as required by scouting.</p> <p>¹Suppression only.</p>

Restrictions:

- **DO NOT** apply more than 0.24 lb ai (1.92 pts) per acre per year.

CONIFER AND DECIDUOUS TREES

Seed Orchards

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
Coneworm spp. Seed Bug spp. Thrips spp.	See Remarks	See Remarks	For high volume sprayers, dilute 5.12 fl oz per 100 gals of water and apply 5-10 gals of finished spray per tree. For low volume sprayers, dilute 20 fl oz per 100 gals of water and apply 100 gals of finished spray per acre. For aerial applications, apply 15 fl oz/A in a minimum of 10 gals finished spray per acre.

Restrictions:

- **DO NOT** apply more than 0.5 lb a.i. (4 pts) per acre per year.

NON-AGRICULTURAL USES

NON-CROPLAND

(Excluding Public Land)

TARGET PESTS	RATE		REMARKS
	lb a.i./A	fl oz/A	
See specific agricultural crop listing on this CRUSADER® 1EC label for target pests and rates.	See specific agricultural crop listing	See specific agricultural crop listing	Spray non-cropland adjacent to agricultural areas to control migratory insects which may threaten crops. Follow use directions, rates, and spray directions found elsewhere on this label for the adjacent crop and target pests. Use highest labeled rates for dense/large foliage, high insect populations and larger larval stages. Repeat as necessary to maintain control.

Restrictions:

- **DO NOT** exceed 0.2 lb a.i. (1.6 pts) per acre per year.
- **DO NOT** graze livestock in treated areas.

RATE CONVERSION CHART

Lb ai/A	Fl Oz/A	Pints/A	Treated Acres/ Gallon of product
0.015	1.92	0.12	66
0.02	2.56	0.16	50
0.025	3.20	0.20	40
0.03	3.84	0.24	33
0.04	5.12	0.32	25

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: DO NOT freeze. **DO NOT** store below 40°F. If solid crystals are observed, warm material to above 60°F by placing container in warm location. Shake or roll container periodically to redissolve solids.

Keep out of reach of children and animals. Store in original container only. Store in cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. **DO NOT** put concentrate or dilute material into food or drink containers. **DO NOT** contaminate other pesticides, fertilizers, water, food or feed by storage or disposal. In case of spill, avoid contact and isolate area. Keep out animal and unprotected persons. Call (800) 424-9300 CHEMTREC.

To confine spill (if liquid) dike surrounding area or absorb with sand, cat litter. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. **DO NOT** contaminate water, food or feed by storage or disposal. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. Dispose of excess waste or pesticide by use according to label directions or contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of federal law. Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to federal, state, or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office. Open dumping is prohibited.

CONTAINER HANDLING:

Non-refillable container. **DO NOT** reuse or refill this container. Triple rinse container (or pressure rinse as follows).

Fill the container ¼ full with appropriate solvent. Replace and tighten closures. Tip container on its side and roll back and forth, ensuring at least one complete revolution for 30 seconds. Stand the container on its end and flip back and forth several times. Turn the container over onto its other end and flip back and forth several times. Empty the rinsate into a storage container for later use or disposal or into a mix tank for product formulation. Repeat this procedure two more times. Pressure rinse. Empty the remaining contents into a storage container or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over the storage container or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rise at about 40 PSI for at least 30 seconds. Drain for 10 seconds after flow begins to drip. An appropriate solvent would include a solvent that will dissolve technical material. The rinsate may be used later to formulate a product or may be disposed in accordance with local state and national environmental laws, rules, standards and regulations.

ALMACENAMIENTO Y DESECHO

No contamine agua, comida ni forrajes mediante el almacenamiento y desecho.

ALMACENAMIENTO DE PESTICIDAS: No congelar. No almacene a menos de 40 °F. Si se observan cristales, caliente el material a más de 60 °F colocando el recipiente en un lugar cálido. Agite o haga rodar el recipiente periódicamente para volver a disolver los sólidos.

Manténgase fuera del alcance de los niños y los animales. Almacene únicamente en los envases originales. Conservar en lugar fresco y seco y evitar el exceso de calor. Abra los recipientes con cuidado. Después de un uso parcial, vuelva a colocar las tapas y cierre herméticamente. No ponga material concentrado o diluido en recipientes de alimentos o bebidas. En caso de derrame, evite el contacto, aisle el área y mantenga fuera a los animales y personas sin protección. Confinar los derrames. Llame a CHEMTREC (Transporte y Derrames): (800)-424-9300.

Para confinar el derrame, haga un dique en el área circundante o absorba con arena, arena para gatos o arcilla comercial. Coloque el paquete dañado en un contenedor de almacenamiento. Identificar contenidos.

DESECHO DE PESTICIDAS: Los residuos de pesticidas son muy peligrosos. La disposición inadecuada del pesticida sobrante, mezcla de aerosol o aguas de enjuague de contenedores, constituye una violación de la Ley Federal. Si estos residuos no pueden ser eliminados de acuerdo a las instrucciones de la etiqueta, contacte la agencia para el manejo de pesticidas o control ambiental de su estado, o el representante para el Manejo de Residuos Peligrosos de la Oficina Regional de la EPA más cercana para obtener indicaciones del proceso a seguir para su disposición.

DESECHO DE PESTICIDAS: Los desechos resultantes del uso de este producto deben eliminarse en el sitio o en una instalación de eliminación de desechos aprobada. Se prohíbe el vertido a cielo abierto.

DISPOSICIÓN DEL ENVASE:

No reutilice ni rellene este recipiente. Enjuague tres veces (o equivalente). Una vez limpio, ofrezca el envase para ser reciclado o reacondicionado, o perfore y deseche en un relleno sanitario apropiado, o deseche utilizando otros métodos aprobados por las autoridades estatales y locales.

Enjuague tres veces de la siguiente manera: Vacíe el contenido restante en el equipo de aplicación o en un tanque de mezcla. Llena el recipiente hasta ¼ de su capacidad con agua. Vuelva a colocar y apriete los cierres. Incline el recipiente de costado y gírelo hacia adelante y hacia atrás, asegurando al menos una revolución completa, durante 30 segundos. Coloque el recipiente sobre su extremo e inclínelo hacia adelante y hacia atrás varias veces. Voltee el recipiente sobre su otro extremo e inclínelo hacia adelante y hacia atrás varias veces. Vacíe el líquido de enjuague en el equipo de aplicación o en un tanque de mezcla o almacene el líquido de enjuague para su posterior uso o eliminación. Repita este procedimiento dos veces más. Enjuague a presión de la siguiente manera: Vacíe el contenido restante en el equipo de aplicación o en un tanque de mezcla y continúe drenando durante 10 segundos después de que el flujo comience a gotear. Sostenga el recipiente boca abajo sobre el equipo de aplicación o el tanque de mezcla o recoja el agua de enjuague para su posterior uso o eliminación. Inserte la boquilla de enjuague a presión en el costado del recipiente y enjuague a aproximadamente 40 PSI durante al menos 30 segundos. Drene durante 10 segundos después de que el flujo comience a gotear.

CONDITIONS OF SALE, WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY

ALBAUGH, LLC and the Seller offer this product subject to the following Conditions of Sale, Warranty Disclaimer and Limitation of Liability, and by purchasing and using this product, Buyer and Users accept such Conditions of Sale, Warranty Disclaimer and Limitation of Liability. No employee or agent of ALBAUGH, LLC or the Seller is authorized to modify the terms of these Conditions of Sale, Warranty Disclaimer and Limitation of Liability unless in writing and signed by a duly authorized representative of ALBAUGH, LLC.

The **DIRECTIONS FOR USE** of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of ALBAUGH, LLC or the Seller. All such risks shall be assumed by the Buyer. **WITHOUT LIMITATION, THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.**

TO THE FULLEST EXTENT PERMITTED BY APPLICABLE LAW, ALBAUGH, LLC MAKES NO EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR ANY OTHER EXPRESS OR IMPLIED WARRANTY THAT EXTENDS BEYOND THE STATEMENTS MADE ON THIS LABEL. BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S AND SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OF HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. When Buyer suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), Buyer must promptly notify Seller in writing of any claims to be eligible to receive either remedy stated above. **IN NO CASE SHALL ALBAUGH, LLC OR THE SELLER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**