

EQUIP®

POSTEMERGENCE, ONE-PASS WEED CONTROL WITH TWO MODES OF ACTION FOR CONTACT AND RESIDUAL CONTROL OF BROADLEAF AND GRASS WEEDS IN FIELD CORN GROWN FOR GRAIN OR SILAGE.



Equip[®] is for use as a postemergence herbicide for selective burndown and residual control of listed annual grass and broadleaf weeds in field corn grown for grain or silage.

To improve residual or burndown control, Equip can be tank mixed with a variety of herbicides. Equip works best when used in a planned sequential application herbicide program, following a preplant application of another labeled corn herbicide. Consult the label of the corn herbicide product for specific use directions and additional information.

For postemergence application of Equip, if activating rainfall or sprinkler irrigation (>1/2") is not received within 5-7 days following application, cultivate or follow with a sequential application of a nicosulfuron-containing product, or glyphosate, as necessary. Equip is rainfast in four hours.

PRODUCT FEATURES

- Long-lasting control with contact plus residual weed control one-pass system.
- Broad-spectrum control of broadleaves and grasses.
- Controls tough weeds found in corn.
- \bullet Two effective modes of action with built-in safener for flexibility.
- Convenient dry formulation for easier clean-up.
- \bullet Mix with at razine or glyphosate to broaden weed spectrum and residual.

ACTIVE INGREDIENTS

31.25% mesotrione + 7.5% rimsulfuron

HRAC GROUPS

27, 2

FORMULATION

Dry-flowable

REI

12 hours

COMPARE TO

Realm® Q

MODES OF ACTION

HPPD inhibitor and acetolactate synthase inhibitor

SIGNAL WORD

Caution

RESTRICTED USE

No

PACKAGING

8 x 50 oz

EPA REG NUMBER

83100-57-83979

See label for complete application guidelines and restrictions.





FREQUENTLY ASKED QUESTIONS

Can I use Equip on seed, sweet or popcorn?

• No. Equip is only labeled for use on field corn grown for grain or silage.

Do I need to use a surfactant with Equip?

• Applications of Equip must include a crop oil concentrate (COC) or a high surfactant oil concentrate (HSOC) for control of emerged weeds. The use of non-ionic surfactant (NIS) instead of COC or HSOC is allowed, but the level of weed control achieved when using a COC or HSOC is consistently better than when using NIS. The use of methylated seed oil (MSO) adjuvants or MSO blend adjuvants may result in severe crop injury. MSO adjuvants are NOT recommended. In addition to COC or HSOC, always add spray grade UAN (e.g. 28-0-0) to the spray solution or AMS, unless precluded elsewhere on the label.

What are my best tank-mix options for one-pass postemergence treatment?

- Application of Equip may be made as a sequential application in a planned postemergence weed control program in corn, following a preemergence herbicide. Consult the preemergence herbicide label for use restrictions, application information, rotational crop guidelines and cautionary statements before making applications of Equip.
- Glyphosate may be used on Roundup Ready® corn after weeds emerge, but before they reach the maximum size listed on the glyphosate herbicide label.
- Glufosinate may be used on LibertyLink® corn after weeds emerge, but before they reach the maximum size listed on the glufosinate herbicide label.

Do I need to use a preemergence product if I use Equip?

• Equip contains both a Group 27 and Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to Group 27 and/or Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies must be followed, including, but not limited to: planting into weed-free fields, rotating the use of this product or other Group 27 and/or Group 2 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field, using tank mixtures with herbicides from a different group if such use is permitted, adopting an integrated weed-management program for herbicide use, and scouting after herbicide application to monitor weed populations for early signs of resistance development.

What are the rotational restrictions for Equip?

- Corn anytime
- Cereals, winter wheat 4 months
- Cereals, spring (wheat, barley, oats) 9 months
- Alfalfa, canola, soybeans, cotton, potatoes 10 months

Refer to label page 13 for all other crop rotation intervals



 \blacktriangleleft See label for complete application guidelines and restrictions.

©2023 Albaugh, LLC. All rights reserved. Albaugh, the Albaugh logo and Equip are registered trademarks of Albaugh, LLC. Realm is a registered trademark of Corteva Agriscience. LibertyLink is a registered trademark of BASF Corporation. Roundup Ready is a registered trademark of Monsanto Technology LLC. Always read and follow label restrictions, precautions and directions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.

