



FIFRA Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

Quinstar® 4L

Herbicide

EPA Reg. No. 42750-169

EPA SLN No. OR-200011

For post-emergence weed control in non-bearing filberts (hazelnuts) only

This SLN label is valid until December 31, 2025 or until otherwise amended, withdrawn, canceled, or suspended.

ACTIVE INGREDIENT:

Quinclorac: 3,7-dichloro-8-quinolinecarboxylic acid..... 40.0%

OTHER INGREDIENTS:..... 60.0%

TOTAL:..... 100.0%

Contains 3.8 pounds of quinclorac per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

ENVIRONMENTAL HAZARDS

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Keep out of lakes, ponds and streams. Do not apply directly to water, areas where surface water is present, or to intertidal areas below the mean high water mark, except as specified on this label for use in rice. Do not contaminate water by cleaning of equipment or disposal of rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This Special Local Need (SLN) label and the federal label for this product must be in the possession of the user at the time of application.

Read this SLN labeling and the entire label affixed to the product container before applying.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the label affixed to the product container, and on this SLN labeling.

POST-EMERGENCE WEED CONTROL IN NON-BEARING FILBERTS (HAZELNUTS)

For post-emergence weed control in non-bearing hazelnuts only to control field bindweed, hedge bindweed, barnyard grass, Canada thistle (suppression only), and large crabgrass.

SPRAY MIXTURE PREPARATION:

Use a minimum of 10 gallons of water per acre. Add one-half the required amount of water to the spray tank, then add this product with agitation, then add any tank mix partners and finally, add the balance of the water with continued agitation. Add crop oil concentrate at 1% v/v (2 pints / acre) in the spray mixture.

APPLICATION INSTRUCTIONS:

Mix 12.6 fl. oz. of Quinstar 4L (0.375 lbs quinclorac) per acre treated. For best performance, apply in the spring when field bindweed shoots are a few inches long. For best long-term bindweed control, make yearly applications of Quinstar 4L in the fall. Apply as a banded, soil application on each side of the hazelnut rows. Quinstar 4L can be applied to the tree base, but it does not control hazelnut suckers. Apply by ground application equipment only.

RESTRICTIONS & LIMITATIONS:

DO NOT apply more than 12.6 fl. oz. of Quinstar 4L (0.375 lbs quinclorac) per acre per application.

DO NOT apply more than 25.2 fl. oz. of Quinstar 4L (0.75 lbs quinclorac) per acre per year.

DO NOT make more than 2 applications per year.

DO NOT make a second application within 30 days after first application.

DO NOT apply by ground when wind speed is greater than 10 mph.

DO NOT apply by air.

DO NOT apply through any type of irrigation equipment (i.e., chemigation).

DO NOT allow livestock to graze in treated areas.

DO NOT harvest hazelnuts treated under this SLN registration for at least 1 year (365 days), starting from the date the product was last applied.

NOTICE

Read the "CONDITIONS OF SALE AND WARRANTY" in the label booklet for this product before using this product. Those terms apply to this SLN labeling and if those terms are not acceptable, return the product unopened at once.

**Section 24(c) Registrant:
Albaugh, LLC
1525 NE 36th Street
Ankeny, Iowa 50021**