



MiCrop™ Fungicide

EPA Reg. No. 45002-24

For Use Only in the State of IDAHO, OREGON, and WASHINGTON

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE USE DIRECTIONS AND RESTRICTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL AND/OR ILLEGAL RESIDUES.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

CANOLA

Target Diseases:

Alternaria black spot (*Alternaria brassicae*), Blackleg/Phoma (*Leptosphaeria maculans*), Cercospora leafspot (*C. brassiciola*), Head rot (*Rhizoctonia solani*), Powdery mildew (*Erysiphe polygoni*), Sclerotica stem rot (*Sclerotinia sclerotiorum*)

Dosage and Frequency/Timing of Applications:

10.5 – 14.5 fl. oz. product/A: For Phoma control, apply during the rosette stage between 2nd true leaf and bolting. For Alternaria and Sclerotica, make an application at 10-25% flowering (3-7 days following first flower). Use the high rate under heavy disease pressure or when conditions are favorable for disease. For Alternaria alone, apply at pod stage (approximately 95% petal fall). Specifically, for blackleg, apply at the 2- to 4-leaf stage. For other foliar diseases, apply at the first sign of disease. For head rot, apply at 50% flowering.

Dilution Rate:

MiCrop™ Fungicide may be applied by ground or aerial application. Use a minimum of 10 gallons of water per acre for ground applications.

Tank Mixes:

Other registered products may be tank mixed with MiCrop™ Fungicide. Read and follow the applicable Restrictions and Limitations and Directions For Use on products involved in tank mixing. The most restrictive labeling applies to tank mixes.

RESTRICTIONS AND LIMITATIONS – CANOLA

1. Do not apply more than 14.5 fl. oz./A/year of MiCrop™ Fungicide.
2. Do not apply more than 0.113 lb. a.i. of propiconazole-containing products/A/year.
3. Do not apply more than 0.45 lb. a.i. of azoxystrobin-containing products/A/year.
4. Do not apply more than 1 application/A/year
5. Do not apply within 30 days of harvest (30-day PHI).

Note: All applicable directions, restrictions, and precautions on the MiCrop™ Fungicide package label are to be followed. This labeling contains supplemental instructions and restrictions for use of this product which may not appear on the package label. Read and follow all precautionary statements and all other information appearing on the product label.

AD051724

Manufactured By:
Albaugh, LLC
Ankeny, IA 50021