



PRODUCT BULLETIN

OXYSTAR™ 2E

EPA Reg. No. 42750-136
Special 2(ee) Recommendation†

**For Distribution and Use in the States of Alabama, Arkansas,
Georgia, Louisiana, Mississippi, Missouri, North Carolina,
Oklahoma, South Carolina, Tennessee, Texas Virginia
(Not For Use On Fallow Beds To Be Planted To Soybeans In California)**

GROUND OR AERIAL APPLICATION OF OXYSTAR™ 2E ON FALLOW BEDS (To Be Planted To Cotton or Soybeans)

ATTENTION

†This recommendation is made as permitted under FIFRA Section 2(ee) and has not been reviewed or approved by EPA.

- This recommendation must be in the possession of the user at the time of application.
- Read the label affixed to the container for Oxystar™ 2E herbicide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Oxystar™ 2E is subject to all use precautions and limitations imposed by the label affixed to the product container.

DIRECTIONS FOR USE

Oxystar™ 2E is effective as a preemergence and/or postemergence herbicide when used alone or in a tank mix combination with Gly Star® Plus or paraquat (Gramoxone) for the control of winter annual broadleaf weeds in fallow beds to be planted to either cotton or soybeans.

WEEDS SUPPRESSED BY PREEMERGENCE APPLICATION

Oxystar™ 2E may be applied at 1 to 2 pints (0.25 to 0.5 lb active) per broadcast acre. Oxystar™ 2E should provide preemergence suppression of the following weeds when used at recommended dosages and weed stage.

HORSEWEED (MARESTAIL) *Coryza Canadensis*
RYEGRASS, ITALIAN *Lolium multiflorum*

Manufactured By:
Albaugh, Inc.
1525 NE 36th Street
Ankeny, IA 50021