


PLOTTER®

SELECTIVE POSTEMERGENCE HERBICIDE FOR CONTROL OR SUPPRESSION OF BROADLEAF AND BRUSH WEEDS.



Plotter controls weeds by postemergence activity. For best results, apply to young, actively growing weeds. The use rate depends upon the weed spectrum and size of weeds at application. The degree and duration of control may depend on weed spectrum and infestation intensity, weed size at application and environmental conditions at and following treatment.

PRODUCT FEATURES

- Effective broadleaf control in wheat and barley crops.
- Excellent tank-mix partner with other registered herbicides for enhanced broadspectrum control.
- Great tool for hard-to-control weeds in pastures and rangeland.
- Controls or suppresses over 30 broadleaf and brush weeds.



ACTIVE INGREDIENT

0.60 lb/gal metsulfuron-methyl

HRAC GROUP

2

FORMULATION

Dry-flowable

REI

4 hours

COMPARE TO

Ally® XP

MODE OF ACTION

ALS inhibitor

SIGNAL WORD

Caution

RESTRICED USE

No

PACKAGING

 8×8 oz bottles (256 oz cases), 8×16 oz bottles (128 oz cases). 2×12 lb

EPA REG NUMBER

83100-58-83979

See label for complete application guidelines and restrictions.





FREQUENTLY ASKED QUESTIONS

Do I need to use a surfactant with Plotter herbicide?

- Applications of Plotter must include either a nonionic surfactant or a crop oil concentrate, except for grain sorghum. In addition, an ammonium nitrogen fertilizer may be used.
- If another herbicide is tank mixed with Plotter agricultural herbicide select adjuvants authorized for use with both products. Products must contain only EPA-exempt ingredients. Antifoaming agents may be needed.
- Refer to label page 7 for specific surfactant recommendations.

Can I tank mix Plotter with other herbicides?

- Plotter may be tank mixed with other suitable registered herbicides to control weeds listed under WEEDS SUPPRESSED, weeds resistant to Plotter, or weeds not listed under WEEDS CONTROLLED. Read and follow all manufacturer's label instructions for the companion herbicide. If those instructions conflict with this label, do not tank mix the herbicide with Plotter.
- Refer to label page 8-9 for specific tank mix recommendations.

What are the rotational restrictions with Plotter?

- Before using Plotter herbicide, carefully consider your crop rotation plans and options. For rotational flexibility, do not treat all of your wheat, barley, grain sorghum, fallow, pasture, or rangeland acres at the same time.
- Do not use Plotter on soils having a pH above 7.9 as extended soil residual activity could extend crop rotation intervals beyond normal. Under certain conditions, Plotter could remain in the soil for 34 months or more, injuring wheat and barley. In addition, other crops planted in high pH soils can be extremely sensitive to low concentrations of Plotter.
- Refer to label page 9-12 for specific rotational restrictions by state and soil pH.



■ See label for complete application guidelines and restrictions.

©2023 Albaugh, LLC. All rights reserved. Albaugh, the Albaugh logo and Plotter are registered trademarks of Albaugh, LLC. Ally is a registered trademark of FMC Corporation.
Always read and follow label restrictions, precautions and directions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.

