

PRIORITY[®] MA

FOR PRE AND POSTEMERGENCE CONTROL OF
GRASS AND BROADLEAF WEEDS IN CORN.



Priority[®] MA is a preemergence and postemergence herbicide for use in corn (field corn, field corn seed, and field corn silage). This product may also be applied to sweet corn, yellow popcorn and grain sorghum as preemergence applications prior to crop emergence. Use Priority MA as a preemergence application for control of most annual grasses and broadleaf weeds, as an early postemergence application for the control of broadleaf weeds in field corn. Priority MA works to control weeds by interfering with the plant's ability to germinate and develop. This product will not provide consistent control of grasses that have emerged at the time of application.

PRODUCT FEATURES

- Contains three herbicide modes of action to increase weed control spectrum and minimize the development of weed resistance.
- Offers long-lasting, broad-spectrum residual control of annual grasses and broadleaf weeds, including waterhemp, lambsquarters, foxtails and *Amaranthus* species.
- Provides application flexibility from 14 days prior to planting through 12-inch corn.
- Can be applied through ground application equipment.

ACTIVE INGREDIENTS

1.70 lbs/gal metolachlor +
1.66 lbs/gal atrazine +
0.218 lbs/gal mesotrione

HRAC GROUPS

15, 5, 27

FORMULATION

Liquid

REI

24 hours

COMPARE TO

Lexar[®]

MODES OF ACTION

Long-chain fatty acid inhibitor (metolachlor), HPPD inhibitor (mesotrione) and photosynthesis inhibitor (atrazine)

SIGNAL WORD

Warning

RESTRICTED USE

Yes

PACKAGING

265 gal, bulk

EPA REG NUMBER

42750-344 | AD07132022

See label for complete application guidelines and restrictions.

PRIORITY[®] MA

FREQUENTLY ASKED QUESTIONS

Do I need to use an adjuvant with Priority MA?

- A non-ionic surfactant at 0.25% v/v (1 qt/100 gal) may be used when this product is applied after field corn has emerged.
- Using a crop oil concentrate (COC) may result in temporary crop injury. If COC is used as an adjuvant, use at a rate that does not exceed 1% v/v (1 qt/100 gal) or no more than 1 qt/A.
- Unless directed for a specific tank mix on this label or as a part of a supplemental label, do not use methylated seed oil (MSO) or nitrogen-based adjuvants (AMS or UAN) with this product when applied alone to emerged field corn, or when application is made as a postemergence tank mixture with other products.
- Any of these adjuvants may be used at preplant or preemergence timing (where the corn plant has not yet emerged) to increase burndown activity on existing weeds.
- Do not make applications of this product to emerged sweet corn, yellow popcorn or grain sorghum as severe adverse crop response or crop injury may result.
- *Refer to label pages 7-8 for specific directions on the use of adjuvants and additives.*

How should I apply Priority MA for best results?

- Use 3.0 qt of this product per acre, if soil organic matter content is less than 3%. Use 3.5 qt of this product per acre, if soil organic matter content is 3% or greater. This product is not recommended on soils that have greater than 10% organic matter, or poor weed control may occur.
- DO NOT make applications to field corn taller than 12 inches, or apply more than 3.5 qt/A per year.
- DO NOT make applications to grain sorghum that has emerged, or to sorghum grown for forage, sweet sorghum

(sorgo), sudangrass, sorghum-sudangrass hybrids, or dual-purpose sorghum. Do not exceed 3.0 qt per acre per year in sorghum.

- *Refer to label pages 13-16 for specific product application directions.*

What are the rotational restrictions for Priority MA?

- If crop is lost, field corn, field corn seed, field corn silage, sweet corn, yellow popcorn and grain sorghum (seed treated with Concep[®] only) may be replanted immediately. Do not make another application of this product.
- If application of this product is made after June 1, rotating to crops other than corn (all types) or sorghum the following spring may result in crop injury.
- Do not rotate to crops other than corn (all types), cotton, small cereal grains, soybeans, sorghum or peanuts the spring following application of this product.
- For soils having a calcareous surface layer, crop injury may result to soybeans planted the year following application of this product (ex. soils found within the Clarion-Nicollet-Webster soil series in Northern Iowa and Southern Minnesota).
- Do not rotate to soybeans for at least 18 months following application of this product if the combined atrazine rate applied was greater than 2.0 lbs. a.i./A, or equivalent band application rate in the Dakotas (eastern region), KS, MN (western region), and NE, or soybean crop injury may result.
- Where rainfall is sparse or erratic or where irrigation is required, in the High Plains and Intermountain areas of the west, use only when corn (all types) or sorghum is to follow field corn, or a crop of untreated corn (all types) or sorghum is to precede other rotational crops.
- Rotational crop interval is 18 months for all other crops.



◀ See label for complete application guidelines and restrictions.

©2024 Albaugh, LLC. All rights reserved. Albaugh, the Albaugh logo and Priority are registered trademarks of Albaugh, LLC. Lexar and Concep are registered trademarks of a Syngenta Group Company. Priority MA and Lexar are restricted use pesticides. Always read and follow label restrictions, precautions and directions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.