



FIFRA §2(ee) Recommendation

PROPI-STAR EC

EPA Reg. No. 42750-211

FOR CONTROL OF RUST, SCAB AND LEAF SPOT ON ALMONDS IN CALIFORNIA ONLY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

The user must refer to the federally approved PROPI-STAR EC label and read and follow all directions for use, restrictions, and precautions. Users should have this recommendation in their possession at the time of use.

This recommendation is permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the EPA.

CROP	PESTS CONTROLLED	RATE OF APPLICATION PER ACRE	APPLICATION INSTRUCTIONS
ALMONDS	Rust (<i>Tranzschelia discolor</i>)	6.0 - 8.0 fl oz	Apply 6.0 to 8 fl oz (0.18 -0.225 lbs. active) per acre. Begin application approximately 2 - 5 weeks after petal fall when environmental conditions are favorable for disease development. For optimum disease control, apply on a 7- to 14-day application schedule.
	Scab (<i>Cladosporium carpophilum</i>)		
	Alternaria Leaf Spot (<i>Alternaria alternata</i>)	8.0 fl. oz	If monitoring or history indicates the presence of Alternaria, apply 8 oz/A of PROPI-STAR EC in the late spring (mid April to beginning of May) and then repeat the treatment 2-3 weeks later.
<ul style="list-style-type: none"> Do not apply more than 32 fl. oz. per acre per season of PROPI-STAR EC. Do not apply more than 0.90 lb. a.i. propiconazole per acre per season. Do not apply within 60 days of harvest. Do not graze livestock in treated areas or cut treated cover crop for feed. Make no more than two consecutive applications before rotating to a fungicide with a different mode of action (not a Group 3). 			

Albaugh, LLC
Ankeny, IA 50021