

PROPISTAR[®] EC

AN EFFECTIVE BROAD-SPECTRUM FUNGICIDE FOR THE CONTROL OF MANY FOLIAR AND SOILBORNE DISEASES.



Propi-Star[®] EC is a broad-spectrum fungicide that provides disease control on a variety of crops, including fruits, vegetables, cereals, tree nuts, sorghum, peanuts and stone fruits, along with turf and ornamental applications. Propi-Star EC stops fungal growth before sporulation through quick uptake and systemic movement through the plant to protect new growth and improve crop quality.

PRODUCT FEATURES

- Offers broad-spectrum curative and preventative activity to control a variety of diseases including leaf spots, powdery mildew, leaf blights, and rusts.
- Effective against foliar and soilborne diseases.
- Excellent choice as part of an integrated disease and pest management program for the control of fungal pathogens.
- Flexible application by ground, aerial or chemigation methods.

ACTIVE INGREDIENT

3.6 lbs/gal propiconazole

FRAC GROUP

3

FORMULATION

Emulsifiable concentrate

REI

12 hours

COMPARE TO

Tilt[®]

MODE OF ACTION

Sterol biosynthesis inhibitor

SIGNAL WORD

Warning

RESTRICTED USE

No

PACKAGING

4 x 1 gal, 2 x 2.5 gal, 265 gal

EPA REG NUMBER

42750-211 | AD041420

See label for complete application guidelines and restrictions.

PROPISTAR[®] EC

FREQUENTLY ASKED QUESTIONS

Is Propi-Star EC effective on almonds?

- Yes, Propi-Star EC is effective against brown rot blossom blight and anthracnose. For best results, apply 4.0-8.0 fl oz/A of Propi-Star EC at 5-10% bloom and at 50-100% bloom for brown rot blossom blight control. Use the higher rate when disease pressure is severe. For anthracnose control, apply 8.0 fl oz/A at bud break on a 7- to 14-day interval.

Propi-Star EC is most effective when applied and allowed to dry before a rainfall. Use sufficient water to provide thorough coverage. Propi-Star EC may be applied by ground or aerial application at a minimum of 15 GPA. Do not apply within 60 days of harvest.

How should Propi-Star EC be used to control foliar diseases on tree nuts?

- Propi-Star EC can be used to control foliar diseases on a variety of tree nuts, including Brazil nut, cashew, macadamia, pecan, walnut, and pistachio. Applications should be made at 4.0-8.0 fl oz/A at the first sign of disease and repeated on a 7- to 14-day interval.

Refer to the label for specific tree nut recommendations.

Propi-Star EC is most effective when applied and allowed to dry before a rainfall. Use sufficient water to provide thorough coverage.

Propi-Star EC may be applied by ground or aerial application at a minimum of 15 GPA. Do not apply within 60 days of harvest, except on pecans. *Refer to label for specific usage instructions for pecan crops.*

When should Propi-Star EC be applied to wheat for best results?

- For early season suppression of powdery mildew, leaf blight, glume blotch and tan spot, apply 2.0-4.0 fl oz/A in the spring and follow-up with a second application up to Feekes growth stage 10.5 for season-long control.
- For control of leaf diseases including rust, powdery mildew, leaf blight, glume blotch, tan spot, *Helminthosporium* leaf blight, spot blotch and net blotch, protecting the flag leaf is important for maximizing the potential yield. Highest yields are normally obtained when Propi-Star EC is applied when the flag leaf is 50% to fully emerged. Applications may be made no closer than a 14-day interval. The use of an oil-based adjuvant may improve the spray coverage and canopy penetration. Do not apply Propi-Star after Feekes 10.5.4. Do not apply after this stage to avoid possible legal residues.
- For foot rot, apply Propi-Star EC at 4.0 fl oz/A plus half rates of other EPA-registered fungicides, such as Topsin[®] M. Apply at tillering, but before elongation has occurred.
- For *Fusarium* head blight, apply Propi-Star at 4.0 fl oz/A at 50% flowering. Addition of a penetrating-type adjuvant may increase *Fusarium* head blight suppression.



◀ See label for complete application guidelines and restrictions.

©2023 Albaugh, LLC. All rights reserved. Albaugh, the Albaugh logo and Propi-Star are registered trademarks of Albaugh, LLC. Tilt is a registered trademark of A Syngenta Group Company. Topsin is a registered trademark of Nippon Soda Company. Always read and follow label restrictions, precautions and directions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.