FUNGICIDE
PROPICONAZOLE
14.3 SELECT™

SYSTEMIC PREVENTIVE AND CURATIVE FUNGICIDE FOR DISEASE CONTROL.

Propiconazole 14.3 Select™ provides preventive and curative control of many turfgrass and ornamental diseases. **Propiconazole 14.3 Select** inhibits the incorporation of ergosterol into the growing fungal cell membranes. Without ergosterol, the fungal cellular growth is stopped. By disrupting fungal growth at the earliest possible stage, the impact and effects of the disease is minimized. **Propiconazole 14.3 Select** is a broad-spectrum product and works to control over 18 turf diseases.

PRODUCT FEATURES

- Excellent residual control.
- Low use rates.
- Low-odor formulation.
- High tank-mix compatibility.
- Labeled for both turf & ornamental applications.
- Both foliar and systemic properties.

LABELED USES

- Residential and commercial turf
- Parks/Rec
- Athletic fields
- Golf turf
- Landscape ornamentals
- Nursery

See label for complete application guidelines and restrictions.

ACTIVE INGREDIENT
14.3% propiconazole

FRAC GROUP
3

FORMULATION
Micro emulsion

CHEMICAL FAMILY
Triazole

COMPARE TO
Banner Maxx®

MODE OF ACTION
Demethylation inhibitor, inhibits sterol biosynthesis in the fungal cell

RESTRICTED USE
No

SIGNAL WORD
Caution

PACKAGING
12 x 1 pt, 4 x 1 gal, 2 x 2.5 gal

EPA REG NUMBER
89442-17

PROPICONAZOLE 14.3 SELECT™

SYSTEMIC PREVENTIVE AND CURATIVE FUNGICIDE FOR DISEASE CONTROL.

APPLICATION INFORMATION

Disease control in turfgrass and dichondra

Use **Propiconazole 14.3 Select™** in a preventive disease control program. Apply in sufficient water to ensure thorough coverage. Apply after mowing OR allow sprayed area to completely dry before mowing. For control of foliar diseases, allow sprayed area to completely dry before irrigation. For control of soil-borne diseases, **Propiconazole 14.3 Select** can be watered in after application. Under conditions optimum for high disease pressure, use the higher rate and the shorter interval. For optimum turf quality and disease control, use **Propiconazole 14.3 Select** in conjunction with turf management practices that promote good plant health. Evaluate spray additives prior to use. Label directions are based on data obtained with no additives. Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.

DO NOT apply more than 16 fl. oz. per 1,000 sq. ft. per calendar year nor apply more than 5.5 gals. of product per acre per calendar year.

DO NOT graze animals on treated areas. **DO NOT** feed clippings from treated areas to livestock or poultry.

Bermudagrass can be sensitive to **Propiconazole 14.3 Select**. **DO NOT** exceed 4 fl. oz. per 1,000 sq. ft. every 30 days on any variety of bermudagrass. In Florida, do not apply **Propiconazole 14.3 Select** to bermudagrass golf course greens when temperatures exceed 90°F.

Disease control in nurseries (field) and landscape plantings

See label for diseases controlled, application instructions, and use rates.

LABELED TURF DISEASES

Anthracnose	Gray leaf spot	*Poa* patch	Spring dead spot
Brown patch	Leaf spot	Powdery mildew	Stripe smut
Dichondra rust	Melting out	Red thread	Summer patch
Dollar spot	Necrotic ring spot	Rust	Take-all patch
Fusarium patch	Pink patch	Snowmold, gray & pink	Yellow patch
			Zoysia patch

LABELED ORNAMENTAL DISEASES

Conifer blights	Leaf blights/spots	Rust
Flower blight	Powdery mildew	Rust powdery mildew

◄ See label for complete application guidelines and restrictions.

ALBAUGH®
SPECIALTY PRODUCTS