

SELECTIVE HERBICIDE

QUINCLORAC 75DF SELECT™

SELECTIVE POSTEMERGENCE HERBICIDE FOR
EFFICIENT WEED CONTROL.



Quinclorac 75 DF Select™ is a selective postemergence herbicide that can be applied to many types of turf, both residential and non-residential, to control a variety of weeds. **Quinclorac 75 DF Select** can be applied from 7 days before seeding and up to 28 days after emergence on selected weeds.

PRODUCT FEATURES

- Controls crabgrass.
- Provides up to 90 days of control.
- Works on both broadleaf and grassy weeds.
- Selective control = safe for use on turf.
- Use in conjunction with seeding/overseeding.
- Convenient dry flowable formulation.

LABELED USES

- Residential and non-residential turfgrass
- Grounds or lawns around residential and commercial establishments and multi-family dwellings
- Schools, parks, picnic grounds, athletic fields, golf courses, and sod farms
- Houses of worship, cemeteries, military and other institutions, airports and roadsides

See label for complete application guidelines and restrictions.

ACTIVE INGREDIENT

75.0% quinclorac

HRAC GROUP

4

FORMULATION

Dry flowable

CHEMICAL FAMILY

Quinoline carboxylic acid

COMPARE TO

Drive® 75 DF

MODE OF ACTION

Synthetic auxin

RESTRICTED USE

No

SIGNAL WORD

Caution

PACKAGING

6 x 1 lb

EPA REG NUMBER

89442-25

QUINCLORAC 75DF SELECT™

GENERAL INFORMATION

Quinclorac 75 DF Select™ is an auxin agonist and is classified as a quinoline carboxylic acid. It is absorbed by foliage and roots and translocated throughout the plant. The control symptoms exhibited by broadleaf weeds include leaf and stem curl or twisting, and chlorosis. Susceptible grasses demonstrate stunting, chlorosis, and gradual reddening followed by necrosis and death. Apply **Quinclorac 75 DF Select** to actively growing weeds as postemergence broadcast or spot sprays using the turf species, rate, and growth stages indicated on the label.

The use of **Quinclorac 75 DF Select** before or after seeding or overseeding a turf area will not significantly interfere with the turfgrass seed germination and growth of those grass types identified as tolerant or moderately tolerant on the label.

BROADLEAF WEEDS CONTROLLED

Bindweed, field	Geranium, Carolina
Clover, hop	Medic, black
Clover, red	Morningglory spp.
Clover, white	Speedwell, common
Daisy, English ^{2,5}	Speedwell, slender
Dandelion, common ²	Speedwell, thymeleaf
Dollarweed	Violet, wild

GRASSY WEEDS SUPRESSED

Barnyardgrass	Foxtail, yellow ¹
Crabgrass, large ^{1,4}	Kikuyugrass ^{2,3}
Crabgrass, smooth ^{1,4}	Signalgrass, broadleaf
Foxtail, giant ¹	Torpedograss ³
Foxtail, green ¹	

¹ Under certain conditions, application of **Quinclorac 75 DF Select** made to annual grasses at 2 to 4 tiller may not provide complete control. A sequential application will be required for grass control in these situations. Optimum control is achieved when applications of **Quinclorac 75 DF Select** + methylated seed oil are applied either before second tiller or as weed grasses mature.

² Tank mix partner or sequential application required.

³ Make 2 sequential applications of 1.0 fl. oz. (0.51 lb. AI/A) of **Quinclorac 75 DF Select** per 1,000 sq. ft. and an additional sequential application up to 0.90 fl. oz. (0.46 lb. AI/A) of **Quinclorac 75 DF Select** per 1,000 sq. ft. at 14- to 21-day intervals.

⁴ Biotypes of large and smooth crabgrass in California have shown varied response to **Quinclorac 75 DF Select**. If control failure occurs following a full or split application, DO NOT reapply **Quinclorac 75 DF Select**. Change to a herbicide with a different mode of action.

⁵ **NOT FOR USE** in California.



◀ See label for complete application guidelines and restrictions.

©2023 Albaugh, LLC. Albaugh and the A logo are registered trademarks and Quinclorac 75DF Select is a trademark of Albaugh, LLC. Drive is a registered trademark of BASF. Always read and follow label restrictions, precautions, and directions for use. This product may not be registered for sale or use in all states or counties. Please check with your state agency responsible for pesticide registration to ensure registration status.