

TAQUET[®]

**SYSTEMIC HERBICIDE FOR BROADLEAF CONTROL
PRE-PLANT AND POST-HARVEST.**

Taquet[®] is a water dispersible granule that is used for selective postemergence weed control in wheat (including durum), barley, oats, triticale, post-harvest burndown, fallow and pre-plant burndown weed control. Best control is obtained when applied to young, actively growing weeds.

Apply Taquet at 0.17 to 0.33 oz/A. Two applications of this product may be made per season, provided the total amount applied does not exceed 0.33 oz/A.

PRODUCT FEATURES

- Systemic activity burns the weed completely to the roots, not allowing for any regrowth.
- Control yield-robbing broadleaf weeds while allowing for maximum cropping flexibility.
- Flexibility as a pre-plant burndown herbicide for sugar beets, winter rape and canola.
- Can also be used pre-plant or at-planting burndown in cotton, field corn, grain sorghum, rice and soybeans.
- Offers powerful control of troublesome weeds including Canada thistle, dandelion, and narrowleaf hawksbeard.
- Can be used as part of an IPM program which can include biological, cultural and genetic practices aimed at preventing economic pest damage.

See label for complete application guidelines and restrictions.



ACTIVE INGREDIENT

75% tribenuron-methyl

HRAC GROUP

2

FORMULATION

Water dispersible granules

REI

12 hours

COMPARE TO

Express[®]

MODE OF ACTION

ALS inhibitor

SIGNAL WORD

Caution

RESTRICTED USE

No

PACKAGING

8 x 10 oz bottles (320 oz cases)

EPA REG NUMBER

83100-31-83979

TAQUET®

FREQUENTLY ASKED QUESTIONS

When should Taquet be applied to wheat, barley, triticale and oats?

- **Wheat, barley, triticale:** Apply Taquet after the crop is in the 2-leaf stage, but before the flag leaf is visible. Do not harvest within 45 days of the last application. Taquet has little-to-no soil activity, so it controls only those weeds that have germinated. For best results, annual broadleaf weeds should be past the cotyledon stage, actively growing and less than 4 inches tall or wide. Several hours of dry weather are needed to allow sufficient absorption by weed foliage.
- **Oats:** For spring oats, make applications after the crop is in the 3-leaf stage, but before jointing. Do not use on Ogle, Porter or Premier varieties as crop injury can occur.
- *For fallow applications, refer to label pages 6-7.*

Can I tank mix Taquet with other herbicides?

- Yes. Taquet may be tank mixed with other suitable registered herbicides to control weeds listed as partially controlled, weeds resistant to Taquet, or weeds not controlled under WEEDS CONTROLLED OR PARTIALLY CONTROLLED sections of the label.
- *Refer to label pages 13-20 for specific tank mix instructions and specific weed problems.*

Can I use Taquet for preplant or at-plant burndown?

- Taquet can be used as a preplant burndown treatment on wheat (including durum), barley or triticale fields to control emerged weeds prior to, or just shortly after planting (prior to emergence). Make applications when the majority of weeds have emerged and are actively growing.
- Cotton, corn, soybeans, and grain sorghum may be planted 14 days after application.
- Sugarbeets, winter rape, and canola can be planted 60 days after application.
- Any other crop may be planted 45 days after application.
- On cotton, field corn, grain sorghum, rice and soybean fields, Taquet may be applied for burndown of emerged weeds before planting or at planting in combination with other suitable registered herbicides. *Refer to label page 7 for specific instructions and use rates.*



◀ See label for complete application guidelines and restrictions.

©2023 Albaugh, LLC. All rights reserved. Albaugh, the Albaugh logo and Taquet are registered trademarks of Albaugh, LLC. Express is a registered trademark of FMC Corporation. Always read and follow label restrictions, precautions and directions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.