

 **FUNGICIDE**

# PROZIO™ SC

**BROAD-SPECTRUM SYSTEMIC FUNGICIDE FOR THE CONTROL OF SPECIFIED DISEASES IN CEREALS, PEANUTS, AND OTHER LABELED CROPS.**



Prozio™ SC is a broad-spectrum systemic fungicide for the control of *Ascomycetes*, *Basidiomycetes* and *Deuteromycetes* diseases in barley, buckwheat, bushberry subgroup, low growing berry subgroup (except strawberry), corn, cotton, cucurbit vegetables, dry shelled pea and bean crop subgroup, millet, oats, peanut, rapeseed subgroup 20A (including canola and *Brassica carinata*), rice, rye, soybean, sugar beets, triticale, wheat; conifer and hardwood nursery seeds and seedlings. Under conditions conducive to extended infection periods or high disease pressure, another registered fungicide may be needed once this product's maximum application rates have been reached. Under these conditions use another fungicide registered for the crop/disease.

## PRODUCT FEATURES

- Advanced triazole technology for powerful protection against a broad range of early season diseases, including *Fusarium*, *Rhizoctonia*, leaf spots, and stem rots.
- Flexible application, in-furrow, at-planting, or in a banded postemergence application to control early season diseases, or by foliar application for management of in-season disease pressure.

See label for complete application guidelines and restrictions.

**ACTIVE INGREDIENT**  
4.0 lbs/gal prothioconazole

**FRAC GROUP**  
3

**FORMULATION**  
Suspension concentrate

**PHI/REI**  
Refer to label

**COMPARE TO**  
Proline® 480 SC

**MODE OF ACTION**  
Demethylation inhibitor/triazoles

**SIGNAL WORD**  
Caution

**RESTRICTED USE**  
No

**PACKAGING**  
2 x 2.5 gal, NR 265

**EPA REG NUMBER**  
45002-56 | AD122122

# PROZIO™ SC

## FREQUENTLY ASKED QUESTIONS

### How should I use Prozio SC on peanuts for best results?

- For in-furrow and banded applications to control Sclerotium rot, white mold, southern blight and other labeled diseases, apply 5.7 fl oz per acre (0.4 fl oz per 1000 row feet if on 36-inch row spacing) in the furrow at planting. Prozio SC may also be applied in a 4- to 6- inch band over the row at or near emergence. Albaugh, LLC recommends a minimum application volume of 20 gpa.
- Refer to label for use instructions for control of soil-borne diseases and for foliar applications.

### How should I use Prozio SC on sugar beets for best results?

- For in-furrow and banded applications for control of *Rhizoctonia* stem canker, root rot, and crown rot, apply 5.7 fl oz/A in the furrow at planting. Refer to label for use directions and rate chart for various row spacing under the SOILBORNE/SEEDLING DISEASE CONTROL section.
- For soil-borne disease control, apply Prozio SC in a seven-inch band at the 4-leaf to row closure growth stage.
- For foliar applications for control of *Cercospora* leaf spot and powdery mildew, apply 5.0-5.7 fl oz/A at the first sign of disease. Use the higher use rate and shorter intervals when conditions are favorable for severe disease pressure and/or when growing less disease resistant varieties.
- Apply up to three (3) applications of Prozio SC per year including all soil and foliar applications.
- Repeat applications as needed using a 14- to 21-day spray interval depending on disease pressure. Use a 14-day spray interval under normal-to-heavy disease pressure and a 21-day spray interval under light disease pressure.
- To optimize disease control, the lowest labeled rate of a spray surfactant may be tank mixed with Prozio SC.
- Refer to label for additional use restrictions.



◀ See label for complete application guidelines and restrictions.

©2024 Albaugh, LLC. All rights reserved. Albaugh and the Albaugh logo are registered trademarks, and Prozio is a trademark of Albaugh, LLC. Proline is a registered trademark of Bayer CropScience. Always read and follow label directions, precautions and restrictions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.