



Premier Supplier of off-patent Crop Protection Products

SUPPLEMENTAL LABELING

(Not for use in California)

READ THE ENTIRE LABEL FOR TRICLOPYR 4E BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABEL

When using TRICLOPYR 4E as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling.

This supplemental labeling must be in the possession of the user at the time of pesticide application.

TRICLOPYR 4E

Specialty Herbicide

APPLICATION FOR ORNAMENTAL TURF

KEEP OUT OF REACH OF CHILDREN.

CAUTION

In case of an emergency involving this product, Call CHEMTREC toll free at 1-800-424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TRICLOPYR 4E may injure certain turfgrass species. Do not apply to bahiagrass, bentgrass, bermudagrass, centipedegrass, St. Augustine grass, or zoysiagrass, unless turf injury can be tolerated.

Established grasses are tolerant to this product, but newly seeded grasses may be injured until well established as indicated by tillering, development of a secondary root system and vigorous growth. Do not reseed treated areas for a minimum of three weeks after treatment.

Do not apply TRICLOPYR 4E to golf course greens.

Do not apply this product through any type of irrigation system.

For spot treatments, do not apply more than 2 qts. of TRICLOPYR 4E per acre in a single application.

Foliar sprays should be applied during warm weather, from early spring through fall, when weeds are actively growing. Broadleaf weeds germinate at different times. Only emerged weeds present at the time of application will be controlled. Newly seeded turf should be mowed 2 or 3 times before being treated.

When making applications to mature plants, hard- to-control species, or during drought conditions, use higher rates. Application under drought conditions may provide less than desirable results. Use low pressure sprays to minimize spray drift. Do not water for 24 hours after application.

Mixing Instructions

When TRICLOPYR 4E is mixed with water it forms an emulsion (not a solution) and separation may occur unless the spray mixture is agitated continuously.

Add about one-half the required amount of clean water to the spray tank. Start agitation and add the specified amount of TRICLOPYR 4E. Provide moderate agitation while completing the addition of water and during application.

Reseeding Note: Do not reseed for 3 weeks after application, (This precaution does not apply when bermudagrass turf is overseeded with perennial ryegrass at a minimum reseeding of 400 lbs. per acre.)

Broadcast Treatment of Ornamental Turf

Apply ½ to 1 quart per acre of TRICLOPYR 4E in enough water to provide uniform coverage of the target area to control actively growing broadleaf weeds growing in perennial bluegrass, perennial ryegrass, or tall fescue. Do not use on other turfgrass species unless injury can be tolerated. To minimize turf injury, do not treat if turf is under heat-or drought-stress and make repeat applications at least 4 weeks apart.

Tank Mixing: To improve the spectrum of activity, TRICLOPYR 4E may be tank mixed at a rate of ½ to 1 pint per acre with specified rates of low volatile amine or ester formulations of 2,4-D, MCPP, or other labeled postemergence broadleaf herbicides. Refer to tank mix product labels for specific use directions, precautions, and limitations before use.

Spot Treatment of Ornamental Turf

Mix 3/8 to ¾ ounces of TRICLOPYR 4E per 1000 square feet in enough water to provide uniform coverage of the target area and apply at any time broadleaf weeds are susceptible.

Note: Do not apply more than 2 quarts per acre or 1.5 ounces per 1000 square feet of TRICLOPYR 4E in a single application.

Control of Kikuyugrass - Apply TRICLOPYR 4E at a rate of ½ to 1 quart per acre. Three to four additional applications at 4 to 6 week intervals may be required to achieve control of kikuyugrass.

Suppression of Bermudagrass - Apply TRICLOPYR 4E at the rate of 1 quart per acre.

Three to four additional applications at 4 week intervals will be required to give adequate suppression of bermudagrass and allow fescue or other desired turfgrass species to dominate. To improve suppression and control of bermudagrass, 1 quart per acre of TRICLOPYR 4E may be tank mixed with a post-emergence grass herbicide registered for this use pattern. Three to four additional applications of this tank mix at 4 week intervals should be made to achieve control. Reseeding following application will accelerate the transition to cool season turf (see Reseeding Note above).

Annual and Perennial Broadleaf Weeds Controlled by TRICLOPYR 4E

Black medic	Curly dock	Matchweed	Sulfur Cinquefoil (2)
Bull Thistle	Dandelion	Mustard	Sweet Clover
Burdock	Dogfennel	Oxalis	Tropical Soda Apple (3)
Canada Thistle	Field Bindweed	Plantain	Vetch
Chicory	Goldenrod	Puple Loosestrife	Wild Carrot (Queen Anne's Lace)
Cinquefoil	Ground Ivy	Ragweed	Wild Lettuce
Clover	Lambsquarters	Sericea Lespedeza (1)	Wild Violet
Creeping Beggarweed	Lespedeza	Smartweed	Yarrow

1. Sericea lespedeza: Apply 1 to 2 pints of TRICLOPYR 4E per acre. For best results, apply after maximum foliage development in the late spring to early summer, but prior to bloom.
2. Sulfur cinquefoil: Apply 1 to 2 pints of TRICLOPYR 4E per acre. For best results, apply to plants in the rosette stage.
3. Tropical soda apple: When plants reach the first flower stage, apply 2 pints of TRICLOPYR 4E per acre. For best results, apply using ground equipment in a total spray volume of 40 gallons per acre. To provide more complete wetting and coverage of the foliage, an agricultural surfactant may be added at the manufacturer's specified rate. To control sparse plant stands, use spot treatments. For spot treatment use a 1 to 1.6% solution of TRICLOPYR 4E in water (1 to 1 ½ gallons of TRICLOPYR 4E in 100 gallons total spray mixture) and spray the entire plant to completely wet the foliage.

In Florida, control of tropical soda apple may be improved by using the following management practices:

- Mow plants to a height of 3 inches every 50 to 60 days or whenever they reach flowering. Continue mowing on this schedule through April.
- In late May to June (50 to 60 days after the April mowing), apply a broadcast treatment of TRICLOPYR 4E.
- To control any remaining plants or to thin stands of plants that germinate following a broadcast treatment, use spot treatments.

EPA Reg. No. 42750-126
AD111014

Manufactured By;
ALBAUGH, LLC
ANKENY, IA 50021