

VOLTA[®] EXTRA

PRE-PLANT AND POSTEMERGENCE CONTROL OF A WIDE SPECTRUM OF BROADLEAF WEEDS ACROSS A NUMBER OF CROPS.



Volta[®] Extra agricultural herbicide is registered for use for selective postemergence control of certain broadleaf weeds in wheat (including durum), barley, oats, triticale, SU Tolerant canola and fallow.

Best results are obtained when Volta Extra is applied to young, actively growing weeds. It stops growth of susceptible weeds rapidly. However, typical symptoms of dying weeds may not be noticeable for 1-3 weeks after application (2-5 weeks for wild garlic), depending on environmental conditions and weed susceptibility. Warm, moist conditions following treatment promote activity, while cold, dry conditions will delay activity.

PRODUCT FEATURES

- Broad-spectrum postemergence weed control in cereal crops.
- Excellent option for pre-plant burndown for cotton, field corn, grain sorghum, rice, sugarbeets, canola, and other crops.
- Tank mix flexibility with grass and broadleaf herbicides to cover full spectrum of weed control.
- Easy-to-use water soluble formulation with easy tank clean-out.
- Excellent performance on winter annuals and wild garlic.

See label for complete application guidelines and restrictions.

ACTIVE INGREDIENTS

50% thifensulfuron-methyl
25% tribenuron-methyl

HRAC GROUP

2

FORMULATION

Dry flowable granule

REI

12 hours

COMPARE TO

Harmony[®] Extra

MODE OF ACTION

ALS inhibitor

SIGNAL WORD

Caution

RESTRICTED USE

No

PACKAGING

8 x 10 oz bottles (320 oz cases)

EPA REG NUMBER

83100-24-83979

VOLTA[®] EXTRA

FREQUENTLY ASKED QUESTIONS

When should I apply Volta Extra on wheat, barley, triticale and oats for best results?

- On wheat (including durum), barley, triticale and winter oats, make applications after the crop is in the 2-3-leaf stage, but before the flag leaf is visible.
- On spring oats, make applications after the crop is in three-leaf stage, but before jointing. Do not use on Ogle, Porter or Premier varieties as crop injury can occur.
- Volta Extra has little to no soil activity and controls only those weeds that have germinated. Annual broadleaf weeds should be past the cotyledon stage, actively growing and less than 4" tall or wide. Wild garlic should be less than 12" tall with 2-4" of new growth.

How should Volta Extra be used to address specific weed problems in cereal crops?

- Volta Extra may be used to control Canada thistle, common cocklebur, common ragweed, lanceleaf sage, wild buckwheat, kochia, Russian thistle, vetch and other tough-to-control weeds in combination with other labeled herbicides.
- *Refer to label pages 10-17 for tank mix options and addressing specific weed problems in cereal crops*



◀ See label for complete application guidelines and restrictions.

©2024 Albaugh, LLC. All rights reserved. Albaugh, the Albaugh logo and Volta are registered trademarks of Albaugh, LLC. Harmony is a registered trademark of FMC Corporation. Always read and follow label restrictions, precautions and directions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.